

Extension of the known range of the snapping shrimp *Alpheus christofferseni* Anker, Hurt and Knowlton, 2007 (Caridea: Alpheidae)

Gabriel Lucas Bochini¹ 

Rodrigo Guéron¹ 

Mariana Terossi² 

Alexandre Oliveira Almeida¹ 

1 Federal University of Pernambuco (UFPE), Biosciences Center, Department of Zoology, Laboratory of Crustacean Biology. Recife, Pernambuco, Brazil.

GLB E-mail: gabriel.bochini@gmail.com

RG E-mail: rggueron@gmail.com

AOA E-mail: aoalmeida.ufpe@gmail.com

2 Federal University of Rio Grande do Sul (UFRGS), Institute of Biosciences, Department of Zoology, Laboratory of Carcinology. Porto Alegre, Rio Grande do Sul, Brazil.

MT E-mail: mterossirm@gmail.com

ZOOBANK: <http://zoobank.org/urn:lsid:zoobank.org:pub:00626966-04C0-4DAE-A412-57C6C352326E>

ABSTRACT

Alpheus christofferseni Anker, Hurt and Knowlton, 2007 was described based on four specimens from Atol das Rocas, northeastern Brazil, and one specimen from Bocas del Toro, the Caribbean coast of Panama. Here, we report the collection of two specimens from Pernambuco, Brazil (~8°41'S), thus increasing the knowledge on species distribution along the Brazilian coast by approximately five latitudinal degrees south of its type locality (Atol das Rocas, Rio Grande do Norte, ~3°51'S). We also report a substantial increase in the known bathymetric distribution from shallow intertidal to approximately 51 m. Notes on morphological variation are provided based on the new material, and the first sequence of the 16S ribosomal subunit gene for the Brazilian coast is provided and compared with congeners.

KEYWORDS

Bathymetric range, Brazilian diversity, Decapoda, distribution, new record

Editor-in-chief
Christopher Tudge

Associate Editor:
Antônio L. Castilho

Corresponding Author
Gabriel Lucas Bochini
gabriel.bochini@gmail.com

Submitted 14 March 2022
Accepted 13 September 2022
Published 27 March 2023

DOI 10.1590/2358-2936e2023001



All content of the journal, except where identified, is licensed under a Creative Commons attribution-type BY.

Nauplius, 31: e2023001

INTRODUCTION

Alpheus christofferseni Anker, Hurt and Knowlton, 2007 has been reported from Atol das Rocas, Brazil (type locality) and from the Atlantic coast of Panama, within burrows in very shallow waters (up to 1.5 m), occasionally associated with echiuran worms (Anker et al., 2007; De Grave and Anker, 2017). Herein, we report two specimens of *A. christofferseni* collected on the coast of Pernambuco State, northeastern Brazil, extending the southernmost known distribution for this shrimp in the western Atlantic by approximately five latitudinal degrees. Additionally, we report a substantial increase in the known bathymetric distribution of the species and the first sequence of the 16S ribosomal subunit gene for one specimen from the Brazilian coast, which is compared with sequences of other *Alpheus* Fabricius, 1798 species and with the sequence of *A. christofferseni* from Panama. We provide drawings illustrating the diagnostic characters and report some morphological variation observed between newly collected material and the description of *A. christofferseni*.

MATERIAL AND METHODS

We collected the first specimen (female) in one of several dredgings carried out off Recife (08°13'25.4"S 34°37'43.2"W, Fig. 1A) in 2018, at a depth of 51 m, with the aim to survey crustacean fauna associated with substrate of biogenic origin (e.g., calcareous algae, and dead coral). Calcareous algae and dead corals were brought to the laboratory, and the endolithic fauna was carefully extracted with a hammer and chisel, via substrate fragmentation. The single specimen obtained was anesthetized on ice and fixed in 70% ethanol for later identification.

We collected the second specimen (male) in 2019, at Praia dos Carneiros, Tamandaré (08°41'32"S 35°04'25"W, Fig. 1A), on the reef near the mouth of the Formoso River. The reefs of Tamandaré are a set of beach rocks approximately 1 km offshore from the beach. The Tamandaré reef complex is part of the Environmental Protection Area "Costa dos Corais", the largest Federal Marine conservation area of Brazil (Maida and Ferreira, 2004). The specimen

was collected by using artificial refuge structures (ARS) as described by Bochini et al. (2020).

The female specimen was deposited in the Museum of Oceanography "Prof. Petrônio Alves Coelho" of the Federal University of Pernambuco, Brazil (MOUFPE) and the male was deposited in the Coleção de Crustáceos do Departamento de Zoologia da Universidade Federal do Rio Grande do Sul, Brazil (DZ/UFRGS). The male specimen was used for drawings made under a dissecting stereomicroscope equipped with a camera lucida and for genetic data. Drawings of the specimen were made by using Adobe Illustrator CS6, according to the methods described by Coleman (2003; 2009). The carapace length (CL, in mm – from the postorbital angle to the posterior margin of the carapace) was measured using a stereomicroscope equipped with a camera lucida and image capture system.

DNA extraction from the male specimen, PCR amplification with specific primers for a region of the 16S ribosomal subunit gene (~550 base pairs), amplicon cleanup, and sequencing were conducted following the laboratory protocols described by Soledade et al. (2019). The obtained sequence and some sequences from the GenBank were used for analysis: one sequence from *A. christofferseni* from Panama, six sequences from species morphologically close to *A. christofferseni* belonging to the *brevirostris* group [*Alpheus barbatus* Coutière, 1897, *Alpheus bellulus* Miya and Miyake, 1969, *Alpheus floridanus* Kingsley, 1878, *Alpheus glaber* (Olivi, 1792) and *Alpheus aff. macroskeles* Alcock and Anderson, 1899] and two sequences used as the outgroup (Fig. 1B). Other 16S gene sequences from closely related species were not available in GenBank. Sequences were aligned by using the MUSCLE software (Edgar, 2004) with default parameters, and Maximum Likelihood performed using RAxML (Stamatakis, 2014) on the platform Cyberinfrastructure for Phylogenetic Research (Miller et al., 2010). The topology consistency was measured by the bootstrap method, and the phylogenetic trees were viewed and edited in MEGA X (Kumar et al., 2018). Only support values above 75% are shown. A genetic distance (p distance) was calculated in MEGAX (Kumar et al., 2018). The sequence obtained in this study was deposited in GenBank (Fig. 1B).




Figure 1. *Alpheus christofferseni* Anker, Hurt and Knowlton, 2007. **A**, Presently known distribution, star represents the type locality, blue and red circles represent the new occurrences, and green circle represents specimen previously reported from Panama. **B**, Phylogram of *A. christofferseni* (bold) and some congeners morphologically similar, using the Maximum Likelihood analysis of the 16S ribosomal subunit gene. Numbers are support values for 960 bootstraps; values < 75% were not included. Atl: Atlantic; PE: Pernambuco; SP: São Paulo. References: A, Chow et al. (2021); B, This study; C, Roura et al. (2018); D, Wang et al. (2020); E, Liao et al. (2019); F, Bracken-Grisson et al. (2014); G, Almeida et al. (2018).

SYSTEMATICS

Family Alpheidae Rafinesque, 1815

Genus *Alpheus* Fabricius, 1798

Alpheus christofferseni Anker, Hurt and Knowlton, 2007

(Fig. 2)

Alpheus christofferseni Anker, Hurt and Knowlton, 2007: 3, figs. 1–6, 11a–c, 12b, c; De Grave and Fransen, 2011: 380; Soledade and Almeida, 2013: 97; De Grave and Anker, 2017: 4.

Material examined. Brazil, Pernambuco: 1 female (CL: 3.8 mm), MOUFPE 20210, of Recife, associated with a biogenic substrate such as calcareous algae, dead coral, and sponges, depth: 51 m, 27.ii.2018, colls. G.L. Bochini, G.O. Soldade and R. Gueron; 1 male (CL: 3.3 mm), DZ/UFRGS 6934 (genetic voucher, GenBank access ON746697), Praia dos Carneiros, Tamandaré, ARS, depth: 3–6 m, 20.viii.2019, colls. G.L. Bochini and G.O. Soledade.

Distribution. The species is found in the Western Atlantic: Panama (Bocas del Toro) and Brazil [Atol das Rocas and Pernambuco (off Recife and Tamandaré)] (Anker et al., 2007; this study).

Habitat. The species is mainly found in burrows on intertidal or shallow seagrass flats (0–1.5 m), associated with echiurans (Anker et al., 2007); reef areas (beach rock) 3–6 m; and substrate of biogenic origin such as calcareous algae, and dead coral, 51 m (this study).

Genetic data. The topology obtained (Fig. 1B) showed a clade formed by the two sequences of *A. christofferseni* and a clear separation of these from

the other species. The intraspecific genetic distance of *A. christofferseni* was 0.9%, much smaller than the divergence of this species from other species of the *brevirostris* group (9.9% - 14.4%).

Remarks. The distribution range of *A. christofferseni* is extended by approximately five degrees south from the type locality (Atol das Rocas, ~3°51'S, Fig. 1A). The current southernmost record of the species is Praia dos Carneiros, Tamandaré, Pernambuco (~8°41'S). Furthermore, the bathymetric distribution increased significantly considering the collection of a specimen at 51 m depth. The specimens from Pernambuco (female and male) resemble the holotype. However, some morphological variations between the two specimens from Pernambuco and the holotype were observed, such as the exopod of the third maxilliped (Fig. 2B, C) reaches the penultimate segment of the endopod (vs. not reaching the penultimate segment of the third maxilliped in the holotype); telson (Fig. 2H, I) with mesial spiniform setae 4.5 times longer than the lateral spiniform setae (vs. mesial spiniform setae twice the size of the lateral spiniform setae in the holotype); third and fourth pereopods (Fig. 2F, G) armed with four spiniform setae along the ventral margin and a pair of distal spiniform setae (vs. five spiniform setae and a pair of distal spiniform setae in the holotype).

ACKNOWLEDGMENTS

We thank Isaac Freitas for helping during collections and Mr. Manuel (Seu Neco) for the vessel and assistance during the scuba diving activities. We thank colleagues from the Laboratory of Crustacean Biology (LBC/UFPE), Guidomar Soledade, Patricia Santos, Ana Carla Costa-Souza, Andressa Cunha, Pedro Paixão, and Renata Lima-Silva who helped with sampling the specimens.




Figure 2. *Alpheus christofferseni* Anker, Hurt and Knowlton, 2007, male from Praia dos Carneiros, Tamandaré, Pernambuco, Brazil (DZ/UFRGS 6934): **A**, Frontal region and cephalic appendages, dorsal view; **B**, third maxilliped, lateral view; **C**, detail of the exopod of the third maxilliped, lateral view; **D**, left major cheliped, lateral view; **E**, right minor cheliped, lateral view; **F**, third pereiopod, lateral view; **G**, fourth pereiopod, lateral view; **H**, telson and uropod, dorsal view; **I**, same, detail of posterior margin. Scale bars: **A**, **D**, **F**, **H**, **I** = 1 mm; **B**, **C**, **E**, **G** = 0.5 mm.

REFERENCES

- Almeida AO; Terossi M; Buranelli RC; Castilho AL; Costa RC; Zara FJ and Mantelatto FL 2018. Checklist of decapods (Crustacea) from the coast of São Paulo State (Brazil) supported by integrative molecular and morphological data: II. Infraorder Caridea: family Alpheidae. *Zootaxa*, 4450(3): 331–358. DOI: 10.11646/zootaxa.4450.3.2
- Alcock A and Anderson ARS 1899. Natural history notes from H.M. Royal Indian marine survey ship Investigator, commander T.H. Heming, R.N., commanding. - Series III, No. 2. An account of the deep-sea Crustacea dredged during the surveying season of 1897-98. *Annals and Magazine of Natural History*, (ser.7), 3: 1-27, 278–292. DOI: 10.1080/00222939908678123
- Anker A; Hurt C and Knowlton N 2007. Three transisthmian snapping shrimps (Crustacea: Decapoda: Alpheidae: *Alpheus*) associated with innkeeper worms (Echiura: Thalassematidae) in Panama. *Zootaxa*, 1626: 1–23. DOI: 10.11646/zootaxa.1626.1.1
- Bochini GL; Cunha MA; Terossi M and Almeida AO 2020. *Chicosciencea*, a new genus of the resurrected family Macromaxillocarididae and a worldwide list of Stenopodidea. *Journal of Crustacean Biology*, 40(6): 704–714. DOI: 10.1093/jcbiol/ruaa064
- Bracken-Grissom HD; Robles R and Felder DL 2014. Molecular phylogenetics of American snapping shrimps allied to *Alpheus floridanus* Kingsley, 1878 (Crustacea: Decapoda: Alpheidae). *Zootaxa*, 3895(4): 492–502. DOI: 10.11646/zootaxa.3895.4.2
- Chow LH; De Grave S; Anker A; Poon KKY; Ma KY; Chu KH; Chan T-Y and Tsang LM 2021. Distinct suites of pre- and post-adaptations indicate independent evolutionary pathways of snapping claws in the shrimp family Alpheidae (Decapoda: Caridea). *Evolution*, 75: 2898–2910. DOI: 10.1111/evo.14351
- Coleman CO 2003. “Digital inking”: How to make perfect line drawings on computers. *Organisms Diversity and Evolution*, 3, Electr. Suppl. 14: 1–14. DOI: 10.1078/1439-6092-00081
- Coleman CO 2009. Drawing setae the digital way. *Zoosystematics and Evolution*, 85: 305–310. DOI: 10.1002/zoos.200900008
- Coutière H 1897. Note sur quelques Alpheïdes nouveaux ou peu connus rapportés de Djibouti (Afrique orientale). *Bulletin du Muséum d’Histoire naturelle*, 3: 233–236. DOI: 10.5962/bhl.part.19259
- De Grave S and Anker A 2017. An annotated checklist of marine caridean and stenopodidean shrimps (Malacostraca: Decapoda) of the Caribbean coast of Panama. *Nauplius*, 25 (e2017015): 1–40. DOI: 10.1590/2358-2936e2017015
- De Grave S and Fransen CHJM 2011. Carideorum Catalogus: The recent species of the dendrobranchiate, stenopodidean, procarididean and caridean shrimps (Crustacea: Decapoda). *Zoologische Mededelingen*, 85(9): 195–588. <https://repository.naturalis.nl/pub/403473>
- Edgar RC 2004. MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Research*, 32: 1792–1797. DOI: 10.1093/nar/gkh340
- Fabricius JC 1798. Supplementum Entomologiae Systematicae: 1–572. Hafniae, Proft et Storch. <https://www.biodiversitylibrary.org/item/132638-page/5/mode/1up>
- Kingsley JS 1878. A synopsis of the North American species of the genus *Alpheus*. *Bulletin of the United States Geological and Geographical Survey*, 4: 189–199. <https://www.marinespecies.org/aphia.php?p=sourceget&id=155546>
- Kumar K; Stecher G; Li M; Knyaz C and Tamura K 2018. MEGA X: Molecular evolutionary genetics analysis across computing platforms. *Molecular Biology and Evolution*, 35: 1547–1549. DOI: 10.1093/molbev/msy096
- Liao Y; Ma KY; De Grave S; Komai T; Chan TY and Chu K 2019. Systematic analysis of the caridean shrimp superfamily Pandaloidea (Crustacea: Decapoda) based on molecular and morphological evidence. *Molecular phylogenetics and evolution*, 134: 200–210. DOI: 10.1016/j.ympev.2019.02.006
- Maida M and Ferreira BP 2004. Recifes de coral brasileiros. p. 617–640. In: Eskinazi-Leça, E; Neumann-Leitão S; Ferreira da Costa M (Orgs.), Oceanografia: um cenário tropical. Recife, Bagaço.
- Miller MA; Pfeiffer W and Schwartz T 2010. Creating the CIPRES Science Gateway for inference of large phylogenetic trees. p. 1–8. In: 2010 Gateway Computing Environments Workshop (GCE), New Orleans, LA, USA, 14 November 2010. IEEE. DOI: 10.1109/GCE.2010.5676129
- Miya Y and Miyake S 1969. Description of *Alpheus bellulus* sp. nov. associated with gobies from Japan (Crustacea, Decapoda, Alpheidae). *Publications from the Seto Marine Biological Laboratory*, 16: 307–314. DOI:10.5134/175556
- Olivi G 1792. Zoologia Adriatica, ossia catalogo ragionato degli animali del golfo e della lagune di Venezia. Bassano, G. Remondini e fl., ix+334+xxxii 366p, 9 pls. <http://www.biodiversitylibrary.org/item/123881>
- Rafinesque CS 1815 Analyse de la Nature, ou Tableau de l’Univers et des Corps Organisés. Palermo, L’Imprimerie de Jean Barravecchia. 224 p. DOI: 10.5962/bhl.title.106607
- Roura Á; Strugnell JM; Guerra Á; González ÁF and Richardson AJ 2018. Small copepods could channel missing carbon through metazoan predation. *Ecology and Evolution*, 8: 10868–10878. DOI: 10.1002/ece3.4546

- Soledade GO and Almeida AO 2013. Snapping shrimps of the genus *Alpheus* Fabricius, 1798 from Brazil (Caridea: Alpheidae): updated checklist and key for identification. *Nauplius*, 21(1): 89–122. DOI: 10.1590/S0104-64972013000100010
- Soledade GO; Terossi M; Scioli JA; Mantelatto FL and Almeida AO 2019. A new western Atlantic snapping shrimp of the *Alpheus macrocheles* group (Caridea, Alpheidae) revealed by morphological, molecular and color data. *European Journal of Taxonomy*, 581: 1–27. DOI: 10.5852/ejt.2019.581
- Stamatakis A 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics*, 30: 1312–1313. DOI: 10.1093/bioinformatics/btu033
- Wang Q; Wang Z; Tang D; Xu X; Tao Y; Ji C and Wang Z 2020. Characterization and comparison of the mitochondrial genomes from two Alpheidae species and insights into the phylogeny of Caridea. *Genomics*, 112(1): 65–70. DOI: 10.1016/j.ygeno.2019.08.013

ADDITIONAL INFORMATION AND DECLARATIONS

Author contributions

Conceptualization and Design: GLB, RG, AOA; Performed research: GLB, RG, AOA; Acquisition of data: GLB, RG, MT, AOA; Analysis and interpretation of data: GLB, RG, MT, AOA; Preparation of figures/tables/maps: GLB, MT; Writing – original draft: GLB; Writing – critical review & editing: GLB, RG, MT, AOA.

Consent for publication

All authors declare that they have reviewed the content of the manuscript and gave their consent to submit the document.

Funding and grant disclosures

Fundação de Amparo à Ciência e Tecnologia do Estado de Pernambuco (FACEPE), Grant/ Award Number: APQ-0196-2.04/16, APQ-0572-2.04/15, and IBPG-

0084-2.05/17; Conselho de Desenvolvimento Científico e Tecnológico (CNPq), Grant/ Award Number: DCR-300067/2018-6, Universal 421963/2016-6, PQ 304235/2019-9, Universal 421193/2018-2, and PQ 311340/2021-0; Coordenação de Aperfeiçoamento de Pessoal de Nível Superior – Brazil (CAPES), Grant/ Award Number: 88887.146252/2017-00.

Conflicts of interest

The authors have no conflicts of interest to declare.

Study permits

Field collection and transportation of specimens were made under the SISBIO permit number 58697-1 issued to G. L. Bochini

Data availability statement

All data generated and analyzed during this study are presented in this article.