



Felipe Turchetto<sup>1a+</sup>, Maristela Machado Araujo<sup>1b</sup>, Suzane Bevilacqua Marcuzzo<sup>1c</sup>, Álvaro Luís Pasquetti Berghetti<sup>1d</sup>, Danieli Guarienti Rorato<sup>1e</sup>, Adriana Maria Griebeler<sup>1f</sup>, Felipe Manzoni Barbosa<sup>1g</sup>

## IMPACT OF SEEDLING REMOVAL ON NATURAL REGENERATION IN THE SOUTHERN ATLANTIC FOREST REMNANT

TURCHETTO, F.; ARAUJO, M. M.; MARCUZZO, S. B.; BERGHETTI, A. L. P.; RORATO, D. G.; GRIEBELER, A. M.; BARBOSA, F. M. Impact of seedling removal on natural regeneration in the southern atlantic forest remnant. *CERNE*, v. 24, n. 2, p. 98-105, 2018.

### HIGHLIGHTS

The seedling bank can be used as a source of propagules.

Forest ecosystems have resilience capacity under low anthropic intervention rates.

Forest species present different levels of tolerance to the transplantation of seedlings.

Species with abundant natural regeneration can be removed from the forest.

### ABSTRACT

The use of a forest seedling bank has been recommended as an alternative to increase species richness in forest nurseries, as well as to produce seedlings of species that are difficult to propagate, especially those that belong to the late secondary and climax successional groups, which are not as commercially available. However, little is known about the impact of this method on forest dynamics. Thus, the present study aimed to examine the resilience and dynamics of a seedling bank in a remnant of a subtropical seasonal forest belonging to the Atlantic Forest Biome when subjected to different intensities of seedling removal. The experiment was conducted in a random block design in a factorial scheme (5 × 4), with treatments composed of five intensities of removal of individuals from the seedling bank and the four seasons. The treatments were distributed into 18 blocks and the experimental units were composed of 1 m × 2.5 m plots. The resilience of the seedling community was assessed by examining effects of the five removal intensities. The dynamics between the evaluation periods within each treatment were verified by comparing the number of species and seedlings present before the treatments with those in the other evaluation periods. After one year, we found that tree-shrub vegetation had a partial capacity for restoration after withdrawal of individuals from the seedling bank. Our results show that the impact on the regeneration community can absorb the effects of up to 25% seedling removal. The technique of seedling transplantation may be recommended for species that have abundant regeneration, such as *Actinostemon concolor*, *Eugenia rostrifolia*, *Trichilia claussenii* and *Nectandra megapota mica*.

**Keywords:**  
Forest dynamics  
Restoration of altered areas  
Seedlings bank

**Historic:**  
Received 05/02/2018  
Accepted 23/05/2018

**+Correspondence:**  
turchetto.felipe@gmail.com

**DOI:**  
10.1590/01047760201824022517

<sup>1</sup> Federal University of Santa Maria, Santa Maria, Brazil. ORCID: 0000-0002-2526-2440<sup>a</sup>, 0000-0003-3751-8754<sup>b</sup>, 0000-0003-1275-6502<sup>c</sup>, 0000-0001-8020-3200<sup>d</sup>, 0000-0003-2751-7336<sup>e</sup>, 0000-0002-9891-5959<sup>f</sup>, 0000-0003-0833-9546<sup>g</sup>

## INTRODUCTION

In the present scenario of a high deforestation rate in tropical forests (Hansen et al., 2013), it is critical to develop strategies to conserve or at least minimize the loss of biodiversity (Rodrigues et al., 2011), as well as to protect soil structure (Rodrigues et al., 2011; Rocha Junior et al., 2013).

Forest restoration approaches vary depending on the level of degradation, residual vegetation, and the desired results (Chazdon, 2008). Environmental analysis is required to define the most appropriate reforestation technique (Chazdon, 2008). In highly degraded environments with impoverished soils and with no potential for natural resilience, the planting of highly diverse forest species (Rodrigues et al., 2009) can speed up the restoration process, providing benefits such as soil and water conservation, carbon sequestration, and other ecosystem services (Rodrigues et al., 2009; Gazell et al., 2012; Ferez et al., 2015).

However, there is a low supply of native forest species available for commercialization in nurseries, as can be seen in the low species diversity of restoration plantations (Viani and Rodrigues, 2008). In addition, two-thirds of these species belong to the early stages of secondary succession (Barbosa et al., 2003) and may reduce the ecological viability of forests in the process of restoration, as is the case for areas in fragmented landscapes (Rodrigues et al., 2011).

Under these circumstances, some studies have shown that transplantation of naturally regenerated seedlings could be a strategy for the production of seedlings in nurseries (Calegari et al., 2011; Viani et al., 2012; Turchetto et al., 2016). The main advantage of this technique is the production of seedlings adapted to their bioclimatic regions, mainly of pioneering species that are not available in nurseries.

The formation of a seedling bank is a strategy that can be used for species of plants that germinate under the canopy (Whitmore, 1989). The process of recruitment and establishment in tropical forests depends on a number of factors, including moisture, nutrients, predation of herbivorous seeds and seedlings, and the occurrence of pathogens (Molofsky and Fisher, 1993; Brearley et al., 2003). Therefore, of the total number of germinated seeds that constitute the seedling bank, approximately 10% survive beyond the juvenile phase. Thus, utilization of these seedlings which would normally be eliminated can be greatly beneficial to restoration practices (Carrington, 2014).

However, according to Viani and Rodrigues (2008), any anthropic action, modification, or management practice in natural areas, in this case specifically related to

the seedling community, requires previous investigation in order to avoid irreparable damage to the diversity, productivity, and connectivity of the population, the community, and the ecosystem as a whole. This information can be obtained by analyzing the restoration capacity of an individual/species/community after exposure to adverse conditions (Higa et al., 2000); these conditions may be related to environmental restrictions or the intensity of anthropic interference, which define the potential for resilience of the community.

In this context, the present study aimed to characterize the resilience and dynamics of the seedling bank of a forest remnant in the extreme south of the Atlantic Forest biome when subjected to seedling transplantation. We sought to answer the following questions: a) Does seedling transplantation impair the dynamics of the community? b) Is there a limit of acceptable removal? and c) Does the impact vary among species?

## MATERIAL AND METHODS

### Study area

The study was carried out in a remnant of the Subtropical Seasonal Forest of approximately 20 ha (29°27'14.71" S and 53°18'17.86" W), in the extreme south of the Atlantic Forest biome, in the central region of the state of Rio Grande do Sul, Brazil. The soil is classified as Neossol Regolithic (EMBRAPA, 2013) and the climate is subtropical, with an average annual precipitation of 1560 mm (Alvares et al., 2013), characterized by well-defined seasons.

### Experimental design

The experiment was conducted as a randomized block design, with five removal intensities from the seedling bank (0%, 25%, 50%, 75%, and 100% removal). The measures were taken in each of the four seasons. The treatments were distributed into 18 blocks (5 m × 5 m) and the experimental units were composed of 1 m × 2.5 m plots.

Plots were distributed so as to avoid interference with natural regeneration during measurements and were therefore arranged with intervals rather than continuously, allowing the evaluator to measure without damaging the vegetation (Turchetto, 2015).

### Data collection

Five evaluations were carried out, the first being the control (a single evaluation), before application of the treatments. Following removal of the seedlings, four

additional evaluations were carried out at 3 (February 2014), 6 (May 2014), 9 (August 2014) and 12 months (November 2014), one for each season—summer, autumn, winter, and spring, respectively.

Data were collected regarding the number of species and seedlings in each plot. Individuals of the tree-shrub category with a height between 5 and 55 cm were considered seedlings and identified according to the APG III classification system (THE ANGIOSPERM PHYLOGENY, 2009). For individuals not identified at the time of sampling, individuals of the same species located outside of the plot were collected.

### Statistical analysis

To evaluate the impact caused by the removal of the seedlings on natural regeneration, a general linear model (GLM) was used to analyze the variance associated with the removal intensities and the evaluation times.

The number of species and seedlings in the control evaluation were compared to those in the other evaluation periods (3, 6, 9 and 12 months) to verify the regeneration dynamics of the seedling community, using the Dunnett test (5% significance level).

## RESULTS

### Impact on the seedling community and regeneration dynamics

Removal of regenerating seedlings at intensities over 50% reduced the number of species and seedlings in each plot ( $p < 0.05$ ). The control plots (without removal) presented the best results (Figure 1).


Recruitment of individuals and species varied over time, with greater recruitment between 6 months (May 2014) and 9 months (August 2014), corresponding to autumn and winter, respectively. The last evaluation, in spring (November/2014), presented the lowest number of both species (Figure 2a) and individuals (Figure 2b).

### Resilience of the seedling community


The number of individuals (Figure 3) and species (Figure 4) did not differ significantly from that in the control for the 25% removal condition, but significant differences were observed for all other treatments.

For 75% and 100% removal, there was over 50% reduction in the number of species and individuals at 12 months compared to the control.


For the most abundant species in the seedling bank (*Actinostemon concolor* (Spreng.) Müll.Arg, *Eugenia rostrifolia* D.Legrand, *Nectandra megapotamica* (Spreng.) Mez and *Trichilia claussenii* C. DC.), we compared the




**FIGURE 1** Effect of different removal intensities on number of individuals (a) and number of species (b) present in the seedling bank in a remnant of a Subtropical Seasonal Forest in the extreme South of the Atlantic Forest Biome. \*Significant 5% probability by GLM test. Vertical bars indicate the confidence interval.



**FIGURE 2** Seasonal variation in number of individuals (a) and number of species (b) present in the seedling bank in a remnant of a subtropical seasonal forest in the extreme South of the Atlantic Forest Biome. Means followed by uppercase letters were compared by the GLM test.



**FIGURE 3** Comparison of the mean number of seedlings per plot between time 0 (control) and the other evaluation periods (3, 6, 9 and 12 months), for each treatment. Witness (a); 25% removal (b); 50% removal (c); 75% removal (d); and 100% removal (e). Means followed by uppercase letters indicate comparison to the control (Dunnnett test 5% probability).




**FIGURE 4** Comparison of the mean number of species per plot between time 0 (control) and the other periods (3, 6, 9 and 12 months), for each treatment. Witness (a); 25% removal (b); 50% removal (c); 75% removal (d); and 100% removal (e). Means followed by uppercase letters indicate comparison to control (Dunnnett test at 5% probability).

number of individuals in each evaluation and removal intensity and found that the species adopted different recruiting strategies to cope with the removal of regenerating individuals (Figure 5).

*A. concolor* was the only species that presented an increase in the number of individuals recruited at 12 months after seedling removal, for all treatments evaluated. However, there was a considerable reduction

of individuals between the 9<sup>th</sup> and 12<sup>th</sup> months (August to November) (Figure 5a).

*E. rostrifolia* presented a considerable reduction in the number of individuals in the 3<sup>rd</sup> month (February 2014) for treatments with 50%, 75%, and 100% removal, and for the control. Additionally, for species *N. megapotamica* and *T. clausenii*, there was a significant reduction in seedlings at removal intensities higher than 75% (Figure 5).



**FIGURE 5** Seasonal dynamics of individuals at different removal intensities (100%; 75%; 50%; 25% and 0%) for *Actinostemon concolor* (a), *Eugenia rostrifolia* (b), *Nectandra megapotamica* (c) and *Trichilia clausenii* (d).

## DISCUSSION

Our results show that the regeneration community can tolerate the removal of up to 25% of regenerating seedlings, while removal intensities over 50% interfere with the resilience of the seedling bank, as the vegetal community was not able to reestablish the species and seedlings initially present. Viani and Rodrigues (2008), investigating the impact on a seedling bank caused by different removal intensities in a semideciduous seasonal forest in the state of São Paulo, found that at 18 months, seedlings and species numbers in plots with 50% removal of individuals did not differ significantly from those in the control (0% removal).

This shows the resilience potential of the environment when the remnant vegetation is not considerably affected. Thus, the results indicate that naturally regenerated seedlings can be used to enrich restoration projects in degraded areas without compromising the regenerative potential of forests.

On the other hand, the removal of more than 50% of the seedlings led to a reduction in the number of individuals and species sampled, indicating a limit on the possible removal of regenerating individuals. In this case, a larger number of species ceased to constitute the seedlings bank, probably due to the need to maintain a “stock” with some species with more difficult dispersion

and/or regeneration. This scenario demonstrates that the impact of the removal of individuals contained in the seedling bank may be significant for some species, especially those with low regeneration density

Comparisons over the different periods of evaluation demonstrated the influence of the season on recruitment of tree-shrub seedlings. There was a greater density of individuals at 6 and 9 months (autumn and winter), with considerable reductions in the density of individuals from the 12<sup>th</sup> month (spring). Other studies of seedling banks in seasonal forests have also found variation over time both in the richness and abundance of regenerating species (McLaren and McDonald, 2003; Ceccon et al., 2004; Venturoli et al., 2011).

The phenology of species in seasonal forests, mechanisms of dormancy (Andeis et al., 2005), germination of many individuals in the seed bank at the end of spring and summer (Viani and Rodrigues, 2008), and seasonality due to supra-annual reproductive characteristics of many tropical tree species, all influence the seedling bank composition over the year.

In this study, increased seedling recruitment at 6 (autumn) and 9 months (winter) may be related to the reproductive characteristics of *A. concolor* (*Iaranjeira-domato*). This species presented the greatest density and was the only species with a greater number of individuals at 3

months (February 2014) than at the initial evaluation, for plots with 100% transplantation of regenerating individuals.

The initial evaluation was carried out in the flowering/fruitlet period for *A. concolor*, which is between September and November (Andreis et al., 2005). In the subsequent evaluations, an increase in the number of emergent individuals is expected due to seed dispersion, since it is an understory species (Scipioni et al., 2013), indicating a seedling bank regeneration strategy.

The decreased number of seedlings at 12 months (November 2014) may be related to the increased temperature at this time of year, since seedlings with a height lower than 55 cm are sensitive to temperature variation. (Huxman et al., 2004), evaluating the efficiency of water utilization in different ecosystems, found that temperature and precipitation interfered in seedling survival and growth. According to Metz et al. (2008), seedlings are more susceptible to seasonal water deficit because they do not have deep enough roots to obtain water from deeper in the soil. The effects of the water deficit on the dry period are greater because of the increased temperatures and intensities of solar radiation, which may lead to plant desiccation and death (Lieberman and Mingguang, 1992; Vieira and Scariot, 2006).

When we analyzed the most abundant species in the seedling bank (*A. concolor*, *E. rostrifolia*, *N. megapotamica*, and *T. clausenii*), different regeneration strategies following seedling bank removal were observed. The variations between treatments and between seasons within each treatment may be related to reproductive and silvicultural characteristics of the species.

According to Janzen (1970), the survival, establishment, and development of seedlings are influenced by several morphological, physiological, and abiotic factors, as well as by biological interactions. The way each species responds to these factors is determined by seedling adaptations, that is, the way each species interacts with the environment and with other organisms (Melo et al., 2004).

*E. rostrifolia* and *A. concolor* presented intraspecific thinning. According to Freckleton and Watkinson (2002), survival rates in most plants decrease as density increases. Moreover, higher temperatures and lack of rain can also contribute to seedling mortality in these species. Because most of the individuals presented a height below 10 cm, they did not possess photosynthetic capacity efficient enough for establishment.

*T. clausenii* and *N. megapotamica* presented a significant reduction in the number of individuals at removal intensities over 75%. *T. clausenii* presents a low germination rate (Backes and Irgang, 2002), while *N.*

*megapotamica* is characterized by abundant regeneration under shade (Backes and Irgang, 2002; Carvalho, 2006). Davide et al. (2003) note that species of the genus *Nectandra* present seeds with short longevity (recalcitrant). Thus, the existence of a seedling bank for these species may constitute a survival strategy, as these remain in the understory until recruitment by higher classes.

According to Viani and Rodrigues (2008), the use of seedling banks of natural areas for seedling production aimed at forest restoration should focus on species with high regeneration density and evident intraspecific thinning, which was the case for *E. rostrifolia* and *A. concolor* in all treatments evaluated.

*T. clausenii* and *N. megapotamica* did not present intraspecific thinning. However, for the treatments up to 50% removal, the number of individuals was similar to that found for the control (0% removal) at 12 months. Of the species studies here, these show good potential for use in forest nurseries, considering that their seedlings are difficult to obtain and produce since they are adapted to regional edaphoclimatic conditions.

## CONCLUSIONS

The remnant studied presented partial capacity for self-regeneration following low-intensity seedling bank removal.

Removal of up to 25% of the seedling bank over a short period of time does not interfere with the dynamics of the most abundant populations in the forest understory and may be a useful strategy to produce seedlings of species presenting difficult propagation.

Seedling removal of up to 25% does not impair the persistence of the communities of the species *A. concolor*, *E. rostrifolia*, *T. clausenii*, and *N. megapotamica*, which present abundant natural regeneration and/or intraspecific thinning. However, more studies with a longer period of evaluation are necessary to more fully understand the interference of seedling removal on vegetal community resilience.

## ACKNOWLEDGEMENTS

This research was supported by the Foundation for Research Support of the Rio Grande do Sul State (FAPERGS) and the Environmental Fund of the Caixa Econômica Federal (AC FSA CAIXA, N°. 015,007 / 2012).

## REFERENCES

- ALVARES, C. A.; STAPE, J. L.; SENTELHAS, P. C.; GONÇALVES, J. L. d. M.; SPAROVEK, G. Köppen's Climate Classification Map for Brazil. *Meteorologische Zeitschrift*, v. 22, n. 6, p. 711-728, 2013.

- ANDREIS, C.; LONGHI, S. J.; BRUN, E. J.; WOJCIECHOWSKI, J. C.; MACHADO, A. A.; VACCARO, S.; CASSAL, C. Z. Estudo Fenológico Em Três Fases Sucessionais de Uma Floresta Estacional Decidual No Município De Santa Tereza, Rs, Brasil. **Revista Árvore**, v. 29, n. 1, p. 55-63, 2005.
- BACKES, P.; IRGANG, B., **Árvores do sul: guia de identificação e reconhecimento ecológico**. Palloti, Porto alegre, 2002, 325p.
- BARBOSA, L. M.; BARBOSA, J. M.; BARBOSA, K. C.; POTOMATI, A.; MARTINS, S. E.; ASPERTI, L. M.; MELO, A. C. G.; CARRASCO, P. G.; CASTANHEIRA, S. A.; PILIACKAS, J. M.; CONTIERI, W. A.; MATTIOLI, D. S.; GUEDES, D. C.; SANTOS JUNIOR, N. A.; SILVA, P. M. S.; PLAZA, A. P. Recuperação florestal com espécies nativas no estado de São Paulo: Pesquisas apontam mudanças Necessárias. **Florestar Estatístico**, v. 6, n. 14, p. 28-30, 2003.
- BREARLEY, F. Q.; PRESS, M. C.; SCHOLE, J. D. Nutrients obtained from leaf litter can improve the growth of dipterocarp seedlings. **New Phytologist**, v. 160, n. 1, p. 101-110, 2003.
- CALEGARI, L.; MARTINS, S. V.; BUSATO, L. C.; SILVA, E.; COUTINHO JUNIOR, R.; GLERIANI, J. M. Produção de mudas de espécies arbóreas nativas em viveiro via resgate de plantas jovens. **Revista Árvore**, v. 35, n. 1, p. 41-50, 2011.
- CARRINGTON, M. E. Seed size and recruitment limitation influence seedling establishment in three tallgrass prairie species. **Plant Ecology**, v. 215, n. 10, p. 1163-1172, 2014.
- CARVALHO, P. E. R. **Espécies Arbóreas Brasileiras**. Embrapa florestas, Colombo, v.2, p. 627, 2006.
- CECCON, E.; SÁNCHEZ, S.; CAMPO, J. Tree seedling dynamics in two abandoned tropical dry forests of differing successional status in Yucatán, Mexico: A Field experiment with N and P fertilization. **Plant Ecology**, v. 170, n. 2, p. 277-285, 2004.
- CHAZDON, R. L. Beyond Deforestation: Restoring Forests and Ecosystem Services on Degraded Lands. **Science**, v. 320, n. 5882, p. 1458-1460, 2008.
- DAVIDE, A. C.; CARVALHO, L. R.; CARVALHO, M. L. M.; GUIMARÃES, R. M. Seed physiology classification of lauraceae forest species in relation to the storage behaviour. **Cerne**, v. 9, n. 1, p. 29-35, 2003.
- EMBRAPA, **Sistema Brasileiro De Classificação De Solos**, 3ª ed. Brasília, Ministério da Agricultura, Pecuária e Abastecimento, p.353, 2013.
- FEREZ, A. P. C.; CAMPOE, O. C.; MENDES, J. C. T.; STAPE, J. L. Silvicultural opportunities for increasing carbon stock in restoration of Atlantic Forests in Brazil. **Forest Ecology and Management**, v. 350, p. 40-45, 2015.
- GAZELL, A. C. F.; RIGHI, C. A.; STAPE, J. L.; CAMPOE, O. C. Tree species richness, does it play a key role on a forest restoration plantation? **Bosque**, v. 33, n. 3, p. 245-248, 2012.
- FRECKLETON, R. P.; WATKINSON, A. R. Are Weed population dynamics chaotic? **Journal of Applied Ecology**, v. 39, n. 5, p. 699-707, 2002.
- HANSEN, M. C.; POTAPOV, P. V.; MOORE, R.; HANCHER, M.; TURUBANOVA, S. A.; TYUKAVINA, A.; THAU, D.; STEHMAN, S. V.; GOETZ, S. J.; LOVELAND, T. R.; KOMMAREDDY, A.; EGOROV, A.; CHINI, L.; JUSTICE, C. O.; TOWNSHEND, J. R. G. High-Resolution Global Maps of 21st-Century Forest Cover Change. **Science**, v. 342, n. 6160, p. 850-853, 2013.
- HIGA, R. C. V.; HIGA, A. R.; TREVISAN, R.; SOUZA, M. V. Resistência e resiliência a geadas em *Eucalyptus Dunnii* Maiden plantados em Campo do Tenente, PR. **Boletim de Pesquisas Florestais**, Colombo, v. 40, p. 67-76, 2000.
- HUXMAN, T. E.; SMITH, M. D.; FAY, P. A.; KNAPP, A. K.; SHAW, M. R.; LOIK, M. E.; SMITH, S. D.; TISSUE, D. T.; ZAK, J. C.; WELTZIN, J. F.; POCKMAN, W. T.; SALA, O. E.; HADDAD, B. M.; HARTE, J.; KOCH, G. W.; SCHWINNING, S.; SMALL, E. E.; WILLIAMS, D. G. Convergence across Biomes to a Common Rain-Use Efficiency. **Nature**, v. 429, n. 6992, p. 651-654, 2004.
- JANZEN, D. H. Herbivores and the number of tree species in tropical forests. **The American Naturalist**, v. 104, n. 940, p. 501-528, 1970.
- LIEBERMAN, D.; MINGGUANG, Li. Seedling recruitment patterns in a tropical dry forest in Ghana. **Journal of Vegetation Science**, v. 3, n. 3, p. 375-382, 1992.
- MCLAREN, K. P.; MCDONALD, M. A. Coppice regrowth in a disturbed tropical dry limestone forest in Jamaica. **Forest Ecology and Management**, v. 180, n. 1-3, p. 99-111, 2003.
- MELO, F. P. L.; AGUIAR NETO, A. V.; SIMABUKURO, E. A.; TABARELLI, M. Recrutamento e estabelecimento de plântulas. In: FERREIRA, A. G.; BORGHETTI, F. **Germinação: do básico ao aplicado**. São Paulo, p. 237-250, 2004.
- METZ, M. R.; COMITA, L. S.; CHEN, Y. Y.; NORDEN, N.; CONDIT, R.; HUBBELL, S. P.; SUN, I. F.; NOOR, N. S. B. M.; WRIGHT, S. J. Temporal and spatial variability in seedling dynamics: a cross-site comparison in four lowland tropical forests. **Journal of Tropical Ecology**, v. 24, n. 1, p. 9-18, 2008.
- MOLOFSKY, J.; FISHER, B. L. Habitat and predation effects on seedling survival and growth in shade-tolerant tropical trees. **Ecology**, v. 74, n. 1, p. 261-265, 1993.
- ROCHA JUNIOR, P. R.; DONAGEMMA, G. K.; ANDRADE, F. V.; PASSOS, R. R.; BALIEIRO, F. C.; MENDONÇA, E. S.; RUIZ, H. A. Can soil organic carbon pools indicate the degradation levels of pastures in the Atlantic Forest Biome? **Journal of Agricultural Science**, v. 6, n. 1, p. 84, 2013.

- RODRIGUES, R. R.; GANDOLFI, S.; NAVE, A. G.; ARONSON, J.; BARRETO, T. E.; VIDAL, C. Y.; BRANCALION, P. H. S. Large-Scale ecological restoration of high-diversity Tropical Forests in Se Brazil. **Forest Ecology and Management**. v. 261, n. 10, p. 1605-1613, 2011.
- RODRIGUES, R. R.; LIMA, R. A. F.; GANDOLFI, S.; NAVE, A. G. On the restoration of high diversity forests: 30 years of experience in the Brazilian Atlantic Forest. **Biological Conservation**. v. 142, n. 6, p. 1242-1251, 2009.
- SCIPIONI, M. C.; GALVÃO, F.; LONGHI, S. J. Composição florística e estratégias de dispersão e regeneração de grupos florísticos em Florestas Estacionais Deciduais no Rio Grande do Sul. **Floresta**. v. 43, n. 2, p. 241-254, 2013.
- THE ANGIOSPERM PHYLOGENY, G. An update of the angiosperm phylogeny group classification for the orders and families of flowering plants: Apg Iii. **Botanical Journal of the Linnean Society**. v. 161, n. 2, p. 105-121, 2009.
- TURCHETTO, F. **Potencial do banco de plântulas como estratégia para restauração florestal no extremo sul do Bioma Mata Atlântica**. 2015 Dissertação. Universidade Federal de Santa Maria, Santa Maria, 135 f. 2015.
- TURCHETTO, F.; ARAUJO, M. M.; TABALDI, L. A.; GRIEBELER, A. M.; RORATO, D. G.; AIMI, S. C.; BERGHETTI, Á. L. P.; GOMES, D. R. Can transplantation of forest seedlings be a strategy to enrich seedling production in plant nurseries? **Forest Ecology and Management**. v. 375, n. 1, p. 96-104, 2016.
- VENTUROLI, F.; FELFILI, J. M.; FAGG, C. W. Avaliação temporal da regeneração natural em uma Floresta Estacional Semidecídua secundária, em Pirenópolis, Goiás. **Revista Árvore**. v. 35, n. 3, p. 473-483, 2011.
- VIANI, R. A. G.; BRANCALION, P. H. S.; RODRIGUES, R. R. Corte foliar e tempo de Transplântio para o uso de plântulas do sub-bosque aa restauração florestal. **Revista Árvore**. V. 36, n. 2, p. 331-339, 2012.
- VIANI, R. A. G.; RODRIGUES, R. R. Impacto da remoção de plântulas sobre a estrutura da comunidade regenerante de Floresta Estacional Semidecidual. **Acta Botanica Brasilica**. v. 22, n. 4, p. 1015-1026, 2008.
- VIEIRA, D. L. M.; SCARIOT, A. Principles of natural regeneration of tropical dry forests for restoration. **Restoration Ecology**. v. 14, n. 1, p. 11-20, 2006.
- WHITMORE, T. C. Canopy gaps and the two major groups of forest trees. **Ecology**. v. 70, n. 3, p. 536-538, 1989.