

Nail psoriasis treated with intralesional methotrexate infiltration*

Artur Antônio Duarte¹
 Getúlio Pisa Carneiro²
 Carolina Malavassi Murari²
 Laura Costa Beber de Jesus²

DOI: <http://dx.doi.org/10.1590/abd1806-4841.20198170>

Dear Editor,

Psoriasis is a chronic inflammatory skin disease characterized by recurrent events. Its prevalence around the world is approximately 1-3%.^{1,2} It's estimated that 80-90% of patients with psoriasis will present with nail disorders throughout their lives.^{2,3} Studies have shown that nail psoriasis is the cause of a significant social problem, often associated with pain.² Different therapeutic options can be found for nail psoriasis, which include topical medication, intralesional treatment, systemic therapy, conventional therapies with immunobiological drugs, as well as non-pharmacological treatments.^{2,3} Methotrexate (MTX) is a folic acid analogue, responsible for inhibiting the synthesis of deoxyribonucleic acid.^{2,3} A literature review shows that intralesional MTX has been successfully used for several indications with no complications reported. The treatment of nail psoriasis has still yielded unsatisfactory results, leading doctors to search for more effective therapeutic approaches.²

Case 1. We report a 45-year-old female patient with nail dystrophy for 2 years referred for onychomycosis treatment. She presented with exuberant subungual hyperkeratosis and discrete cupuliform depressions, in addition to erythema, periungual edema, and pain in the right hallux (Figure 1). No changes were observed in the other nails. We observed no other alterations on physical examination. Direct mycological examination, repeated weekly for 4 weeks, was always negative. MRI of the hallux confirmed the

diagnosis of psoriatic arthritis, initially monoarticular. We started treatment with systemic MTX and continued for 3 months with no response. Then, we used leflunomide for 12 months. The patient reported improvement of joint pain, but we observed a worsening of nail dystrophy. No clinical change was observed after one year. Then, we opted for intralesional infiltration with methylprednisolone, with no improvement. Finally, we used intralesional infiltration with MTX 25 mg/ml (Figure 2). Case 2. A 60-year-old male patient with psoriasis in small plaques affecting seven fingernails. The initial treatment consisted of topical occlusive therapy with calcipotriol and betamethasone associated with infiltrations of methotrexate at 25 mg/ml (Figure 3). The treatment with MTX consisted of the infiltration of 0.1 ml (2.5 mg) per affected nail, every 30 days, for 3 consecutive months. As the applications occurred at the level of the proximal nail fold, in the case of the hallux, the dose was divided with an interval of 1cm. No anesthetic nerve block was performed and patients reported pain. After the third infiltration, the patients demonstrated an evident improvement. Five months after the first application, they still presented a good evolution.



FIGURE 1: Left hallux, pre-treatment. Subungual hyperkeratosis, edema and periungual erythema



FIGURE 2: Progressive improvement of nail dystrophy five months after the first infiltration

Received 07 February 2018.

Accepted 19 August 2018.

* Study conducted at Universidade de Santo Amaro, São Paulo, SP, Brasil.

Financial support: None.

Conflict of interest: None.

¹ Connective Tissue Disease Unit, Dermatology Service, Universidade de Santo Amaro, São Paulo, SP, Brasil.

² Dermatology Service, Universidade de Santo Amaro, São Paulo, SP, Brasil.

MAILING ADDRESS:
 Getúlio Pisa Carneiro
 E-mail: getulliopisa@gmail.com

©2019 by Anais Brasileiros de Dermatologia





FIGURE 3: Right first digit, pre-treatment (left) and after 60 days (2 infiltrations), 1 infiltration every 30 days (right)


The prevalence of nail psoriasis in patients with psoriatic arthritis may exceed 80%. However, nail psoriasis, in the absence of cutaneous or articular disease, is present in 5-10% of patients.³ Nail psoriasis may also be considered as an indicator for patients at risk for future psoriatic arthritis.⁴ Topical treatments mentioned in the literature include calcipotriol, topical steroids, 5-fluorouracil, antralin, tazarotene, and cyclosporine in oil solution. The application of these agents is recommended for up to 6 months, period during which the nail plate grows completely from the matrix to the hyponychium.² The chronic use of corticosteroids has some local side effects such as periungual telangiectasia, skin atrophy, and appearance of the underlying phalanx. In addition, an important complication of intraleisional treatment is phalangeal atrophy and the risk of the disappearing digit.³ MTX, acitretin, and ciclosporin are conventional drugs indicated for the treatment of plaque psoriasis, but with non uniform results in nail psoriasis.¹ However, non-pharmacological therapies such as phototherapy, superficial radiotherapy, grenz rays, and electron therapy show good results in some reports.² Recently, biological agents, such as TNF-alpha inhibitors (etanercept, infliximabe, and adalimumab), anti-IL-12-23 (ustekinumab), and anti-IL-17 (secukinumab) have been used in the treatment of nail psoriasis with good results. However, risks, benefits, and costs must be weighed when the indication is exclusively for the treatment of nail disorders.³ MTX is the first-line drug for the systemic treatment of plaque psoriasis with or without nail injury and in joint disease, and can be administered orally or subcutaneously.³ Intraleisional injection has been successfully used in ectopic pregnancy, early gastric cancer, and keratoacanthoma in dermatology.^{2,4} In the literature, only two cases of intraleisional MTX treatment have been reported, one of which compares the use of MTX, ciclosporin, and intraleisional triamcinolone acetone.⁵ Eight nails were treated and it was possible to notice a significant reversion of onychodystrophy and a sustained improvement of the lesions. However, the second left fingernail showed a slower response to therapy. Therefore, the use of intraleisional MTX, although off-label,


may be a new therapeutic alternative for nail injury. The dosages used were empirical and more studies are needed to determine a pattern of response and adequacy of the dosage. □


REFERENCES

- Schons KRR, Knob CF, Murussi N, Beber AC, Neumaier W, Monticelo OA. Nail psoriasis: a review of the literature. *An Bras Dermatol.* 2014;89:312-7.
- Saricaogly H, Oz A, Turan H. Nail psoriasis successfully treated with intraleisional methotrexate: case report. *Dermatology.* 2011;222:5-7.
- Pasch MC. Nail Psoriasis: A Review of Treatment Options. *Drugs.* 2016;76:675-705.
- Haneke E. Nail psoriasis: clinical features, pathogenesis, differential diagnoses, and management. *Psoriasis (Auckl).* 2017;7:51-63.
- Mittal J, Mahajan BB. Intramatrix injections for nail psoriasis: An open-label comparative study of triamcinolone, methotrexate, and cyclosporine. *Indian J Dermatol Venereol Leprol.* 2018;84:419-23.

AUTHORS' CONTRIBUTIONS

Artur Antônio Duarte  ORCID 0000-0003-0361-9776
Approval of the final version of the manuscript; conception and planning of the study; obtaining, analyzing and interpreting the data; effective participation in research orientation; Intellectual participation in propaedeutic and/or therapeutic approach of the studied cases; critical review of the literature; critical review of the manuscript.

Getullio Pisa Carneiro  ORCID 0000-0001-7748-0282
Approval of the final version of the manuscript; conception and planning of the study; elaboration and writing of the manuscript; obtaining, analyzing and interpreting the data; effective participation in research orientation; critical review of the literature; critical review of the manuscript.

Carolina Malavassi Murari  ORCID 0000-0001-6688-5145
Conception and planning of the study; obtaining, analyzing and interpreting the data; intellectual participation in propaedeutic and/or therapeutic approach of the studied cases.

Laura Costa Beber de Jesus  ORCID 0000-0003-2347-5554
Statistical analysis; obtaining, analyzing and interpreting the data; Effective participation in research orientation; critical review of the literature.

How to cite this article: Duarte AA, Carneiro GP, Murari CM, Jesus LCB. Nail psoriasis treated with intraleisional methotrexate infiltration. *An Bras Dermatol.* 2019;94(4):491-2.