

that correlated CPK variations after physical exercise with body composition and BMI. The present study is the first to demonstrate that patients using ISO that have a high BMI may show a higher elevation in CPK levels. Studies with a larger number of patients are required to confirm this association.

Financial support

None declared.

Authors' contribution

Bruna Maggioni Busetti: Statistical analysis; design and planning of the study; drafting and editing of the manuscript; collection, analysis and interpretation of data.

David Rubem Azulay: Effective participation in research orientation; critical review of the manuscript.

Felipe Aguinaga: Approval of the final version of the manuscript; design and planning of the study; drafting and editing of the manuscript; collection, analysis and interpretation of data; effective participation in research orientation.

Edgar Ollague Cordova: Drafting and editing of the manuscript; collection, analysis and interpretation of data.

Conflicts of interest

None declared.

References

1. Kaymak Y. Creatine phosphokinase values during isotretinoin treatment for acne. *Int J Dermatol.* 2008;47:398–401.
2. Kim J, Lee J. The relationship of creatine kinase variability with body composition and muscle damage markers following eccentric muscle contractions. *J Exerc Nutrition Biochem.* 2015;19:123–9.
3. Heled Y, Bloom MS, Wu TJ, Stephens Q, Deuster PA. CM-MM and ACE genotypes and physiological prediction of the creatine kinase response to exercise. *J Appl Phys.* 2007;103:504–10.
4. Paschalis V, Nikolaidis MG, Giakas G, Theodorou AA, Sakellariou GK, Fatouros IG, et al. Beneficial changes in energy expenditure and lipid profile after eccentric exercise in overweight and lean women. *Scand J Med Sci Sports.* 2010;20:103–11.
5. Kriketos AD, Baur LA, O'Connor J, Carey D, King S, Caterson ID, et al. Muscle fibre type composition in infant and adult populations and relationships with obesity. *Int J Obes Relat Metab Disord.* 1997;21:796–801.

Bruna Maggioni Busetti *, David Rubem Azulay , Felipe Aguinaga , Edgar Ollague Cordova 

Instituto de Dermatologia Professor Rubem David Azulay, Santa Casa de Misericórdia do Rio de Janeiro, Rio de Janeiro, RJ, Brazil

* Corresponding author.

E-mail: drabrunambusetti@gmail.com (B.M. Busetti).

Received 13 December 2019; accepted 16 August 2020

Available online 12 July 2021

<https://doi.org/10.1016/j.abd.2020.08.020>

0365-0596/ © 2021 Sociedade Brasileira de Dermatologia.

Published by Elsevier España, S.L.U. This is an open access article under the CC BY license

(<https://creativecommons.org/licenses/by/4.0/>).

Evaluation of depression, stress and quality of life indexes in patients with atopic dermatitis^{☆,☆☆}



Dear Editor,

Atopic Dermatitis (AD) is a recurrent chronic inflammatory dermatosis, characterized by pruritus, sleep disorders and multiple comorbidities, with significant psychological alterations and impact on mental health.¹ In many cases, there is loss of quality of life, with deep psychosocial and economic impact.²

Some authors have described different degrees of depression and stress in patients with AD.^{1,3} In this study, the authors evaluated depression, stress and quality of life indexes in adults with AD using:

- Beck depression inventory (Beck-BDI-II depression inventory - validated as of 15 years old)⁴;
- Inventory of stress symptoms for adults (ISSL - validated as of 15 years old) Classifies into 4 phases: alert (contact with aggressive stimulus); resistance (adaptation of the body); near exhaustion (depletion) and exhaustion (physical and/or psychological impairment)⁵;
- Dermatology life quality index (DLQI-BRA).

The analyzed variables were: age; gender; the visual scale of daytime and nighttime pruritus, the severity of AD (SCORAD - atopic dermatitis score) using Pearson's correlation with indexes of depression, stress and quality of life.

Statistical analysis was performed using the Adans Statistics Company (CONRE 8250-A).

Thirty-one patients were included, of which 18 were women, 13 men, with a mean age of 29 years, and a standard deviation of 12.2 (minimum 15; maximum 54 years). Of the 31 patients, 7 (22.6%) had mild depressive symptoms, while 5 (16.1%) had mild to moderate; 18 (38.7%) had moderate to severe and 7 (22.6%) had severe symptoms (Fig. 1). Sleep pattern disorders had the highest score (1.74), followed by indecision (1.58), low self-esteem (1.42), and self-criticism (1.39). Suicidal thoughts or ideation had the lowest score (0.26), followed by pessimism (0.55) and loss of sexual interest (0.55).

[☆] How to cite this article: Castro CR, Andrade MEB, Pires RMG, Pires MC. Evaluation of depression, stress and quality of life indexes in patients with atopic dermatitis. *An Bras Dermatol.* 2021;96:627–9.

^{☆☆} Study conducted at the Complexo Hospitalar Padre Bento de Guarulhos, Guarulhos, SP, Brazil.




Figure 1 Beck II depression inventory.




Figure 2 Stress phases.

Of the 30 patients with symptoms of physical stress, 18 (60%) were in the resistance phase, while 11 (36.7%) were in the near-exhaustion phase and only 1 (3.3%) was in the alert phase (Fig. 2). Symptoms of psychological stress were manifested by 22 patients (73.3%), while 4 (13.3%) also reported physical symptoms and another 4 (13.3%) manifested both. There was no association between stress, depression and impaired quality of life with gender or age.

The DLQI-BRA classification showed patients with significant quality of life impairment as follows: 7 (22.6%) showing mild impact, 6 (19.4%) moderate, 14 (45.2%) severe and 4 (12.9%) very severe impact (Fig. 3). The correlation coefficient between daytime pruritus and physical stress was 0.385; with psychological stress, 0.369; with both, 0.412; with depression, 0.311; with quality of life, 0.127. A moderate correlation was observed with stress and depression, and a weak correlation with quality of life. The correlation between depression and anxiety was 0.608, considered moderate to strong.

The correlation coefficients of SCORAD with physical/psychological stress and depression were non-significant (-0.02 and -0.06 respectively, p-value 0.911 and 0.739). There was a weak positive correlation between the degree of severity and impaired quality of life (0.268, p-value = 0.145), which is in accordance with Carvalho et al.² Because it is a reference for the health network for dermatological diseases, most patients had moderate to severe disease; thus, it was concluded that the stress, depression and impaired quality of life indexes do not depend on AD severity.

Psychosocial and emotional factors affect the course of the disease, as well as the response to therapeutic interventions. Other authors have obtained similar indexes of depression (around 15% for severe cases).^{1,3} Silverberg et al. stated that dermatologists should consider these associations when evaluating AD patients.¹ Unlike other authors, the present study found no association with suicidal thoughts or ideation (0.26).³

The present study found a high prevalence of stress (96.8% of participants), with the majority (60.0%) in the resistance phase and a significant predominance of psychological symptoms (73.3%), which was more relevant than in other studies.^{1,3}

The most common psychological symptoms were: sudden increase in motivation; enthusiasm; insecurity; feeling of incompetence; constant thinking on a single subject; excessive irritability; nightmares and fatigue. Classically, these findings are observed when living with greater vulnerability to psychological stress, as it occurs in AD crises. On the other hand, in our experience with atopic dermatitis patients we observe anticipatory distress and self demand, which in turn can lead to crises, creating a vicious circle (personal opinion of the authors).

The authors conclude that individuals affected with atopic dermatitis often show symptoms of depression, stress and impaired quality of life. Itching is among the symptoms that were most often related to psychological distress and reduced quality of life. Thus, it is recommended that dermatologists, when treating patients with AD, also analyze aspects such as stress, depression and quality of life.




Figure 3 Dermatology Life Quality Index (DLQI) Questionnaire.

Financial support

None declared.

Authors' contributions

Cleide Rodrigues de Castro: Design and planning of the study; drafting and editing of the manuscript; collection, analysis, and interpretation of data; critical review of the literature; critical review of the manuscript.

Maria Elisa Bertocco Andrade: Intellectual participation in the propaedeutic and/or therapeutic conduct of the studied cases.

Renata Marli Gonçalves Pires: Critical review of the literature; critical review of the manuscript.

Mario Cezar Pires: Approval of the final version of the manuscript; design and planning of the study; effective participation in research orientation; intellectual participation in the propaedeutic and/or therapeutic conduct of the studied cases; critical review of the literature; critical review of the manuscript.

Conflicts of interest

None declared.

References

- Silverberg JI, Gelfand JM, Margolis DJ, Boguniewicz M, Fonacier L, Grayson MH, et al. Symptoms and diagnosis of anxiety and depression in atopic dermatitis in US adults. *Br J Dermatol.* 2019;181:554–65.
- Carvalho D, Aguiar P, Mendes-Bastos P, Palma-Carlos A, Freitas J, Ferrinho P. Quality of life and characterization of patients with atopic dermatitis in Portugal – QUADep study. *J Invest Allergol Clin Immunol.* 2020;30:430–8.
- Cheng BT, Silverberg JI. Depression and psychological distress in US adults with atopic dermatitis. *Ann Allergy Asthma Immunol.* 2019;123:179–85.
- Cunha JA. Manual da versão em português das Escalas Beck. São Paulo: Casa do Psicólogo; 2001.
- Lipp MEN. Mecanismos neuropsicofisiológicos do stress: teoria e aplicações clínicas, 2003, Casa do Psicólogo; São Paulo. In: Lipp ME, editor. Mecanismos neuropsicofisiológicos do stress: teoria e aplicações clínicas. São Paulo: Casa do Psicólogo; 2003.

Cleide Rodrigues de Castro ^{a,*},
 Maria Elisa Bertocco Andrade ^a,
 Renata Marli Gonçalves Pires ^b, Mario Cezar Pires ^a

^a Complexo Hospitalar Padre Bento de Guarulhos, Guarulhos, SP, Brazil

^b Faculty of Medicine, Universidade de São Paulo, São Paulo, SP, Brazil

* Corresponding author.

E-mail: cleiderodriguesdecastro@gmail.com (C.R. Castro).

Received 7 December 2019; accepted 1 September 2020

Available online 17 July 2021

<https://doi.org/10.1016/j.abd.2020.09.010>

0365-0596/ © 2021 Sociedade Brasileira de Dermatologia.

Published by Elsevier España, S.L.U. This is an open access article under the CC BY license

(<https://creativecommons.org/licenses/by/4.0/>).