

Differences between attendance in emergency care of male and female victims of traffic accidents in Porto Alegre, Rio Grande do Sul State, Brazil

Diferenças entre homens e mulheres vítimas de acidente de trânsito atendidos em emergências de Porto Alegre, RS, Brasil

Raquel Forgiarini Saldanha¹

Flavio Pechansky³

Daniela Benzano³

Carlos Alberto Sampaio Martins de Barros¹

Raquel Brandini De Boni²

Abstract *Driving under the influence of alcohol/drugs (DUI) is a well-established risk factor for traffic accidents, and men and women have different consumption patterns. The scope of this paper is to analyze differences in alcohol and drug consumption, as well as on behavior associated with traffic accidents among men and women. A cross-sectional study was conducted with 609 sequential traffic accident victims attended in emergency care from Porto Alegre. Subjects gave a structured interview, were breathalyzed and had a saliva test for alcohol/drug screening. Results showed that women were mainly passengers or pedestrians ($p < 0.001$). There was no significant difference in positive blood alcohol concentration. However, men reported more binge drinking and THC use, while women had more benzodiazepine in their saliva ($p < 0.05$). This is the first Brazilian study to compare alcohol and drug use among men and women who were the victims of traffic accidents. Results point to differences in the pattern of substance abuse, as well on risk behavior. Data may be useful for specific prevention strategies that take gender differences into consideration.*

Key words *Traffic accident, Gender differences, Alcohol consumption, Drug abuse, Driving under the influence (DUI)*

Resumo *Dirigir sob a influência de álcool/drogas (DUI) contribui para ocorrência de acidentes de trânsito, sendo que homens e mulheres diferem quanto ao seu consumo. Objetivo: Analisar as diferenças no consumo de álcool/drogas e nos comportamentos de risco para dirigir entre homens e mulheres. Método: Estudo transversal, com amostra consecutiva de 609 vítimas de acidentes de trânsito atendidas nas emergências de Porto Alegre. Realizou-se entrevista estruturada, teste de bafômetro e saliva para screening de álcool e drogas. Resultados: As mulheres acidentaram-se principalmente como passageiras e pedestres, ($p < 0.001$). Não houve diferença na triagem para abuso/dependência ou alcoolemia positiva. Porém, os homens referiram mais “beber pesado” e utilizaram mais THC e cocaína, enquanto as mulheres utilizaram benzodiazepínicos ($p < 0.05$). Conclusão: Este é o primeiro estudo brasileiro a comparar uso de álcool e drogas entre homens e mulheres vítimas de acidentes de trânsito. Os dados podem ser úteis na elaboração de estratégias específicas de prevenção que considerem as diferenças de gênero.*

Palavras-chave *Acidente de trânsito, Diferenças de gênero, Álcool e drogas, Dirigir sob a influência (DUI)*

¹ Centro de Estudos Jose de Barros Falcão. Av. Prof. Oscar Pereira 4821, Cascata. 91.712-320 Porto Alegre RS Brasil. raquelsaldanha@gmail.com

² Instituto Nacional de Infectologia Evandro Chagas, Fiocruz.

³ Center for Drug and Alcohol Research, Hospital de Clínicas de Porto Alegre.

Introduction

Traffic accidents (TA) represent a high cost to society, causing more than 1 million deaths worldwide each year, and an even greater number of hospitalizations, physical and psychological sequelae¹. Literature indicates that most TA involve young men, and they suffer such accidents especially as drivers, while women predominate as passengers and pedestrians^{2,3}.

Driving under the influence of alcohol (DUI) increases the risk of TA, and data from a study conducted in São Paulo showed a prevalence of 21.9% of positive of Blood alcohol concentration (BAC) readings on weekends in the city of Diadema⁴, which is much higher than other countries (usually less than 10%). Some of the DUI-associated factors include: being male, unmarried, binge drinking (five or more drinks on one occasion)⁵⁻⁷, having been a passenger of a DUI driver, little insight about punishment, low socioeconomic status, and low education⁶. Some studies also show that driving a vehicle alone increases the chance of TA involving alcohol^{6,8,9}. This, however, does not apply to young people who usually DUI and have passengers in the car. In addition, drivers who suffer TA involving alcohol tend to drink more in bars, and on weekends^{6,7}.

Other psychoactive substances also affect driving skills, and may increase TA risk¹⁰. In a cohort study conducted in New Zealand involving 969 drivers who suffered TA, it was found that 24% of those who had drunk also consumed cannabis and LSD⁶.

Consistently, the literature indicates that men have higher risk for both TA, and for DUI. However, there is no Brazilian data comparing risk behaviors associated with the TA between men and women. Thus, the aim of this study was to analyze the differences in alcohol and drugs consumption, as well as risk behaviors among men and women who were attended in emergencies of Porto Alegre after a traffic accident.

Methods

This was a cross-sectional study with consecutive sampling conducted in the two hospitals that attend more than 90% of trauma victims of the city of Porto Alegre, the southernmost state capital of Brazil. All TA victims who were over 18 years of age and who suffered TA either as driv-

ers, pedestrians or riders, between october-november 2009, were approached after initial evaluation. The study was approved by the Ethics Committee in Research of Hospital de Clinicas de Porto Alegre, and methodological details are described elsewhere¹¹.

Demographic data and data related to risk behaviors were collected through structured interviews. BAC was measured using breath test (model ALCO-SENSOR IV™, *Intoximeters* Inc, Devon, UK) and substance use (cannabis, cocaine and benzodiazepines) through saliva test (Quantisal™, Immunalysis Corporation, Pomona, CA, USA), analyzed by ELISA. Drug abuse and dependence were evaluated using the Mini International Neuropsychiatric Interview (MINI).

Data were analyzed with SPSS.16, chi-square test for categorical variables and Student t test for quantitative variables were used.

Results

The sample comprised 441 men (72.4%) and 168 (27.6%) women with a mean age of 32 and 34 years, respectively. 92.6% of those eligible were approached, and 10.9% refused. Women had higher education, lower proportion of driver's license, and were injured mainly as passengers and pedestrians ($p < 0.001$), as can be seen in Table 1.

There was no statistical difference between gender neither for alcohol abuse/dependence nor for positive BAC. Men reported more binge drinking and used more cocaine and THC, while women used more benzodiazepines ($p < 0.05$) (Table 2).

Discussion

According to the literature, regardless of the type of accident, the men are always more involved¹². The larger number of women passengers and pedestrians in our sample is consistent with the literature, and may reflect social norms in which men usually takes the driving^{2,3,13,14}. These standards may also be associated with differences in substance use among men and women. Studies have shown that in developing countries, where the social norms are more traditional, women have higher rates of abstinence from alcohol while men drink more and have more binge drinking^{15,16}. In communities where women have less conservative atti-

Table 1. Demographics and risk behavior for traffic accidents among men and women on emergency rooms of Porto Alegre, in 2008.

	N*	Men	Women	P**
Age	609	32.0 (+/- 12.1)	34.9 (+/- 15.5)	0.027
Schooling	439			< 0.001
Until 8 th grade		167 (38.0)	39 (23.4)	
High school		218 (49.7)	73 (43.7)	
College or more		54 (12.8)	55 (32.9)	
Had driver license	603	345 (78.9)	56 (33.7)	< 0.001
Situation in the Traffic accident	606			< 0.001
Driver		330 (75.2)	31 (18.6)	
Passenger		54 (12.3)	84 (50.3)	
Pedestrian		55 (12.5)	52 (31.1)	
Time of Traffic accident	581			0,107
22h to 8h		112 (23.4)	29 (18.5)	
8h to 16h		159 (37.5)	70 (44.36)	
16h to 22h		153 (36.1)	58 (36.9)	
Weekend	580	193 (45.6)	69 (43.9)	0.790
Coming from	426			0.012
Pub/ Party		20 (6.0)	12 (12.9)	
Home		186 (55.9)	58 (62.4)	
Work		127 (38.1)	23 (24.7)	
Accompanied by under 18	474	28 (7.4)	15 (15.6)	0.021
Had use seat belt	132	47 (67.1)	31 (50)	0.068
Previous Traffic accident	609	176 (40.0)	30 (18.1)	< 0.001
Hospitalization	605	46 (10.5)	16 (9.6)	0.557

* N of men and women analyzed for each variable. **Chi-square for categorical variables and t Student test for continuous variables.

Table 2. Alcohol and other drugs among men and women on emergency rooms of Porto Alegre, in 2008.

	Men N (%)	Women N (%)	P*	RP (CI95%)**
Binge drinking in the last 12 months	163 (38.4)	31 (19.5)	< 0.001	2.0 (1.4-2.8)
Alcohol Abuse/dependence	38 (9.0)	9 (5.7)	0.257	1.6 (0.8-3.2)
Reported drinking in last 24h	95 (29.6)	27 (26)	0.557	1.3 (0.9-2.0)
Positive Blood alcohol concentration	34 (8.8)	10 (7.1)	0.662	1.2 (0.6-2.4)
Cannabis (THC)	38 (11.9)	4 (3.3)	0.010	3.6 (1.3-9.9)
Benzodiazepines	8 (2.8)	9 (8.3)	0.025	0.3 (0.1-0.9)
Cocaine	28 (8.9)	1 (0.8)	0.005	10.7 (1.5-78.0)
Drive under the influence in the last 12 months	55 (16.9)	10 (9.3)	0.083	1.8 (1.0-3.4)

*chi square. **Prevalence rate.

tudes, their drinking patterns are similar to men, as well as the woman's role in society. This difference may also be associated with the lower proportion of women with driver's license in the sample, which is consistent with data from

the general population, where 75% of drivers are men¹⁷.

Although most drivers used seat belts, contrary to international studies women in our sample used tended to use less¹⁸. We do not know

whether these women were in the backseat of the cars - and therefore did not use the device, which is common practice in Brazil. Another explanation would be the high prevalence of substance use, since individuals who drink and drive adopt less safety behaviors even when sober⁷.

Alcohol was the most frequent drug used by victims, followed by cannabinoids. Although the prevalence of positive BAC measured by breathalyzer was 7.22%, 27.8% of patients reported drinking within 24 hours prior to the TA. This value is consistent with another Brazilian study, where the prevalence of reported alcohol use was 27.7 %¹⁴. This difference can be justified by the time between the accident and sample collection. A Brazilian study that aimed to describe the frequency of reports of alcohol consumption among emergency room care (victims of accidents and violence) found the proportion of alcohol consumption among men was three times greater than among women¹⁹.

Consistent with previous studies, women had used more benzodiazepines and men had more cannabis and alcohol^{9,20}. Even though Brazilian

studies on drug use conducted in emergency departments are scarce, and generally restricted to the use of alcohol, a study in São Paulo found the following prevalence of substance abuse: THC 13.6%, cocaine 3.3% and benzodiazepines 4.2%²¹. Data are similar to those we found, but draw attention to the high prevalence of cocaine use among the men in our sample, which may be associated to the increase in crack use in Brazil^{22,23}.

This is the first Brazilian study to compare risk behaviors among men and women with traffic accidents, and it must be read under the scope of its limitations: 1) the sample was consecutive and cannot be generalized to all TA victims; 2) individuals under 18 years were excluded, which may have underestimated the use of substances; 3) The time between the TA and the collection of material for analysis may have underestimated drug use prevalence. However, data confirm international studies, and point to the need of specific prevention strategies that consider gender differences in Brazil.

Collaborations

RF Saldanha, R De Boni, F Pechansky, D Benzano e CASM Barros participated equally in all stages of preparation of the article.

References

1. Peden M SR, Slet D, Mohan D, Hyder AA, Jarawan E, Mathers C, editors. *World report on traffic injury prevention*. Geneva: World Health Organization; 2004.
2. Andrade S, Jorge M. Victims' characteristics by road accidents in a city of Southern Brazil. *Rev Saude Publica* 2000; 34(2):149-156.
3. Gawryszewski V, Coelho H, Scarpelini S, Zan, Jorge M, Rodrigues E. Land transport injuries among emergency department visits in the state of São Paulo, in 2005. *Rev Saude Publica* 2009; 43(2):275-282.
4. Duailib S PI, Laranjeira R. Prevalência do beber e dirigir em Diadema, estado de São Paulo. *Rev Saude Publica* 2007; 41(5):1058-1061.
5. Caetano R, McGrath C. Driving under the influence (DUI) among U.S. ethnic groups. *Accident Analyses and Prevention* 2005; 37(2):217-224.
6. Morrisson L, Begg DJ, Langley JD. Personal and situational influences on drink and driving and sober driving among a cohort of young adults. *Injury Prevention* 2002; 8(2):111-115.
7. Bingham C, Elliot M, Shope J. Social and Behavioral Characteristics of Young Adult Drink/Drivers Adjusted for Level of Alcohol Use. *Alcohol Clin Exp Res* 2007; 31(4):655-664.
8. Nelson T, Isaac N, Kennedy BP, Graham JD. Factors associated with planned avoidance of alcohol-impaired driving in high-risk men. *J Stud Alcohol* 1999; 60(3):407-412.
9. Longo M, Hunter C, Lokan R, White J, White M. The prevalence of alcohol, cannabinoids, benzodiazepines and stimulants amongst injured drivers and their role in driver culpability Part I: the prevalence of drug use in drivers, and characteristics of the drug-positive group. *Accid Anal Prev* 2000; 32(5):613-622.
10. Ojaniemi KK, Lintonen TP, Impinim AO, Lillsunde PM, Ostamo AL. Trends in driving under the influence of drugs: A register-based study of DUID suspects during 1977-2007. *Accid Anal Prev* 2009; 41(1):191-196.
11. Soibelman M, Benzano D, Von Diemen L, De Boni R, Pechansky F. *Consumo de álcool e outras drogas entre vítimas de acidentes de trânsito atendidas em emergências de Porto Alegre. Uso de bebidas alcoólicas e outras drogas nas rodovias brasileiras e outros estudos*. Porto Alegre: Secretaria Nacional de Políticas sobre Drogas; 2010.
12. Golias A, Caetano R. Acidentes entre motocicletas: análise dos casos ocorridos no estado do Paraná entre julho de 2010 e junho de 2011. *Cien Saude Colet* 2013; 18(5):1235-1246.
13. Marmor M, Parnes N, Aladgem D, Birshan V, Sorkine P, Halpern P. Characteristics of Road Traffic Accidents Treated in an Urban Trauma Center. *Isr Med Assoc J* 2005; 7(1):9-12.
14. Rodrigues CS, Ladeira RM, Pereira JC, Paula IM. Saúde em trânsito: Pesquisa de acompanhamento de vítimas de acidentes de trânsito em Belo Horizonte. Belo Horizonte: Secretaria de Saúde de Belo Horizonte - SM-SA-BH; [2005]. Empresa de Transportes e Trânsito de Belo Horizonte - BHTRANS, 2005.
15. Almeida-Filho N, Lessa I, Magalhães L, Araújo M, Aquino E, Kawachi I, James SA. Alcohol drinking patterns by gender, ethnicity, and social class in Bahia, Brazil. *Rev Saude Publica* 2004; 38(1):45-54.
16. Wilsnack R, Vogelanz N, Wilsnack S, Harris T, Ahlström, S, Bondy S, Csémy L, Ferrence R, Ferris J, Fleming J, Graham K, Greenfield T, Guyon L, Haavio-Mannila E, Kellner F, Knibbe R, Kubicka L, Loukomskaja M, Mustonen H, Nadeau L, Narusk A, Neve R, Rahav G, Spak F, Teichman M, Trocki K, Webster I, Weiss S. Gender differences in alcohol consumption and adverse drinking consequences: cross-cultural patterns. *Addiction* 2008; 95(2):251-265.
17. Denatran AdI. *Dia Nacional do Motorista 2006*. [cited 2010 out 14]. Available from: http://www.denatran.gov.br/ultimas/20060725_diamotorista.htm
18. Ouimet MC, Morton BGS, Noelcke EA, Williams AF, Leaf WA, Preusser DF, Hartos JL. Perceived risk and other predictors and correlates of teenagers safety belt use during the first year of licensure. *Traffic Inj Prev* 2008; 9(1):1-10.
19. Mascarenhas M, Malta D, Silva M, Gazal-Carvalho C, Monteiro R. Consumo de álcool entre vítimas de acidentes e violências atendidas em serviços de emergência no Brasil, 2006 e 2007. *Cien Saude Colet* 2009; 14(5):17896-1796.
20. Blows S, Ivers R, Connor J, Ameratunga S, Woodward M, Norton R. Marijuana use and car crash injury. *Addiction* 2005; 100(5):605-611.
21. Reis A, Figlie N, Laranjeira R. Prevalence of substance use among trauma patients treated in a Brazilian emergency room. *Rev Bras Psiquiatr* 2006; 28(3):191-195.
22. Ferri C, Laranjeira R, Silveira Dd, Dunn J, Formiconi M. Aumento da procura de tratamento por usuários de crack em dois ambulatórios na cidade de São Paulo, nos anos de 1990 a 1993. *Rev. Assoc Médica Brasil* 1997; 43(1):25-28.
23. Duailibi L, Ribeiro M, Laranjeira R. Profile of cocaine and crack users in Brazil. *Cad Saude Publica* 2008; 24(4):545-557.

Artigo apresentado em 29/06/2013

Aprovado em 25/07/2013

Versão final apresentada em 30/07/2013

