



# The Brazilian Journal of INFECTIOUS DISEASES

www.elsevier.com/locate/bjid



## Original article

# Aeromycological study at the intensive care unit of the “Dr. Manuel Gea Gonzalez” General Hospital

José Manuel Ríos-Yuil<sup>a,b,\*</sup>, Roberto Arenas<sup>a</sup>, Ramón Fernández<sup>a</sup>,  
María Calderón-Ezquerro<sup>c</sup>, Raymundo Rodríguez-Badillo<sup>a</sup>

<sup>a</sup> Hospital General “Dr. Manuel Gea González”, Ciudad de México, Mexico

<sup>b</sup> Hospital General de México, Ciudad de México, Mexico

<sup>c</sup> Instituto Nacional de Ciencias de la Atmósfera, Universidad Nacional Autónoma de México, Ciudad de México, Mexico

## ARTICLE INFO

### Article history:

Received 9 April 2012

Accepted 24 June 2012

Available online 10 September 2012

### Keywords:

Fungi

Air

Sampling

Colony-forming units

## ABSTRACT

**Introduction:** An aeromycological study verifies the presence and quantifies the concentration of fungal propagules in the air. It is very important in the hospital setting because of the increasing numbers of immunosuppressed and severely ill patients. The objective of this study was to determine the concentration of fungi in the air of the intensive care unit (ICU) of “Dr. Manuel Gea González” General Hospital.

**Methods:** This is a descriptive, observational cross-sectional study. Air samples were obtained with a single stage Thermo-Andersen Viable Particle Sampler (Thermo Electron Corporation - Massachusetts, U.S.A.) in a Petri dish with potato dextrose agar for 15 minutes at two different times (morning and afternoon) and heights (1 and 1.5 meters). The Petri dishes were incubated for five to seven days at 27° C, the number of colonies was counted, and the total CFU/m<sup>3</sup> was determined. The isolated fungal genera were identified by morphological features. Epi Info v. 3.4.3 © was used for statistical analysis.

**Results:** The mean concentration of fungi in the air of the ICU was 85.08 ± 29.19 CFU/m<sup>3</sup>; while in the outside air it was 84.3 ± 17.23 CFU/m<sup>3</sup> (p = 0.96). The fungi isolated were: *Cladosporium* spp., *Penicillium* spp., *Aspergillus* spp. (non-*fumigatus*), *Fusarium* spp., *Exophiala* spp., *Syncephalastrum* spp., and *Acremonium* spp.

**Discussion:** Fungal spores were found in the air of the ICU and *Cladosporium* spp. was the most frequently isolated fungi. There was no difference according to sampling time or height.

© 2012 Elsevier Editora Ltda. All rights reserved.

## Introduction

Bioaerosols are aerial suspensions of particles from live organisms, microorganisms or other biological materials.<sup>1–3</sup>

Aeromycology is the branch of aerobiology that studies the dispersion of spores and other fungal elements in indoor and outdoor air, the changes in their concentrations, and the factors that affect those changes.<sup>3</sup> Fungal spores enter hospitals

\* Corresponding author at: Servicio de Dermatopatología, Hospital General de México, Dr. Balmis 148, Colonia Doctores, Delegación Cuauhtémoc, México DF, Mexico.

E-mail address: jmriosyuil@hotmail.com (J.M. Ríos-Yuil).

1413-8670/\$ – see front matter © 2012 Elsevier Editora Ltda. All rights reserved.

<http://dx.doi.org/10.1016/j.bjid.2012.08.012>

through ventilation systems and fungi develop on multiple surfaces, releasing more spores.<sup>1,3</sup> The number of spores in indoor air varies depending on climate, weather, air currents, humidity, temperature, time of the day, type and maintenance of ventilation systems, age of the buildings, movement of people, cleaning, and the presence of plants or food.<sup>1,3-7</sup>

Immunosuppressed patients with severe neutropenia, chronic granulomatous disease, and acquired immunodeficiency syndrome (AIDS) have the highest risk of developing invasive fungal infection.<sup>8-10</sup> The main fungal genera related with these diseases are *Aspergillus*, *Candida*, *Fusarium*, *Penicillium*, *Mucor*, and *Rhizopus*.<sup>11-17</sup> Invasive aspergillosis (especially by *A. fumigatus*), candidemia, disseminated fusariosis, infections by *P. marneffeii*, and zygomycosis have a mortality rate that can reach 100%.<sup>10-17</sup>

The single stage Thermo-Andersen Viable Particle Sampler (Thermo Electron Corporation – Massachusetts, USA) is the most frequently used air sampler.<sup>8,18,19</sup> In the detection phase, the amount of colony forming units (CFU) per cubic meter of air is determined, and then the identification of the fungi is performed by culture or by molecular techniques.<sup>7,9,18-21</sup>

Because the number of immunosuppressed and severely ill patients is increasing worldwide, especially in the hospital setting, this study was designed to determine the concentration of fungi in the air of the intensive care unit (ICU) of this hospital.

## Materials and methods

This was a descriptive, observational, cross-sectional study. The universe was the air of the ICU of “Dr. Manuel Gea Gonzalez” General Hospital. The ICU is divided in different sections with different air volumes: five single rooms (30 m<sup>3</sup> each), one double room (120 m<sup>3</sup>), one aisle (67.5 m<sup>3</sup>), and one nurse module (37.5 m<sup>3</sup>). For convenience, an air sample was taken from the area just outside the ICU (one meter away from the entrance door), from one single room (picked randomly), from the double room, from the aisle, and from the nurse module. The total air volume of the areas that were sampled inside the ICU was 255 m<sup>3</sup>. The air volume sampled from the five selected areas (inside and outside the ICU) was 0.1415 m<sup>3</sup>.<sup>14</sup> The unit is ventilated by a central air conditioning unit without fans or open windows. The unit can accommodate up to seven patients and always has at least ten staff members working. The highest level of activity occurs during morning hours. Among the studied variables are: humidity, temperature, CFU per air volume, and fungal agents.

Air samples were obtained with a single stage Thermo-Andersen Viable Particle Sampler (Thermo Electron Corporation – Massachusetts, USA) in a Petri dish with potato dextrose agar. The sampler was placed in the center of each room, and each sample was collected with a vacuum flow of 28.3 liters/min for 15 minutes.<sup>14,22</sup> In each sampling area, four samples were collected: one at each time period (morning and afternoon) and one at each height (one and 1.5 meters). Humidity and temperature were measured with a hygrothermograph (Control Company – Friendswood, Texas, USA). Nobody was allowed in or out of the room during sampling.

After sampling, the Petri dishes were incubated for five to seven days at 27° C.<sup>7,14</sup> After that, the number of colonies was counted in every Petri dish. The total number of fungal colonies present in the ICU was determined by obtaining the mean of the colony count in the four areas sampled. This count was stratified by time and height of the sample. The concentration of fungal propagules in the air was expressed in CFU/m<sup>3</sup>. For its calculation, a correction factor was used for the colony count, based on the probability that more than one viable propagule could have passed through the same hole and impacted the culture medium. The formula used was:

$$CFU = N \ln[N/N-P]$$

Where CFU was the corrected colony count, N was the number of holes in the perforated plate of the sampler (400), and P was the number of colonies that grew in the medium.

The corrected count represents the real number of CFUs present in the sampled air. After that, the CFU/m<sup>3</sup> of air was determined with the following formula:

$$CFU/m^3 = \frac{CFU}{t} \times K$$

Where CFU was the corrected colony count, t was the total sampling time expressed in minutes, and K was a conversion factor from cubic feet to cubic meters (K = 35).

The identification of fungi was performed by observation of the macroscopic characteristics of the colonies and the microscopic characteristics of the sporulating hyphae.<sup>5</sup>

The statistical analysis was performed with the Epi Info v. 3.4.3© software, and the means and standard deviations of humidity, temperature, and colony forming units per air volume were determined according to sampling time and height. Independent Student's t-test was used to determine whether there was a statistically significant difference between the means of CFU per air volume according to sampling area, time, and height. Differences were considered significant when the p-value was below 0.05.

All procedures performed were in accordance with and approved by the ethical standards of the “Dr. Manuel Gea Gonzalez” General Hospital Review Board and Ethics Committee. The principles of the Helsinki Declaration of 1975 with the modifications of 1983, and the Mexican General Health Law were followed.

## Results

There was no difference in the fungal spore concentration of the air inside the ICU when compared with the air outside the unit (Table 1). These concentrations were not affected by the time of sampling (p = 0.13 and p = 0.14, respectively) or the height of sampling (p = 0.99 and p = 0.92, respectively) (Table 1). Several colonies were cultured, and the fungal genera isolated were: *Cladosporium* spp., *Penicillium* spp., *Aspergillus* spp. (non-*fumigatus*), *Fusarium* spp., *Exophiala* spp., *Syncephalastrum* spp. and *Acremonium* spp. (Table 2). The mean humidity during sampling was 38.88 ± 3.20% and the mean temperature was 24.09 ± 0.82° C.

**Table 1 – Fungal spore concentrations in the air according to different sampling conditions.**

Sampling conditions	Global	Time of the day		Height of the sampling	
		AM	PM	1 meter	1.5 meters
Inside the ICU*	85.08 ± 29.19	96.7 ± 36.76	73.47 ± 12.9	84.99 ± 32.42	85.18 ± 27.83
Outside the ICU*	84.3 ± 17.23	97.06 ± 9.16	71.54 ± 12.47	83.14 ± 28.86	85.48 ± 7.23

\* In CFU/m<sup>3</sup>.**Table 2 – Fungal genera isolated.**

Fungal genus	Concentration of spores inside the ICU*
<i>Cladosporium</i> spp.	47.34
<i>Penicillium</i> spp.	19
<i>Aspergillus</i> spp.	8.35
<i>Fusarium</i> spp.	2.63
<i>Exophiala</i> spp.	0.44
<i>Syncephalastrum</i> spp.	0.29
<i>Acremonium</i> spp.	0.15
Colony with sterile mycelia	6.88

\* In CFU/m<sup>3</sup>.

## Discussion

In the present study, the mean concentration of fungi in the air of the ICU was 85.08 ± 29.19 CFU/m<sup>3</sup>. This result is similar to that obtained by a study performed in the pulmonology ward at a hospital in Lublin, Poland, where the annual fungal concentration varied from 9.9 to 96.1 CFU/m<sup>3</sup>.<sup>21</sup> No difference was found in this study when the fungal concentration in the air from inside the ICU was compared with the air from outside the unit (p = 0.96). The present findings differ from the results of a study conducted at a university hospital in Rotterdam, Netherlands, where the spore concentration was higher in open areas inside the hospital when compared to the hematology ward.<sup>17</sup>

No differences were found in the concentration of spores in the ICU according to the time of sampling (p = 0.13). This result contrasts with the previously mentioned Polish study, where the concentration of fungal spores was significantly higher in the morning than in the afternoon (p < 0.01).<sup>21</sup>

The fungal genera isolated in the present study were: *Cladosporium* spp., *Penicillium* spp., *Aspergillus* spp., *Fusarium* spp., *Exophiala* spp., *Syncephalastrum* spp. and *Acremonium* spp. None of the fungal colonies isolated in this study corresponded to *Aspergillus fumigatus*. The present results are in accordance with the results of other aeromycology studies. An aeromycology study performed in the ICUs and operating rooms of a hospital in the city of Araraquara, Sao Paulo, Brasil, found that the most isolated fungal genera were *Cladophialophora* spp., *Fusarium* spp., *Penicillium* spp., *Chrysosporium* spp. and *Aspergillus* spp.<sup>5</sup> A study performed in the waiting room of the allergy service of the Infanta Cristina Hospital in Spain showed that most of the fungal spores belonged to the genus *Cladosporium* (*C. cladosporioides* and *C. herbarum*).<sup>6</sup> A research project performed in the ICUs, operating rooms, biomedical laboratories, and lobbies of five general hospitals in Seoul, South Korea, found that the most frequently isolated fungal genera

in the air were: *Cladosporium* spp. (30%), *Penicillium* spp. (20%-25%) and *Aspergillus* spp. (15%-20%).<sup>7</sup> A study conducted at the transplantation ward of a hospital in Bogota, Colombia, found a mean of 2.8 CFU/L of *Aspergillus*.<sup>19</sup>

The presence of the fungi that were isolated from the air of the present ICU, especially *Aspergillus* spp. and *Fusarium* spp., is worrisome because the patients of these units are usually severely ill or immunosuppressed, and this is the group at the highest risk of suffering from potentially fatal invasive fungal diseases.<sup>10,12,13</sup> However, some of these fungal genera also represent a threat to immunocompetent hosts, such as the healthcare staff of the ICU. Fungal spores of several genera, such as *Cladosporium* spp., have been related to asthma exacerbation.<sup>3</sup> *Aspergillus* spp. has been associated with several diseases such as rhinosinusitis, chronic cavitary pulmonary aspergillosis, aspergilloma, allergic bronchopulmonary aspergillosis, and skin and wound infections, among others.<sup>3,23</sup> Meanwhile, spores of *Aspergillus* spp., *Fusarium* spp., *Cladosporium* spp., and *Penicillium* spp. can produce mycotoxin-related disease.<sup>9,18</sup>

The present results underline the importance of establishing control measures to improve air quality inside the ICU, aiming to reduce fungal-related morbidity and mortality. Among these measures, the following can be mentioned: periodic measurements of fungal propagules in the air with aeromycological studies, rigorous cleaning with disinfectants and dust removal, routine equipment maintenance, humidity control, air filtration with high efficiency particle air filters, and use of laminar flow ventilation systems, among others.<sup>1,4,24</sup>

## Conclusion

Fungal spores were found in the air of the ICU (85.08 ± 29.19 CFU/m<sup>3</sup>), and *Cladosporium* spp. was the most isolated fungi. There was no difference according to sampling time or height.

## Conflict of interest

All authors declare to have no conflict of interest.

## Acknowledgements

To the staff of the research unit and the ICU of "Dr. Manuel Gea Gonzalez" General Hospital, and to Dr. Yamilett Morales, for their collaboration.

## REFERENCES

1. Srikanth P, Sudharsanam S, Steinberg R. Bio-aerosols in indoor environment: Composition, health effects and analysis. *Indian J Med Microbiol.* 2008;26:302-12.
2. Angenent LT, Kelley ST, St Amand A, Pace NR, Hernandez MT. Molecular identification of potential pathogens in water and air of a hospital therapy pool. *Proc Natl Acad Sci USA.* 2005;102:4860-5.
3. Kasprzyk I. Aeromicrobiology – main research fields of interest during the last 25 years. *Ann Agric Environ Med.* 2008;15:1-7.
4. Cordeiro RA, Brilhante RS, Pantoja LD, et al. Isolation of pathogenic yeasts in the air from hospital environments in the city of Fortaleza, northeast Brazil. *Braz J Infect Dis.* 2010;14:30-4.
5. Martins-Diniz JN, da Silva RA, Miranda ET, Mendes-Giannini MJ. Monitoring of airborne fungus and yeast species in a hospital unit. *Rev Saude Publica.* 2005;39:398-405.
6. Tormo Molina R, Gonzalo Garijo MA, Muñoz Rodríguez AF, Silva Palacios I. Pollen and spores in the air of a hospital out-patient ward. *Allergol Immunopathol (Madr).* 2002;30:232-8.
7. Kim KY, Kim YS, Kim D. Distribution characteristics of airborne bacteria and fungi in the general hospitals of Korea. *Ind Health.* 2010;48:236-43.
8. Eames I, Tang JW, Li Y, Wilson P. Airborne transmission of disease in hospitals. *J R Soc Interface.* 2009;6 Suppl 6:S697-702.
9. Humphreys H. Microbes in the air - when to count! (The role of air sampling in hospitals). *J Med Microbiol.* 1992;37:81-2.
10. de Pauw BE. What are fungal infections? *Mediterr J Hematol Infect Dis.* 2011;3:e2011001.
11. Chazalet V, Debeaupuis JP, Sarfati J, et al. Molecular typing of environmental and patient isolates of *Aspergillus fumigatus* from various hospital settings. *J Clin Microbiol.* 1998;36:1494-500.
12. Mayayo E, Landeyro J, Stchigel AM, Gazzoni A, Capilla J. Infiltración perineural por células fúngicas. Presentación de un caso y revisión de la literatura. *Rev Iberoam Micol.* 2010;27:94-7.
13. Távora LG, Gambale W, Heins-Vaccari EM, et al. Comparative performance of two air samplers for monitoring airborne fungal propagules. *Braz J Med Biol Res.* 2003;36:613-6.
14. Nunes ZG, Martins AS, Altoe AL, et al. Indoor air microbiological evaluation of offices, hospitals, industries, and shopping centers. *Mem Inst Oswaldo Cruz.* 2005;100:351-7.
15. Nielsen PV. Control of airborne infectious diseases in ventilated spaces. *J R Soc Interface.* 2009;6 Suppl 6:S747-55.
16. Maschmeyer G, Neuburger S, Fritz L, et al. A prospective, randomized study on the use of well-fitting masks for prevention of invasive aspergillosis in high-risk patients. *Ann Oncol.* 2009;20:1560-4.
17. Leenders AC, van Belkum A, Behrendt M, Luijendijk A, Verbrugh HA. Density and molecular epidemiology of *Aspergillus* in air and relationship to outbreaks of *Aspergillus* infection. *J Clin Microbiol.* 1999;37:1752-7.
18. Mobin M, do Amparo Salmato M. Fungus microbiota in air conditioners in intensive care units in Teresina, Piauí. *Rev Soc Bras Med Trop.* 2006;39:556-9.
19. Cárdenas MX, Cortes JA, Parra CM. *Aspergillus* spp. in risk areas of transplant patients in a university hospital. *Rev Iberoam Micol.* 2008;25:232-6.
20. Calderon C, Ward E, Freeman J, McCartney A. Detection of airborne fungal spores sampled by rotating-arm and Hirst-type spore traps using polymerase chain reaction assays. *Aerosol Science.* 2002;33:283-96.
21. Augustowska M, Dutkiewicz J. Variability of airborne microflora in a hospital ward within a period of one year. *Ann Agric Environ Med.* 2006;13:99-106.
22. Andersen Viable Particle Sampler, Thermo Electron Corporation. Environmental Instruments. Series 10-800. Single Stage Viable Sampler. Instruction manual P/N 100074-00. Massachusetts, USA: Thermo Electron Corporation; 2003. p. 13.
23. Hedayati MT, Pasqualotto AC, Warn PA, Bowyer P, Denning DW. *Aspergillus flavus*: human pathogen, allergen and mycotoxin producer. *Microbiology.* 2007;153 Pt 6: 1677-92.
24. Warris A, Voss A, Verweij PE. Hospital sources of *Aspergillus* species: new routes of transmission? *Rev Iberoam Micol.* 2001;18:156-62.