



ELSEVIER

The Brazilian Journal of INFECTIOUS DISEASES

www.elsevier.com/locate/bjid



Original article

Sexually transmitted diseases among psychiatric patients in Brazil

Maria Rita Teixeira Dutra^{a,b}, Lorenza Nogueira Campos^{a,b},
Mark Drew Crosland Guimarães^{a,c,*}

^a Epidemiology and Health Evaluation Research Group, Universidade Federal de Minas Gerais (UFMG), Belo Horizonte, MG, Brazil

^b FHEMIG – Hospital Eduardo de Menezes, Belo Horizonte, MG, Brazil

^c Department of Preventive and Social Medicine, School of Medicine, UFMG, Belo Horizonte, MG, Brazil

ARTICLE INFO

Article history:

Received 30 November 2012

Accepted 18 April 2013

Available online 6 August 2013

Keywords:

Psychiatric patients

Sexually transmitted infections

Brazil

ABSTRACT

Sexually transmitted diseases are still highly prevalent worldwide and represent an important public health problem. Psychiatric patients are at increased risk of sexually transmitted diseases but there are scarce published studies with representative data of this population. We sought to estimate the prevalence and correlates of self-reported sexually transmitted diseases among patients with mental illnesses under care in a national representative sample in Brazil ($n = 2145$). More than one quarter of the sample (25.8%) reported a lifetime history of sexually transmitted disease. Multivariate analyses showed that patients with a lifetime sexually transmitted disease history were older, had history of homelessness, used more alcohol and illicit drugs, suffered violence, perceived themselves to be at greater risk for HIV and had high risk sexual behavioral: practised unprotected sex, started sexual life earlier, had more than ten sexual partners, exchanged money and/or drugs for sex and had a partner that refused to use condom. Our findings indicate a high prevalence of self-reported sexually transmitted diseases among psychiatric patients in Brazil, and emphasize the need for implementing sexually transmitted diseases prevention programs in psychiatric settings, including screening, treatment, and behavioral modification interventions.

© 2013 Elsevier Editora Ltda. All rights reserved.

Introduction

Sexually transmitted diseases (STDs) are still highly prevalent worldwide. According to the 2005 World Health Organization (WHO) estimates,¹ 448 million new cases of curable STDs (e.g., syphilis, gonorrhea, chlamydia, and trichomoniasis) occur annually throughout the world, not including HIV and other STDs (like herpes or hepatitis), and, among these, 50 million cases occur in the Americas.¹

In 2005, the Brazilian Ministry of Health evaluated the prevalence of STDs on three different populations in six Brazilian capitals: pregnant women, male workers from small industries (both considered to be at an average risk for STDs), and patients from an STD clinic (male and female sample of a group regarded as having a high-risk sexual behavior).³ The prevalence of at least one STD in each of these three populations was, respectively, 42.0%, 5.2% and 51.0%. The prevalence of STDs was higher among younger age groups (20 years old

* Corresponding author at: Av. Professor Alfredo Balena, 190, Belo Horizonte, MG 30130-100, Brazil.

E-mail address: drew@medicina.ufmg.br (M.D.C. Guimarães).

1413-8670/\$ – see front matter © 2013 Elsevier Editora Ltda. All rights reserved.

<http://dx.doi.org/10.1016/j.bjid.2013.04.004>

or less), among those with multiple partners, who practised unsafe sex, anal sex, and in those who used injected drugs. In another national survey,⁴ 35.6% of truck drivers reported a lifetime history of STD and, relative to those without STD history, they were older, consumed more illicit drugs, had been incarcerated more often, and had more sex with sex workers.

STDs also represent an important public health problem due to their complications such as neonatal syphilis – which may result in stillbirth, neonatal death or malformation¹; cervical cancer caused by human papilloma virus (HPV)⁵; and pelvic inflammatory disease and infertility due to *Chlamydia*.⁵ In addition, STDs, especially genital ulcerative diseases, can facilitate transmission and acquisition of HIV.⁶ A study⁷ using anonymous HIV testing and counseling program, conducted in Taiwan from 2006 to 2010, showed 10.7% of participants with at least one STD and 3.5% with HIV in that sample. In the same study, the HIV prevalence was 14.7% among patients with previous STDs, while among those without STDs history it was only 3.0%.⁷

Patients with chronic mental illness are at increased risk of STDs and present elevated rates of high risk sexual behavior.⁸ A national sample of psychiatric patients in Brazil⁹ demonstrated that these patients are sexually active at rates that are similar to those of the overall adult Brazilian population.¹⁰ Furthermore, only 16.0% of these psychiatric patients used condoms in the last six months, in comparison to 20.6% among estimates for the overall Brazilian adults during the last year.¹¹ In addition, the use of psychoactive drugs was higher among psychiatric patients compared to the general population, 25.1%⁹ and 8.9%,¹² respectively. In the same sample of psychiatric patients,⁹ unprotected sexual behavior was associated with having sex under the influence of alcohol and having multiple sexual partners.

However, there are few published studies addressing STDs among patients with psychiatric illnesses.¹³ A systematic review indicated that most of the published studies on this population are from developed countries and are based on relatively small and non-representative samples.¹⁴ The prevalence of selected STDs among psychiatric patients varied from 0.8% to 29.0% for HIV; from 1.6% to 66.0% for Hepatitis B; from 0.4% to 38.0% for Hepatitis C; and from 1.1% to 7.6% for syphilis.¹⁴ Over one-third (38.0%) of a sample of patients from two American psychiatric hospitals reported a lifetime history of one or more STDs.¹⁵ HIV-related knowledge, greater self-perceived HIV risk, higher intention to use condom and higher rates of unprotected vaginal and anal sex emerged as significant correlates of a prior STD.

The objective of this study was to estimate the prevalence and correlates of lifetime self-reported STDs among patients with mental illnesses under care in a national representative sample in Brazil.

Materials and methods

Study design and participants

The sample for the current analysis was drawn from a larger national multicenter cross-sectional study conducted in 11 public psychiatric hospitals and 15 public mental health

outpatient clinics (CAPS) in Brazil in 2006, designed to assess risk behavior and sexually transmitted infections/HIV prevalence in a national representative sample of patients with chronic mental illnesses under care ($n=2475$) previously described.^{16,17} Briefly, participants had to be 18 years old or over and be capable of providing written informed consent.

A preliminary assessment was carried out by mental health professionals in order to evaluate the subjects' capacity to participate and sign the consent form – those with delusional symptoms, acute psychosis and a severe degree of mental retardation were not eligible. Ethical approval was obtained from the Federal University of Minas Gerais (UFMG/ETIC 125/03) and the National Ethical Review Board (CONEP 592/2006), and all participants signed written informed consent. For the current analysis, only those participants who reported being sexually active at least once in life were included ($n=2145$).

Data collection

A semi-structured person-to-person interview was conducted using a previously tested and validated questionnaire.¹⁷ It was administered by experienced and trained mental health care professionals and aimed at obtaining sociodemographic, clinical and behavioral data. The main outcome in this analysis was lifetime self-reported history of STDs and it was assessed with the question: "Have you had any disease transmitted through sexual intercourse or venereal disease?". Additional questions were asked regarding three main syndromic groups (genital/anal ulcerative diseases, warts, and discharge) and specific STD medical diagnoses (e.g. syphilis, herpes, chancroid, chlamydia, gonorrhoea, lymphogranuloma venereum, and condyloma).

Good to excellent reliability (intra and inter-rater) was found for all self-reported data used in the present analysis, including being sexually active ($\kappa=0.76$) and having an STD history ($\kappa=0.70$).¹⁸

Explanatory variables included in this analysis were: sociodemographic data (e.g. age, gender, skin color, marital status, schooling, family income, living situation, history of homelessness); clinical characteristics (e.g. type of recruitment center, psychiatric diagnosis, STD history, previous psychiatric hospitalization, previous HIV testing); and, behavioral data (e.g. lifetime tobacco, alcohol, illicit drug and injection drug use, sex under the influence of alcohol/drugs, lifetime unprotected sex, having a partner that refused to use condom, exchanging money/drugs for sex, number of sexual partners, age at first sexual intercourse, lifetime verbal physical or sexual violence, lifetime incarceration, HIV/AIDS knowledge, and self-perception of HIV risk).

Psychiatric diagnoses were obtained from medical charts and were coded according to the International Classification of Diseases-10th Edition (ICD-10). When more than one psychiatric diagnosis was present, these were hierarchically grouped according to clinical severity as follows: (1) schizophrenia (and other psychotic disorders), depression with psychotic symptoms and bipolar disorder; (2) depression, anxiety and others; (3) substance use disorder. Living conditions at the time of the interview were defined as stable, when participants reported living in houses, apartments, or hospitals, or unstable, when

participants reported living on the streets, public shelters, garages, rooms or other ill-defined places. Condom use was assessed as always, sometimes (more than 50% of the time), rarely (less than 50% of the time), and never. For this analysis, unprotected sex was defined as not always using condoms in all sexual practices during lifetime. HIV/AIDS knowledge was assessed through 10 true or false questions and good knowledge was defined as having eight or more correct answers. These statements have been used in prior studies and were derived from Brazilian population-based studies.^{19,20} Self-perception of HIV risk was categorized as high or some, no risk, and did not know.

Statistical analysis

The prevalence of lifetime self-reported STDs was expressed as a percentage and it was calculated as the number of participants with prior history of STD divided by the total number of patients included in this analysis. Descriptive analysis was carried out and Chi-square statistic was used to assess differences of categorical data. The magnitude of the associations between explanatory variables and the outcome of interest – history of STD – was estimated by the odds ratio (OR) with 95% confidence interval (95% CI). The independent effect of potential exposure variables was assessed by multivariate analysis using logistic regression. All variables with *p*-values equal to or less than 0.20 in the univariate analysis were included in multivariate modeling. A backward deletion strategy was applied and those variables with *p*-values equal to or less than 0.05 remained in the final model. SAS System[®] was used for data analysis and Paradox[®] Windows for the database management.

Results

Among the 2475 patients interviewed, 2145 (86.7%) were sexually active (ever) and were included in this analysis. Among these, most participants were under treatment in outpatient centers (CAPS) (64.7%), older than 40 years old (55.5%), white (51.5%), female (52.6%), and 62.3% were single, divorced or widowed (Table 1). Almost half of the sample had less than five years of schooling (48.3%) and for 18.0% of patients the family income was lower than the Brazilian minimum wage at the time of the interview (<US 200). Four percent were living on the streets or in public shelters while 18.5% had a history of homelessness. Previous psychiatric hospitalization was reported by 57.5% of the patients. Psychoses, depression with psychotic symptoms and bipolar disorders were the most common psychiatric diagnoses (55.8%), while substance use disorder represented 7.9%. The remaining 36.3% consisted of anxiety, depression and others, mostly personality disorders, dementias and diagnoses under investigation. Only 30% of the participants had been previously tested for HIV. While 73.6% had a history of cigarette smoking, 68.1% consumed alcohol, 27.9% reported use of any illicit drug, and 3.3% reported injection drug use. Almost 32.0% of the participants reported having had sex under the influence of alcohol and/or drugs, 91.6% reported lifetime unprotected sex, and 28.2% reported having had a partner who refused to use condom at least once.

Table 1 – Descriptive characteristics among the 2145 eligible psychiatric patients.

Characteristics	n (%)
Sociodemographic	
Age (>40 years old)	1190 (55.5)
Skin color (white)	1105 (51.5)
Gender (female)	1129 (52.6)
Marital status (single, divorced or widowed)	1337 (62.3)
Schooling (<5 years)	1036 (48.3)
Family income (<1 minimum wage ^a)	305 (18.0)
Living in unstable situation ^b	84 (3.9)
Lifetime homelessness	397 (18.5)
Clinical	
Type of recruitment center	
CAPS	1387 (64.7)
Hospital	758 (35.3)
Previous psychiatric hospitalization	1228 (57.5)
Main psychiatric diagnosis	
Psychoses, DPS, BPD	1197 (55.8)
Substance use	169 (7.9)
Others	779 (36.3)
Previous HIV testing	646 (30.1)
Behavioral	
Lifetime cigarette smoking	1574 (73.6)
Lifetime alcohol use	1461 (68.1)
Lifetime illicit drug use	599 (27.9)
Lifetime injection drug use	70 (3.3)
Sex under the influence of alcohol and/or drugs	672 (31.6)
Lifetime unprotected sex ^c	1944 (91.6)
Partner refused to use condom anytime	588 (28.2)
Exchange money and/or drugs for sex	642 (30.1)
Number of sexual partners	
Only one	496 (24.8)
2-9	969 (48.4)
10 or more	536 (26.8)
Age of first sexual intercourse (years old)	
<14	306 (15.0)
14-18	1125 (55.0)
>18	616 (30.0)
Lifetime verbal violence	1533 (71.6)
Lifetime physical violence	1279 (59.8)
Lifetime sexual violence	460 (21.6)
Lifetime incarceration	584 (27.2)
Good HIV/AIDS knowledge	1249 (58.6)
High HIV risk self-perception	415 (20.1)

DPS, depression with psychotic symptoms; BPD, bipolar disorder.

^a Minimum wage = US\$200.

^b Living at the streets or public shelters.

^c Not always uses condoms in all practices.

Thirty percent exchanged money or drugs for sex, 26.8% had more than 10 sexual partners, and 15.0% had their first sexual intercourse before 14 years old. There was a high proportion of lifetime verbal, physical or sexual violence (71.6%, 59.8% and 21.6%, respectively) and incarceration (27.2%). Finally, 58.6% of the participants had good HIV/AIDS knowledge and 20.1% perceived themselves as being at high risk for HIV/AIDS.

More than one quarter of the participants (25.8%) reported a history of STD. Lifetime STD symptoms, characterized in three main syndromic groups, were genital discharges (35.8%), genital ulcers (14.2%), and warts (7.6%). Four percent of the participants received a medical diagnosis of syphilis, 3.7%

Table 2 – Characteristics and prevalence of lifetime sexually transmitted diseases among the 2145 eligible psychiatric patients.

Characteristics	n (%)
Self-reported STD ^a	554 (25.8)
<i>Syndromic groups</i>	
Genital discharge	761 (35.8)
Genital ulcers	301 (14.2)
Genital warts	161 (7.6)
<i>Medical diagnosis</i>	
Syphilis	79 (3.8)
Herpes	79 (3.7)
Chancroid	50 (2.4)
Gonorrhea	227 (10.7)
Chlamydia	66 (3.1)
Lymphogranuloma venereum	16 (0.8)
Condyloma	67 (3.2)

^a Sexually transmitted diseases.

of herpes, 2.4% of chancroid, 10.7% of gonorrhea, 3.1% of chlamydia, 0.8% of lymphogranuloma, and 3.2% of condyloma (Table 2).

Univariate analysis (Table 3) indicated that the prevalence of self-reported STD was almost two fold higher among men and among those over 40 years old ($p < 0.001$). The odds ratio of self-reported STD was two times higher for those living on the streets or in public shelters than among those living in houses, apartments or hospitals. Also, a previous history of homelessness was statistically associated with past STD. The participants who had a previous psychiatric hospitalization and those who were tested for HIV were more likely to have an STD history ($p < 0.002$). Likewise, the use of cigarette, alcohol, illicit drugs and injected drugs were statistically associated with self-reported STD. Several high risk sexual behavior characteristics were also associated with STD history ($p < 0.001$), as follows: sex under the influence of alcohol/drugs, previous unprotected sex, partner refusing to use condom and receiving or offering money/drugs for sex. In addition, the odds ratio of STD history was more than three times higher among those with 10 or more sexual partners and more than two times higher among those who had their first sexual intercourse before 14 years old. Moreover, patients with a history of verbal, physical or sexual violence and those with past incarceration were also more likely to report previous STD ($p < 0.001$). Finally, the univariate analysis indicated that the prevalence of self-reported STDs was higher among patients who perceived themselves as being at high risk for HIV/AIDS ($p < 0.001$).

The multivariate analysis (Table 3) indicated 12 variables which were independently associated ($p < 0.05$) with self-reported STD: being 30–39 years old and older than 40 years old; history of homelessness; lifetime alcohol or illicit drugs use; lifetime unprotected sex; partners' refusal to use condom at least once; exchanging money/drugs for sex; having more than 10 sexual partners; early age of first sexual intercourse (<14 years old and between 14 and 18 years old); history of verbal or physical violence; and those who perceived themselves as being at high risk for HIV/AIDS.

Discussion

Our results indicate that more than a quarter (25.8%) of participants in this national representative sample of psychiatric patients under care in Brazil had a lifetime history of at least one STD. This result is of public health concern and it is corroborated by other studies among psychiatric patients. In a study of patients with chronic mental illnesses, designed to assess the efficacy of an intervention for HIV prevention, in the USA, the prevalence of self-reported STD was 38%.¹⁵ However, their sample was only hospital-based and with psychiatric patients with substance use disorders, which can partially explain their higher proportion of history of STD. King et al.²¹ demonstrated 40.4% of self-reported STD among patients attending an out-patient university-based psychiatric clinic which included diverse psychiatric diagnoses in USA. Studies carried out in India among psychiatric patients seeking care for substance use disorder²² and among patients hospitalized in a general psychiatric hospital²³ found STD rates of 25.1% and 16.2%, respectively. Both studies used serologic tests to investigate syphilis, chlamydia, HIV and hepatitis B.

The prevalence of previous STD found in our study is high when compared with other Brazilian non-psychiatric populations. Many authors studied the prevalence of STDs in the general population, ranging from 5.2% to 21.3%.^{3,24,25} Among pregnant women the prevalence of STDs ranged from 6.7% to 42.0%.^{3,26} Population with well-known higher risk sexual behavior were assessed and STDs prevalence ranged from 22.0% to 51.0% when clients of STD clinics were studied.^{3,27}

It should be pointed out the difficulty in comparing data across studies that vary widely in their sampling and assessment methods. Nonetheless, it is important to notice that the prevalence of STDs was higher when compared to our study only when the population studied had well-known higher risk sexual behavior, such as STD clinics clients, or when serologic testing for STD increased the detection rates of the addressed population.

Our study identified several factors associated with history of STD among this representative sample of psychiatric patients. These factors may serve as potential indicators of risk behaviors suitable to preventive actions in these services. The prevalence of self-reported STD was higher among those older than 30 years old. In contrast to higher risk of STD among younger populations described by other authors,^{2,3} the association found with older age groups in our study can be explained by a lifetime cumulative exposure, usually detected in prevalence as opposed to incidence studies. A similar result was found in a study conducted among 641 truck drivers in Brazil.⁴ As demonstrated by other studies, substance use is a strong predictor of STD in several populations, including patients with mental illnesses.^{4,22,24,27} Carey et al.²² demonstrated that psychiatric patients in risk for alcohol and drug abuse had higher prevalence of STDs. In a controlled trial²⁸ addressing HIV prevention, patients who received a substance use reduction intervention were more likely than controls to reduce the number of sexual partners and to endorse stronger intention to use condoms. In our study, alcohol and illicit drug use were independently associated with STD history, while use of tobacco and injected drugs was only found to be

Table 3 – Univariate and multivariate logistic analysis of self-reported STD^a among the 2145 eligible psychiatric patients.

Characteristics	Total	STD ^b n (%)	Univariate analysis		Multivariate analysis	
			Odds ratio (95% CI)	X ² (p-value)	Odds ratio ^c (95% CI)	X ² (p-value)
Sociodemographic						
<i>Age (years old)</i>						
18–29	369	68 (18.4)	1.00		1.00	
30–39	586	145 (24.7)	1.46 (1.05–2.01)	5.18 (0.023)	1.83 (1.27–2.62)	10.75 (0.001)
>40	1190	341 (28.7)	1.78 (1.33–2.38)	14.96 (<0.001)	3.04 (2.16–4.30)	39.94 (<0.001)
<i>Skin color</i>						
Non-white	1040	285 (27.4)	1.00		–	
White	1105	269 (24.3)	0.85 (0.70–1.03)	2.62 (0.106)		
<i>Gender</i>						
Female	1129	251 (22.2)	1.00		–	
Male	1016	303 (29.8)	1.49 (1.22–1.81)	16.08 (<0.001)		
<i>Marital status</i>						
Married or in union	808	197 (24.4)	1.00		–	
Single, divorced or widowed	1337	357 (26.7)	1.13 (0.92–1.38)	1.41 (0.234)		
<i>Schooling (years)</i>						
≥5	1109	306 (27.6)	1.00		–	
<5	1036	248 (23.9)	0.83 (0.68–1.00)	3.73 (0.053)		
<i>Family income</i>						
≥1 minimum wage ^d	1386	346 (25.0)	1.00		–	
<1 minimum wage	305	88 (28.9)	1.22 (0.92–1.60)	1.98 (0.159)		
<i>Living situation</i>						
House, apartment, hospital	2058	523 (25.4)	1.00		–	
Unstable situation ^e	84	30 (35.7)	1.63 (1.03–2.58)	4.47 (0.035)		
<i>Lifetime homelessness</i>						
No	1748	394 (22.5)	1.00		1.00	
Yes	397	160 (40.3)	2.32 (1.84–2.92)	53.28 (<0.001)	1.35 (1.03–1.76)	4.74 (0.029)
Clinical						
<i>Type of recruitment center</i>						
CAPS	1387	355 (25.6)	1.00		–	
Hospital	758	199 (26.3)	1.03 (0.85–1.27)	0.11 (0.739)		
<i>Previous psychiatric hospitalization</i>						
No	908	204 (22.5)	1.00		–	
Yes	1228	348 (28.3)	1.36 (1.12–1.67)	9.39 (<0.002)		
<i>Main psychiatric diagnosis</i>						
Others	779	177 (22.7)	1.00		–	
Psychoses, DPS, BPD	1197	328 (27.4)	1.28 (1.04–1.58)	5.42 (0.020)		
Substance use	169	48 (28.4)	1.35 (0.93–1.96)	2.46 (0.117)		
<i>Previous HIV testing</i>						
No	1499	353 (23.6)	1.00		–	
Yes	646	201 (31.1)	1.47 (1.19–1.80)	13.48 (<0.001)		
<i>Lifetime cigarette smoking</i>						
No	564	111 (19.7)	1.00		–	
Yes	1574	439 (27.9)	1.58 (1.25–2.00)	14.64 (<0.001)		
<i>Lifetime alcohol use</i>						
No	684	121 (17.7)	1.00		1.00	
Yes	1461	433 (29.6)	1.96 (1.56–2.46)	34.70 (<0.001)	1.37 (1.05–1.79)	5.43 (0.020)
<i>Lifetime illicit drug use</i>						
No	1546	323 (20.9)	1.00		1.00	
Yes	599	231 (38.6)	2.38 (1.94–2.92)	70.35 (<0.001)	1.65 (1.27–2.14)	14.06 (<0.001)
<i>Lifetime injected drug use</i>						
No	2065	515 (24.9)	1.00		–	
Yes	70	37 (52.9)	3.37 (2.09–5.45)	27.51 (<0.001)		
<i>Sex under the influence of alcohol/drugs</i>						
No	1455	313 (21.5)	1.00		–	
Yes	672	237 (35.3)	1.99 (1.62–2.43)	45.35 (<0.001)		
<i>Lifetime unprotected sex^f</i>						
No	179	27 (15.1)	1.00		1.00	
Yes	1944	524 (27.0)	2.08 (1.36–3.17)	12.02 (<0.001)	1.80 (1.13–2.84)	6.24 (0.013)
<i>Partner refused to use condom anytime</i>						
No	1496	356 (23.8)	1.00		1.00	
Yes	588	183 (31.1)	1.45 (1.17–1.79)	11.81 (<0.001)	1.37 (1.08–1.73)	6.75 (0.009)

Table 3 (Continuación)

Characteristics	Total	STD ^b n (%)	Univariate analysis		Multivariate analysis	
			Odds ratio (95% CI)	X ² (p-value)	Odds ratio ^c (95% CI)	X ² (p-value)
<i>Exchange money/drugs for sex</i>						
No	1488	299 (20.1)	1.00		1.00	
Yes	642	251 (39.1)	2.55 (2.08–3.13)	84.51 (<0.001)	1.48 (1.15–1.89)	9.56 (0.002)
<i>Number of sexual partners</i>						
Only one	496	82 (16.5)	1.00		1.00	
2–9	969	222 (22.9)	1.50 (1.13–1.97)	8.05 (0.011)	1.09 (0.80–1.49)	0.27 (0.600)
10 or more	536	219 (40.9)	3.49 (2.60–4.68)	69.89 (<0.001)	1.72 (1.20–2.46)	8.88 (0.003)
<i>Age of first sexual intercourse</i>						
>18 years old	616	117 (19.0)	1.00		1.00	
14–18 years old	1125	314 (27.9)	1.65 (1.30–2.10)	16.80 (<0.001)	1.32 (1.01–1.72)	4.08 (0.043)
<14 years old	306	104 (34.0)	2.20 (1.61–3.00)	24.63 (<0.001)	1.53 (1.07–2.19)	5.32 (0.021)
<i>Lifetime verbal violence</i>						
No	607	106 (17.5)	1.00		1.00	
Yes	1533	447 (29.2)	1.95 (1.54–2.47)	31.02 (<0.001)	1.52 (1.16–1.99)	9.07 (0.003)
<i>Lifetime physical violence</i>						
No	861	158 (18.4)	1.00		1.00	
Yes	1279	394 (30.8)	1.98 (1.61–2.44)	41.68 (<0.001)	1.40 (1.10–1.78)	7.28 (0.007)
<i>Lifetime sexual violence</i>						
No	1675	396 (23.6)	1.00		–	
Yes	460	155 (33.7)	1.64 (1.31–2.05)	19.04 (<0.001)		
<i>History of incarceration</i>						
No	1561	337 (21.6)	1.00		–	
Yes	584	217 (37.2)	2.15 (1.75–2.64)	53.75 (<0.001)		
<i>HIV/AIDS knowledge</i>						
Good	1249	340 (27.2)	1.00		–	
Poor	882	211 (23.9)	0.84 (0.69–1.03)	2.93 (0.087)		
<i>HIV risk self-perception</i>						
Some risk. None or do not know	1650	389 (23.6)	1.00		1.00	
High	415	147 (35.4)	1.78 (1.41–2.24)	24.20 (<0.001)	1.56 (1.21–2.00)	11.92 (<0.001)

DPS, depression with psychotic symptoms; BPD, bipolar disorder.

^a Sexually transmitted diseases.

^b Number and proportion of self-reported STD.

^c Hosmer–Lemeshow test: X² = 9400; df = 8; p = 0.3474.

^d Minimum wage = US\$200.

^e Living at the streets or public shelters.

^f Not always using condoms in all practices.

associated in univariate analysis. The association of substance use with STD can be explained by poor access to prevention and care and also due to a potential impairment of judgment to practise safe sex, which may lead to poor decisions regarding sexual behavior. Effective alcohol and drug prevention and treatment may reduce sexual risk behavior, and consequently the STDs prevalence.

We have also identified important behavioral factors associated with an increased likelihood of STDs: lifetime unprotected sex, partner's refusal to use condom, exchanging money and/or drugs for sex, having multiple sexual partners, and early age of first sexual intercourse. High risk sexual behaviors among patients with mental illnesses have been highlighted by other investigators. Vanable et al.¹⁵ found that patients with mental illness receiving care in a hospital with a previous STD practised unprotected sex more often. Carey et al.,²³ also studying psychiatric inpatients, indicated that lifetime STD was statistically associated with exchanging money for sex and having multiples partners. King et al.²¹ demonstrated that STD was associated with exchanging sex for drugs in a sample of psychiatric outpatients in USA. This

high risk sexual behavior was also described in Brazil, in a sample of patients of a psychiatric hospital in the Southeast Region,²⁹ where 68.2% of patients had previous unprotected sex and 2.6% exchanged sex for money or drugs.

In addition to behavioral and substance use correlates, our data also point out the increased social vulnerability profile of these psychiatric patients, including suffering violence and presenting higher rates of homelessness. Those who reported having suffered verbal and physical violence, and those with a history of homelessness had statistically higher proportions of STDs. Our findings are consistent with recent data published by Oliveira³⁰ which indicates that verbal and physical violence against women and men were independently associated with a previous diagnosis of STD among psychiatric patients. In addition, suffering violence was also associated with lifetime homelessness in the same study. People living with mental illnesses are frequently exposed to many factors that leave them in extreme vulnerable conditions, especially those who have experienced homelessness situations. They are often subject to violence, and are more prone to alcohol and drug abuse as well as

exchanging sex as a means of obtaining food, shelter, money or drugs. All these factors exacerbate the already fragile individuals and it certainly leaves them more exposed to STDs.

On the other hand, patients who perceived themselves at a higher risk for acquisition of HIV presented higher prevalence of history of STD. This result may actually reflect a dynamics between risk behavior, HIV awareness, and the perception of need and access to care. As proposed by other authors,¹⁵ it seems that diagnosis and treatment of STDs may increase HIV-related awareness and knowledge, but they do not necessarily lead to changes in sexual risk behavior. This is consistent with our data, since in our sample, patients with a STD history reported higher rates of sexual risk behavior in comparison to the patients without a previous STD history.

Our study has some limitations: STDs were self-reported, and since they are often asymptomatic, it could lead to a potential underestimation of their prevalence. However, the proportion found was considerably high, which actually reinforces the magnitude of the problem. In addition, reliability of the data was adequate as shown.¹⁸ Also, because this is a cross-sectional study, we can only indicate statistical correlations rather than direct causal effect between potential explanatory variables and a history of STDs. Nevertheless, our findings are consistent with the literature and have important clinical and public health implications.

In conclusion, as shown, most factors associated with STDs among this representative sample of patients with mental illnesses in Brazil are potentially modifiable behavioral factors. As patients are admitted for care in these mental health services, a good opportunity for STD and risk behavioral screening is created. Although programs designed to integrate STD prevention with the psychiatric care have been successfully implemented,²⁸ only eight (31.0%) of the 26 centers included in our study had specific educational programs for STDs,³¹ indicating a clear lack of opportunity for prevention and care for STDs. Screening for STDs, including HIV, hepatitis B and C, and syphilis should be part of the routine care in these centers. In addition, counseling, behavioral modification intervention and referral for treatment and follow-up of those with positive results should be implemented on a routine basis.

Conflicts of interest

The authors declare no conflicts of interest.

Acknowledgments

This work was carried out by the Federal University of Minas Gerais with technical and financial support of the Ministry of Health/Secretariat of Health Surveillance/Department of STD, AIDS and Viral Hepatitis through the Project of International Technical Cooperation 914/BRA/1101 between the Brazilian Government and the United Nations Educational, Scientific and Cultural Organization – UNESCO.

REFERENCES

1. World Health Organization. Regional HIV/STI Plan for the Health Sector 2006–2015; November, 2005. Available from: <http://www.paho.org/aids> [accessed on 01.11.12].
2. Centers for Disease Control and Prevention. Sexually Transmitted Disease Surveillance 2010. Atlanta: U.S. Department of Health and Human Services; 2011. Available from: <http://www.cdc.gov/std/stats> [accessed on 01.11.12].
3. Brazil, Ministry of Health. Secretaria de Vigilância em Saúde. Programa Nacional de DST e Aids. Prevalências e frequências relativas de Doenças Sexualmente Transmissíveis (DST) em populações selecionadas de seis capitais brasileiras, 2005. Brasília: Ministério da Saúde; 2008. p. 224.
4. Teles AS, Matos MA, Caetano KAA, et al. Comportamentos de risco para doenças sexualmente transmissíveis em caminhoneiros no Brasil. *Rev Panam Salud Publica*. 2008;24:25–30.
5. Brazil, Ministry of Health. Secretaria de Vigilância em Saúde. Programa Nacional de DST e Aids. Manual de Controle das Doenças Sexualmente Transmissíveis/Ministério da Saúde, Secretaria de Vigilância em Saúde, Programa Nacional de DST e Aids. Brasília: Ministério da Saúde; 2005.
6. Cohen MS. Sexually transmitted diseases enhance HIV transmission: no longer a hypothesis. *Lancet*. 1998;351: 5–7.
7. Wu H, Wu P-Y, Li S-Y, et al. Maximising the potential of voluntary counselling and testing for HIV: sexually transmitted infections and HIV epidemiology in a population testing for HIV and its implications for practice. *Sex Transm Infect*. 2012;88:612–6.
8. Meade CS, Sikkema KJ. HIV risk behavior among adults with severe mental illness: a systematic review. *Clin Psychology Rev*. 2005;25:433–57.
9. Guimarães MDC, McKinnon K, Campos LN, Melo APS, Wainberg M. HIV risk behavior of psychiatric patients with mental illness: a sample of Brazilian patients. *Rev Bras Psiquiatr*. 2010;32:351–60.
10. Barbosa RM, Koyama MA, Grupo de Estudos em População, Sexualidade e Aids. Sexual behavior and practices among men and women, Brazil 1998 and 2005. *Rev Saude Publica*. 2008;42:21–33.
11. Brazil, Ministry of Health. Pesquisa sobre Comportamento, Atitudes e Práticas Relacionadas às DST e Aids na População Brasileira; 2009. Available from: <http://www.aids.gov.br/data/Pages/LUMISE77B47C8ITEMIDE56B57FF739940039DB3112DF74DE47FPTBRIE.htm> [cited 23.07.06].
12. Bastos FI, Bertoni N, Hacker MA, Grupo de Estudos em População, Sexualidade e Aids. Drug and alcohol use: main findings of a national survey, Brazil 2005. *Rev Saude Publica*. 2008;42:109–17.
13. Lagios K, Deane FP. Severe mental illness is a new risk marker for bloodborne viruses and sexually transmitted infections. *Aust N Z J Public Health*. 2007;31:562–6.
14. Campos LN, Guimarães MDC, Carmo RA, et al. HIV, syphilis, and hepatitis B and C prevalence among patients with mental illness: a review of the literature. *Cad Saúde Pública*. 2008;24:607–20.
15. Vanable PA, Carey MP, Carey KB, Maisto SA. Differences in HIV-related knowledge, attitudes, and behavior among psychiatric outpatients with and without a history of a Sexually Transmitted Infection. *J Prev Interv Community*. 2007;33:79–94.
16. Guimarães MDC, Campos LN, Melo APS, Carmo RA, Machado CJ, Acúrcio FA, et al. Prevalence of HIV, syphilis, hepatitis B and C among adults with mental illness: a multicenter study in Brazil. *Rev Bras Psiquiatr*. 2009;31:43–7.

17. Guimarães MDC, Oliveira HN, Campos LN, et al. Reliability and validity of a questionnaire on vulnerability to sexually transmitted infections among adults with chronic mental illness. *Rev Bras Psiquiatr.* 2008;30:55-9.
18. Oliveira HN, Machado CJ, Campos LN, Acúrcio FA, Guimarães MD. Avaliação da confiabilidade de entrevista aplicada a indivíduos com transtornos mentais: Projeto PESSOAS. *Rev Med Minas Gerais.* 2008;18:40-5.
19. Ministry of Health-Brazil. National program for STDs and AIDS. Positive response: Experiences of the Brazilian AIDS program. Brasília, Brazil: Ministry of Health of Brazil; 2005. Available from: http://www.aids.gov.br/final/biblioteca/resposta/resposta_2005.pdf [cited 23.07.06].
20. Melo APS, Cesar CC, Acúrcio FA, et al. Individual and treatment setting predictors of HIV/AIDS knowledge among psychiatric patients and their implications in a national multisite Study in Brazil. *Community Ment Health J.* 2010;46:505-16.
21. King C, Feldman J, Waithaka Y, et al. Sexual risk behaviors and sexually transmitted infection prevalence in an outpatient Psychiatry Clinic. *Sex Transm Dis.* 2008;35:877-82.
22. Carey MP, Ravi V, Chandrac PS, Desai A, Neald DJ. Screening for sexually transmitted infections at a deaddictions service in South India. *Drug Alcohol Depend.* 2006;82:127-34.
23. Carey MP, Ravi V, Chandrac PS, Desai A, Neald DJ. Prevalence of HIV, Hepatitis B, Syphilis, and chlamydia among adults seeking treatment for a mental disorder in Southern India. *AIDS Behav.* 2007;11:289-97.
24. Codes JS, Cohen DA, Melo NA, et al. Detecção de doenças sexualmente transmissíveis em ambientes clínicos e não clínicos na Cidade de Salvador, Bahia, Brasil. *Cad Saúde Pública.* 2006;22:325-34.
25. Oliveira FA, Pflieger V, Lang K, et al. Sexually transmitted infections, bacterial vaginosis, and candidiasis in women of reproductive age in rural Northeast Brazil: a population-based study. *Mem Inst Oswaldo Cruz.* 2007;102:751-6.
26. Lima LHM, Viana MC. Prevalence and risk factors for HIV, syphilis, hepatitis B, hepatitis C, and HTLV-I/II infection in low-income postpartum and pregnant women in Greater Metropolitan Vitória, Espírito Santo State, Brazil. *Cad Saúde Pública.* 2009;25:668-76.
27. Cook R, May S, Harrison LH, et al. High prevalence of sexually transmitted diseases in young women seeking HIV testing in Rio de Janeiro, Brazil. *Sex Transm Dis.* 2004;31:67-72.
28. Carey MP, Carey KB, Maisto SA, Gordon CM, Schroder KE, Vanable PA. Reducing HIV-risk behavior among adults receiving outpatient psychiatric treatment: results from a randomized controlled trial. *J Consult Clin Psychol.* 2004;72:252-68.
29. Almeida RC, Pedroso ERP. Vulnerabilidade e exposição a marcadores sorológicos do Vírus da Imunodeficiência Humana, hepatites B e C, vírus linfotrópico de células T humanas e sífilis em pacientes psiquiátricos internados em hospital público. *Rev Méd Minas Gerais.* 2004;14:244-50.
30. Oliveira HN. Violência contra indivíduos com transtorno mental: Projeto Pessoas [thesis]. Belo Horizonte (MG): Universidade Federal de Minas Gerais; 2012.
31. Melo APS, Acúrcio FA, Cherchiglia ML, Veloso CCG, Guimarães MDC. Avaliação de serviços de saúde mental: assistência e prevenção às doenças sexualmente transmissíveis no contexto do Projeto Pessoas. *Rev Méd Minas Gerais.* 2007;17:233-9.