



The Brazilian Journal of INFECTIOUS DISEASES

www.elsevier.com/locate/bjid



Original article

Modulation of drug resistance and biofilm formation of *Staphylococcus aureus* isolated from the oral cavity of Tunisian children



Tarek Zmantar^a, Rihab Ben Slama^a, Kais Fdhila^a, Bochra Kouidhi^b, Amina Bakhrouf^a, Kamel Chaieb^{c,*}

^a Faculty of Pharmacy, Laboratory of Analysis, Treatment and Valorization of Pollutants of the Environment and Products, Monastir University, Tunisia

^b College of Applied Medical Sciences, Medical Laboratory Technology Department, Yanbu al Bahr, Taibah University, Al Madinah Al Monawarah, Saudi Arabia

^c College of Sciences, Biology Department, Yanbu al Bahr, Taibah University, Al Madinah Al Monawarah, Saudi Arabia

ARTICLE INFO

Article history:

Received 24 March 2016

Accepted 26 October 2016

Available online 1 December 2016

Keywords:

Antibacterial

Staphylococcus aureus

Biofilm

Antibiotics

Synergetic

Lactobacillus

ABSTRACT

Objectives: This study aims to investigate the antimicrobial and the anti-biofilm activities of *Lactobacillus plantarum* extract (LPE) against a panel of oral *Staphylococcus aureus* ($n=9$) and *S. aureus* ATCC 25923. The *in vitro* ability of LPE to modulate bacterial resistance to tetracycline, benzalchonium chloride, and chlorhexidine were tested also.

Methods: The minimum inhibitory concentrations (MICs) and the minimal bactericidal concentrations of *Lactobacillus plantarum* extract, tetracycline, benzalchonium chloride and clohrhexidine were determined in absence and in presence of a sub-MIC doses of LPE (1/2 MIC). In addition, the LPE potential to inhibit biofilm formation was assessed by microtiter plate and atomic force microscopy assays. Statistical analysis was performed on SPSS v. 17.0 software using Friedman test and Wilcoxon signed ranks test. These tests were used to assess inter-group difference ($p < 0.05$).

Results: Our results revealed that LPE exhibited a significant antimicrobial and anti-biofilm activities against the tested strains. A synergistic effect of LPEs and drug susceptibility was observed with a 2–8-fold reduction.

Conclusion: LPE may be considered to have resistance-modifying activity. A more detailed investigation is necessary to determine the active compound responsible for therapeutic and disinfectant modulation.

© 2016 Sociedade Brasileira de Infectologia. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

* Corresponding author.

E-mail address: chaieb_mo@yahoo.fr (K. Chaieb).

<http://dx.doi.org/10.1016/j.bjid.2016.10.009>

1413-8670/© 2016 Sociedade Brasileira de Infectologia. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Staphylococci are important causes of infections associated with various devices. *Staphylococcus aureus* is one of the most frequent human pathogen associated to medical implant.¹ It has been isolated from parotitis,² gingival pockets,³ periodontitis,⁴ carious lesions,⁵ and gingivitis.⁶ Orthodontic and other oral appliances may act as reservoir of resistant opportunistic *S. aureus*.⁷

S. aureus has the capacity to adhere to various medical devices and form biofilm.⁸ Biofilm is associated to the *ica* gene encoding for polysaccharide intercellular adhesion (PIA)⁹ which leads to increased bacterial drug resistance compared to planktonic cells.^{10,11}

S. aureus is able to acquire resistance to antibiotics and to antiseptic agents via efflux pumps (TetK, *msrA* transporters) systems, which export certain tetracyclines and macrolides molecules outside the cells. On the other hand, some multidrug resistance (MDR) proteins like NorA and QacA confer resistance to a wide range of structurally unrelated antiseptics.¹² In clinical practice, chlorhexidine (CHX) and quaternary ammonium compounds (QAC) are the most frequently used disinfectants to reduce and prevent the spread of pathogens. Multidrug resistance pumps have been recognized as mediators of a number of commonly used ammonium compounds and detergents.¹³ However, the use of these disinfectants in hospitals may contribute to the rise of disinfectant-resistant bacteria^{14,15} due to QAC-resistant genes (*qacA*, *qacB*, and *qacC*), which have been identified in several staphylococcal species.¹⁶⁻¹⁸

In this study, the antibacterial activity of *Lactobacillus plantarum* extract (LPE) was investigated. In addition, the ability of LPE to modulate the susceptibility of *S. aureus*, isolated from the oral cavity of Tunisian children, to tetracycline (TET), benzalchonium chloride (BC) and clohrhexidine (CHX) was tested. Secondly, the effects of disinfectant and antibiotic associated with LPE were tested for biofilm inhibition of *S. aureus* to polystyrene and glass using microtiter plate and atomic force microscopy assays, respectively.

Material and methods

Antibacterial activity of *L. plantarum*

L. plantarum strain was isolated from Tunisian traditional fermented milk (ricotta cheese), identified with a conventional method using Api-50 CHL system (BioMerieux, Marcy-l'Étoile, France) and by polymerase chain reaction (PCR) technique using *L. plantarum*-specific primers: forward IDL04F: 5'-AGGGTGAAGTCGTAACAAGTAGCC-3' and reverse IDL62R 5'-CTAGTGGTAACAGTTGATTAATAACTGC-3', giving a product size of about 428 bp as described previously.¹⁹

Cell-free supernatant was obtained by centrifuging (10,000 × *g* for 10 min at 4 °C) *L. plantarum* culture, grown in MRS broth for 16 h at 30 °C. The supernatant was adjusted at neutral pH and filter-sterilized (0.22 μm, Millipore). The obtained *L. plantarum* extract (LPE) was conserved at 4 °C until use.

Antibacterial activity of LPE was tested against nine strains (Table 1) isolated from the oral cavity of Tunisian children and *S. aureus* ATCC 25923 using the broth microdilution method.^{20,21} The LPE was tested alone or in combination with antibiotics.

Microorganisms

The *S. aureus* strains (*n*=9) used in this study were isolated from the oral cavity of Tunisian children from the dental clinic of dentistry (Monastir, Center of Tunisia).

The criteria for inclusion were: no antibiotic treatment four weeks prior to sampling, no use of mouth rinses or any other preventive measure that might involve exposure to antimicrobial agents, and no systemic disease. A sterile swab was used for sample collection from the oral cavity of each patient. After incubation (24 h at 37 °C), swabs were plated on blood agar plates supplemented with 5% sheep blood (24 h at 37 °C) and bacterial identification was achieved using conventional methods.

Minimal inhibitory concentration determination

The broth microdilution method was used to determine the minimum inhibitory concentration (MIC) of LPE (5–90%, v/v), BC (Acros organics, USA) (0–1024 μg/mL), CHX (Sigma-Aldrich, USA) (0–1024 μg/mL) and tetracycline (TET) (0–1024 μg/mL) according to the Clinical and Laboratory Standards Institute (2006) guidelines.²²

After 24 h of incubation, bacterial growth was evaluated by the presence of turbidity and a pellet on the well bottom. MIC was defined as the lowest concentration of the compound that had no macroscopically visible growth. All experiments were carried out three times.

Minimal bactericidal concentration determination (MBC)

To determine the MBC values, 10 μL of each well medium, with no visible growth was removed and inoculated in Muller Hinton agar plates. MBC was defined as the lowest concentration at which 99% of the bacteria were killed. Each experiment was repeated at least twice.²³

Modulation of *S. aureus* susceptibility to BC, CHX and TET by LPE

To determine the potential effect of LPE to modulate drug resistance of *S. aureus*, MICs of BC, CHX and TET (ranging from 0.5 to 2048 μg/mL) were determined alone and combined with a sub-MIC of LPE (1/2 MIC, v/v) using the microtiter plates assay.²⁴

Determination of anti-biofilms activity by microtiter plates assay

The anti-adhesion properties of LPE (ranging from 10% to 90%, v/v) to *S. aureus* strains were tested as previously described by Merritt et al.²⁵ Briefly, the bacterial culture was grown in Tryptone soy broth (TSB) at 37 °C for 24 h. Two microliters were disposed into each well of 96-well plates in the presence of

Table 1 – Minimum inhibitory and minimum bactericidal concentrations in $\mu\text{g/mL}$ of *Lactobacillus plantarum* extract alone or in combination with benzalchonium chloride, chlorhexidine, and tetracycline.

Strains	LPE (%v/v)		BC ($\mu\text{g/mL}$)		BC ($\mu\text{g/mL}$) + 1/2 MIC (LPE %, v/v) ^a		CHX ($\mu\text{g/mL}$)		CHX ($\mu\text{g/mL}$) + 1/2 MIC (LPE %, v/v) ^a		TET ($\mu\text{g/mL}$)		TET ($\mu\text{g/mL}$) + 1/2 MIC (LPE%, v/v) ^a	
	MIC	MBC	MIC	MBC	MIC	MBC	MIC	MBC	MIC	MBC	MIC	MBC	MIC	MBC
S1	20	60	8	16	4 (2)	8 (2)	256	512	128 (2)	512	8	16	4 (2)	8 (2)
S2	10	60	4	8	4	8	128	512	128	512	4	16	1(4)	16
S3	40	40	32	64	16 (2)	64	128	256	32 (4)	256	32	64	16 (2)	64
S4	20	30	2	16	2	16	64	128	64	64 (2)	1	8	1	8 (2)
S5	20	70	16	32	8 (2)	32	128	512	64	512	32	32	8 (4)	16 (2)
S6	50	80	16	64	4 (4)	32 (2)	64	512	8 (8)	256 (2)	16	32	16	32
S7	10	30	8	16	4 (2)	16	128	128	32 (4)	64 (2)	8	16	8	16
S8	20	20	8	16	4 (2)	16	256	256	32 (8)	256	16	32	2 (8)	32
S9	20	30	128	256	128	128 (2)	128	256	128	128 (2)	64	128	32 (2)	128
<i>S. aureus</i> ATCC25923	20	20	16	32	4 (4)	16 (2)	32	64	16 (2)	16 (4)	4	8	2 (2)	4 (2)

LPE, *Lactobacillus plantarum* extract; BC, benzalchonium chloride; CHX, chlorhexidine; TET, tetracycline; MIC, minimal inhibitory concentration; MBC, minimal bactericide concentration.

^a Fold reductions are given in parentheses.

198 µL of the medium supplemented with 2% glucose (w/v) containing LPE. Plates were incubated in aerobic conditions at 37 °C for 24 h. The anti-biofilm activity of TET, CHX, and BC ranging from 2 to 1024 µg/mL were determined against *S. aureus* strains with or without sub-MIC of the LPE (1/2 MIC, v/v), using the microtiter plates assay. After incubation, the plate was washed with phosphate buffer saline, stained with 100 µL of 1% (w/v) crystal violet and incubated at room temperature for 15 min. Then, the absorbance at 590 nm was determined using a microplate reader (D.E.E.D reader, Bio-Rad instrument). The mean absorbance (OD_{590nm}) of the sample was determined, and the percentage of inhibition obtained for each concentration of LPE and drugs was determined by the following formula:

$$\left[\frac{(\text{OD growth control} - \text{OD sample})}{\text{OD growth control}} \right] \times 100$$

Evaluation of LPE biofilm inhibition by atomic force microscopy (AFM)

To evaluate biofilm inhibition potency of LPE, *S. aureus* ATCC 25923 was grown in Tryptone soy broth (TSB) at 37 °C for 24 h and then diluted into the TSB medium (10⁵ cells/mL) supplemented with 2% glucose (w/v). For each concentration of the tested drugs (2–1024 µg/mL) and LPE (5–90%), 10 mL of bacterial suspension was incubated in a 6-well plate containing round glass cover slips for 24 h at 37 °C. After adhesion, the non-adhering bacteria were removed by rinsing the substrate four times with sterile PBS. Then the substrate was air dried and observed using AFM. All AFM images were then processed to analyze the biofilm formation behavior.

Statistical analysis

Data were analyzed using SPSS v. 17.0 software. The Friedman test, followed by the Wilcoxon signed ranks test were used to assess inter-group differences. In addition, statistical significance was set at *p*-value <0.05.

Results

Antimicrobial activity of LPE

As presented in Table 1, LPE demonstrated selective antimicrobial properties. Seven *S. aureus* strains (S3, S4, S5, S6, S1, S8 and S9) and *S. aureus* ATCC 25923 have a MICs value ranged from 20% to 50% (v/v). The remaining two strains (S2, S7) were more susceptible (MIC 10%). In addition, a variable MBCs (10–90%) (v/v) was observed according to the tested strain (Table 1). On the other hand, out of the nine oral *S. aureus* strains, eight (Table 1) were considered resistant to BC (MICs between 4 and 128 µg/mL) and only one strain (S4) was sensitive (MICs ≤ 2 µg/mL); all strains were resistant to CHX (MICs > 2 µg/mL). Moreover, only one strain (S4) was susceptible to TET (MIC < 2 µg/mL).

Modulation of drug resistance by LPE

Data presented in Table 1 showed that the presence of LPE at 1/2-MIC (v/v) in combination with BC resulted in a 4-fold reduction of MICs and MBCs of *S. aureus* ATCC 25923 and one oral strain (S6). Concerning the synergistic effect of CHX and LPE, there was an important diminution of MIC and MBC of CHX (2–8-fold potentiation) in six strains (Table 1). Furthermore, a 2–8-fold reduction of MIC (µg/mL) of TET against seven strains when combined with LPE was noticed (Table 1).

Biofilm inhibition by LPE

Biofilm formation of oral *S. aureus* strains was evaluated in 96 microtiter wells plate with LPE (ranging from 5% to 90%, v/v). Results were expressed as minimum biofilm inhibition concentration (BIC50 and BIC90). As presented in Table 2, BIC90 of LPE was reached at 54% LPE supplementation, for two oral strains (S4 and S7). While, BIC50 ranged from 28 to 77% (v/v) LPE supplementation (Table 2).

In addition, BC showed various effects on the development of *S. aureus* biofilms with BIC50 values ranging from 12 to 112 µg/mL for the tested strains. Moreover, the combination of BC and 1/2 MIC LPE resulted in a reduction of BIC50 against eight strains (Table 2). In the same way BIC90 was also reduced with the combination of BC and LPE.

Concerning the effect of CHX alone and in combination with LPE, BIC50 values ranged from 34–355 µg/mL for CHX for the tested strains (Table 2) whereas the combination of CHX and 1/2 MIC of LPE (v/v) exhibited a reduction of BIC50 and BIC90 against all the tested strains (Table 2).

Our results revealed that TET may reduce *S. aureus* attachment to polystyrene (BIC90 reach 30 µg/mL for S4). In addition, the combination of TET and 1/2 MIC of LPE exhibited a reduction of BIC50 and BIC90 against nine strains (Table 2).

Evaluation of LPE biofilm inhibition by AFM

AFM revealed that the tested strains were strongly biofilm positive and had the potency to adhere to glass surface (Fig. 1A). On the other hand, LPE decreased significantly the adherence to glass (Fig. 1D). In addition, the combination of LPE with TET, BC and CHX decreased significantly the biofilm formation (Fig. 1B–E). As presented in Fig. 1F, almost total biofilm inhibition was observed with 16 µg/mL of BC combined with 1/2 MIC of LPE % (v/v).

Discussion

A wide ranges of *L. plantarum* produce anti-microbial peptides with important effects against Gram-positive bacteria.²⁶ Lactic acid bacteria (LAB) are known to have bactericidal activity against pathogenic bacteria. For that, they develop many mechanisms like hydrogen peroxide production causing denaturation of many enzymes and peroxidation of membrane leading to increased permeability.²⁷ Also, the LAB are able to produce bacteriocins which have a bactericidal effect on other species.²⁸ In order to find a natural source able to

Table 2 – Anti-biofilm effect of *Lactobacillus plantarum* extract alone or in combination with benzalchonium chloride, chlorhexidine, and tetracycline against oral staphylococci.

Strains	Inhibition of biofilm development (%)													
	LPE (% v/v)		BC (µg/mL)		BC (µg/mL) + 1/2 MIC (LPE %, v/v)		CHX (µg/mL)		CHX (µg/mL) + 1/2 MIC (LPE %, v/v)		TET (µg/mL)		TET (µg/mL) + 1/2 MIC (LPE %, v/v)	
	BIC50	BIC90	BIC50	BIC90	BIC50	BIC90	BIC50	BIC90	BIC50	BIC90	BIC50	BIC90	BIC50	BIC90
S1	32	84	16	84	16	80	50	510	32	62	59	220	10	190
S2	63	81	29	77	15	63	34	90	20	111	10	514	2	126
S3	77	93	30	90	14	60	355	532	126	321	130	321	83	319
S4	28	54	12	64	9	8	60	520	35	132	18	30	16	30
S5	31	92	20	90	8	85	277	556	19	120	29	421	11	216
S6	54	84	30	71	30	67	132	455	60	321	16	120	16	120
S7	31	54	21	16	7	5	66	432	40	509	27	37	11	23
S8	48	75	42	62	16	35	126	300	54	129	31	65	19	44
S9	29	51	112	200	22	14	133	266	65	113	34	73	17	51
<i>S. aureus</i> ATCC 25923	38	82	24	1015	12	27	46	64	26	72	16	82	9	46

LPE, *Lactobacillus plantarum* extract; BIC50, minimum biofilm inhibition concentration that showed 50% inhibition; BIC90, minimum biofilm inhibition concentration that showed 90% inhibition; BC, benzalchonium chloride; CHX, chlorhexidine; TET, tetracycline.




Fig. 1 – AFM images of biofilm formed by *S. aureus* ATCC 25923 to glass surface: (A) TSB alone; (B) TSB with 4 µg/mL of BC + 1/2 MIC LPE; (C) TSB with 8 µg/mL of CHX + 1/2 MIC LPE; (D) TSB with + 40% LPE (v/v); (E) TSB with 8 µg/mL TET + 1/2 MIC LPE; (F) TSB with 16 µg BC + 1/2 MIC LPE.

inhibit pathogenic bacteria and prevent oral biofilm formation, we tested in the effect of LEP on biofilm formation of nine *S. aureus* isolated from the oral cavity. In addition, we tested the potential effect of LPE to modulate susceptibility of some drugs and disinfectants of 10 *S. aureus*.

In our study, neutralized LPE supplementation (MIC values 10–50%, v/v) had potential bactericidal effect on oral *S. aureus* (Table 1). As presented in Table 1, the extract is very effective against S2 and S7 and may constitute a valuable agent to counter the growth of biofilms and undesirable

microorganisms on surfaces. Similar results have been found concerning the antibacterial activity of LPE against *Vibrio* sp.²⁹ and *Listeria monocytogenes*.³⁰ Even with sub-inhibitory concentrations (1/2 MIC) LPE showed a potential role to modulate susceptibility of the tested *S. aureus*. Since BC, TET, and CHX resistance is modulated by efflux pump systems, we suggest that LPE may potentiate, partially, the role of the disinfectant by blocking the respective efflux pump. The efflux pump is the one of major antibiotic resistance mechanisms utilized by bacterial cells to develop resistance. Thus, this resistance mechanism eliminates several classes of antibiotics such as fluoroquinolones, TET, and other compounds such as BC and CHX.

We also noted that the MIC of BC (2–4-fold reduction), CHX (2–8-fold reduction), and TET (2–8-fold reduction) was reduced when the medium was supplemented with LPE at a sub-inhibitory concentration (Table 1). The results showed a statistically significant difference between the antibacterial effects of drugs with and without LPE supplementation ($p < 0.05$). Thus, this finding confirms our hypothesis that the ability to modify susceptibility tested in this study (Table 1) is related to components that have a potential role to inhibit *S. aureus* efflux pump.

Biofilm formation is an important virulence factor of oral bacteria.⁵ Our results demonstrate that LPE exhibited a good anti-biofilm activity for all tested strains (Table 2). Similar results were found concerning the anti-biofilm activity of LPE against *L. monocytogenes*.³⁰ Previous studies have shown that the cells in a biofilm were more resistant to antimicrobial agents compared to free-floating cells.^{31,32} On the other hand, the anti-biofilm activity of disinfectants and antibiotic (BC, CHX and TET) to polystyrene and glass was enhanced when the medium was supplemented with LPE at different percentages (Table 2 and Fig. 1). The statistical analysis showed a correlation between the percentage inhibition of biofilm of different drugs alone or combined with a sub-MIC of LPE % (1/2 MIC, v/v) ($p < 0.05$). As shown in Fig. 1, AFM demonstrates that *S. aureus* biofilm were strongly affected by the supplementation of LPE. The identification of anti-biofilm constituents will be essential to be included as alternatives in the control of bacterial biofilms.

The MICs modifying ability of LPE tested in this study may be related to components that have a potential role to inhibit *S. aureus* efflux pump. Further studies are required to assess their composition and potential clinical relevance.

Conflicts of interest

The authors declare no conflicts of interest.

REFERENCES

- Huang H-L, Chang Y-Y, Lai M-C, Lin C-R, Lai C-H, Shieh T-M. Antibacterial TaN-Ag coatings on titanium dental implants. *Surf Coat Technol.* 2010;205:1636–41.
- Smith A, Jackson M, Bagg J. The ecology of *Staphylococcus* species in the oral cavity. *J Med Microbiol.* 2001;50:940–6.
- Colombo APV, Teles RP, Torres MC, et al. Subgingival microbiota of Brazilian subjects with untreated chronic periodontitis. *J Periodontol.* 2002;73:360–9.
- Souto R, Andrade AFBD, Uzeda M, Colombo APV. Prevalence of non-oral pathogenic bacteria in subgingival biofilm of subjects with chronic periodontitis. *Braz J Microbiol.* 2006;37:208–15.
- Kouidhi B, Zmantar T, Hentati H, Bakhrouf A. Cell surface hydrophobicity, biofilm formation, adhesives properties and molecular detection of adhesins genes in *Staphylococcus aureus* associated to dental caries. *Microb Pathog.* 2010;49:14–22.
- Koukos G, Sakellari D, Arsenakis M, Tsalikis L, Slini T, Konstantinidis A. Prevalence of *Staphylococcus aureus* and methicillin resistant *Staphylococcus aureus* (MRSA) in the oral cavity. *Arch Oral Biol.* 2015;60:1410–5.
- Merghni A, Nejma MB, Dallel I, et al. High potential of adhesion to biotic and abiotic surfaces by opportunistic *Staphylococcus aureus* strains isolated from orthodontic appliances. *Microb Pathog.* 2016;91:61–7.
- Proctor RA. Toward an understanding of biomaterial infections: a complex interplay between the host and bacteria. *J Lab Clin Med.* 2000;135:14–5.
- Mckenney D, Pouliot KL, Wang Y, et al. Broadly protective vaccine for *Staphylococcus aureus* based on an in vivo-expressed antigen. *Science.* 1999;284:1523–7.
- Carpentier B, Cerf O. Biofilms and their consequences, with particular reference to hygiene in the food industry. *J Appl Bacteriol.* 1993;75:499–511.
- Costerton JW, Lewandowski Z, Caldwell DE, Korber DR, Lappin-Scott HM. Microbial biofilms. *Annu Rev Microbiol.* 1995;49:711–45.
- Marshall NJ, Piddock LJ. Antibacterial efflux systems. *Microbiologia.* 1997;13:285–300.
- Littlejohn TG, Paulsen IT, Gillespie MT, et al. Substrate specificity and energetics of antiseptic and disinfectant resistance in *Staphylococcus aureus*. *FEMS Microbiol Lett.* 1992;74:259–65.
- Reverdy ME, Bes M, Brun Y, Fleurette J. Evolution of resistance to antibiotics and antiseptics of hospital *Staphylococcus aureus* strains isolated from 1980 to 1991. *Pathol Biol (Paris).* 1993;41:897–904.
- Russell AD. Do biocides select for antibiotic resistance. *J Pharm Pharmacol.* 2000;52:227–33.
- Anthonisen I-L, Sunde M, Steinum T, Sidhu M, Sørum H. Organization of the antiseptic resistance gene *qacA* and Tn552-related β -lactamase genes in multidrug-resistant *Staphylococcus haemolyticus* strains of animal and human origins. *Antimicrob Agents Chemother.* 2002;46:3606–12.
- Bjorland J, Steinum T, Kvitle B, Waage S, Sunde M, Heir E. Widespread distribution of disinfectant resistance genes among staphylococci of bovine and caprine origin in Norway. *J Clin Microbiol.* 2005;43:4363–8.
- Zmantar T, Kouidhi B, Miladi H, Bakhrouf A. Detection of macrolide and disinfectant resistance genes in clinical *Staphylococcus aureus* and coagulase-negative staphylococci. *BMC Res Notes.* 2011;4:453.
- Kwon HS, Yang EH, Yeon SW, Kang BH, Kim TY. Rapid identification of probiotic *Lactobacillus* species by multiplex PCR using species-specific primers based on the region extending from 16S rRNA through 23S rRNA. *FEMS Microbiol Lett.* 2004;239:267–75.
- Bagamboula CF, Uyttendaele M, Debevere J. Inhibitory effects of spices and herbs towards *Shigella sonnei* and *S. flexneri*. *Meded Rijksuniv Gent Fak Landbouwkd Toegep Biol Wet.* 2001;66:523–30.
- Erdemoglu N, Kupeli E, Yesilada E. Anti-inflammatory and antinociceptive activity assessment of plants used as remedy in Turkish folk medicine. *J Ethnopharmacol.* 2003;89:123–9.

22. CLSI Performance standards for antimicrobial susceptibility testing. Clinical and Laboratory Standards Institute. Wayne, PA: Clinical and Laboratory Standards Institute; 2006. Supplement M100-S16.
23. Magina M, Dalmarco E, Wisniewski A, et al. Chemical composition and antibacterial activity of essential oils of *Eugenia* species. *J Nat Med*. 2009;63:345–50.
24. Kouidhi B, Zmantar T, Jrah H, et al. Antibacterial and resistance-modifying activities of thymoquinone against oral pathogens. *Ann Clin Microbiol Antimicrob*. 2011;10: 29.
25. Merritt JH, Kadouri DE, O'toole GA. Growing and analyzing static biofilms. *Curr Protoc Microbiol*. 2005 [chapter 1]: Unit 1B 1.
26. Eijsink VG, Axelsson L, Diep DB, Havarstein LS, Holo H, Nes IF. Production of class II bacteriocins by lactic acid bacteria; an example of biological warfare and communication. *Antonie Van Leeuwenhoek*. 2002;81:639–54.
27. Kong S, Davison AJ. The role of interactions between O_2 , H_2O_2 , $\bullet OHe$ - and O_2 -in free radical damage to biological systems. *Arch Biochem Biophys*. 1980;204:18–29.
28. Garneau S, Martin NI, Vederas JC. Two-peptide bacteriocins produced by lactic acid bacteria. *Biochimie*. 2002;84:577–92.
29. Lamari F, Sadok K, Bakhrouf A, Gatesoupe F-J. Selection of lactic acid bacteria as candidate probiotics and in vivo test on *Artemia nauplii*. *Aquacult Int*. 2014;22:699–709.
30. Ben Slama R, Kouidhi B, Zmantar T, Chaieb K, Bakhrouf A. Anti-listerial and anti-biofilm activities of potential probiotic *Lactobacillus* strains isolated from Tunisian traditional fermented food. *J Food Saf*. 2013;33:8–16.
31. Frank JF, Koffi RA. Surface-adherent growth of *Listeria monocytogenes* is associated with increased resistance to surfactant sanitizers and heat. *J Food Protect*. 1990;53:550–4.
32. Krysinski E, Brown L, Marchisello T. Effect of cleaners and sanitizers on *Listeria monocytogenes* attached to product contact surfaces. *J Food Protect*. 1992;55:246–51.