



Genetic structure and gene flow in *Eugenia dysenterica* DC in the Brazilian Cerrado utilizing SSR markers

Maria Imaculada Zucchi¹, Rosana Pereira Vianello Brondani², José Baldin Pinheiro³,
Lázaro José Chaves³, Alexandre Siqueira Guedes Coelho⁴ and Roland Vencovsky^{1,3}

¹Universidade de São Paulo, ESALQ, Departamento de Genética, Piracicaba, SP, Brazil.

²Embrapa Arroz e Feijão, Santo Antônio de Goiás, GO, Brazil.

³Universidade Federal de Goiás, Escola de Agronomia, Campus Samambaia, Goiânia, GO, Brazil.

⁴Universidade Federal de Goiás, Instituto de Ciências Biológicas, Campus Samambaia, Goiânia, GO, Brazil.

Abstract

The “cagaita tree” (*Eugenia dysenterica*) is a plant found widespread in the Brazilian Cerrado. Its fruit is used for popular consumption and for industrial purposes. This study opens a new perspective for the generation of population genetic data and parameters estimates for devising sound collection and conservation procedures for *Eugenia dysenterica*. A battery of 356 primer pairs developed for *Eucalyptus* spp. was tested on the “cagaita tree”. Only 10 primer pairs were found to be transferable between the two species. Using a polyacrilamide gel, an average of 10.4 alleles per locus was detected, in a sample of 116 individuals from 10 natural “cagaita tree” populations. Seven polymorphic loci allowed estimation of genetic parameters, including expected average heterozygosity $H_e = 0,442$, among population diversity, $R_{ST} = 0,268$ and gene flow $Nm = 0,680$. Results indicated a potential of SSR locus transferability developed for *Eucalyptus* to other species of different genera, such as in the case of the “cagaita tree”. The high genetic diversity among populations detected with SSR markers indicated that these markers are highly sensitive to detect population structure. Estimated Nm values and the existence of private alleles indicated reduced gene flow and consequently possible damage to the metapopulation structure.

Key words: SSR, Myrtaceae, Cerrado, tropical tree, genetic diversity and transferability.

Received: November 21, 2002; Accepted: August 12, 2003.

Introduction

Eugenia dysenterica, commonly known as the “cagaita tree” is a fruit species native to the Cerrado (Brazilian Savannah) region belonging to the Myrtaceae family that presents potential for use in agricultural production systems (Almeida, 1998). It is outstanding for its social and economic potential for processing many sub-products and may contribute to increase income as well as jobs in the regional communities. Besides being an ornamental and honey-bearing plant, it can be used for cork extraction and in small buildings or in the manufacture of charcoal, providing good quality firewood; its bark is used in tanneries. Its leaves have anti-diarrhea and jaundice properties and its fruits are laxative (Heringer and Ferreira, 1974).

Send correspondence to: Maria Imaculada Zucchi. Universidade de São Paulo, ESALQ, Departamento de Genética, Av. Pádua Dias, 11 Caixa Postal 83, 13400-970 Piracicaba, SP, Brazil. E-mail: mizucchi@carpa.ciagri.usp.br.

The fruit trees native to the Cerrado are species from several genera and families that produce fruits of interest for food and industrialization. There is a potential and growing market for the fruit trees, which, however, is not much exploited by farmers. Fruit harvesting is mostly extractive or predatory.

The Cerrado vegetation in Brazil has been fragmented by expanding agricultural frontiers, which have affected the population dynamic of many species, including the “cagaita tree” populations. Alteration in these areas may reduce the genetic variability by founder or bottleneck effects. Genetic drift and restricted gene flow increase inbreeding and also the genetic divergence among populations. Inbreeding can lead to the fixation of deleterious alleles, threatening certain populations present in this habitat with extinction (Gilpin and Soulé, 1986; Young *et al.*, 1996).

The diversity among populations, gene flow and other genetic parameters should be evaluated in a study of natural populations of native species. Knowledge of native

species populations has been widened with the advent of molecular markers. Microsatellite markers (SSR) have been used in natural population studies (Collevatti *et al.*, 1999; Daynandan *et al.*, 1997) as they are highly polymorphic when compared with other classes of markers. SSR markers have been widely used as a tool to answer several questions on population genetics, such as gene flow and paternity analysis (Wright and Bentzen, 1994).

However, advances in the use of microsatellites have been hindered due to the high cost and time taken to develop specific primers for each locus of the native species. The chance of success in the transferability (heterologous amplification) of DNA sequences by PCR is inversely related with the evolutionary distance between the two species. Many studies have shown the possibility of using pairs of primers designed for one species belonging to the same genus (Cipriani *et al.*, 1999; Isagi and Suhandono, 1997) or even among different genera (Roa *et al.*, 2000; White and Powell, 1997).

In the present study the transferability to *Eugenia dysenterica* of primer pairs developed for *Eucalyptus* spp., which belongs to the same family, was investigated in order to identify microsatellite markers in the “cagaia tree” for studies of the genetic variability, population structure, gene flow and reproductive system. The main objective of this research was the generation of information for domestication and breeding of the species and its conservation.

Material and Methods

Plant material

The study material was collected in ten locations in southeast Goiás state, forming ten populations, represented by 116 trees (matrices). Plant material (leaves) was collected from each one of the 116 “cagaia trees” for genotypic characterization of the plants. Table 1 shows the locations where the populations were collected, including some characterization of the areas.

SSR locus transferability and amplification by PCR

In this study the transferability of primer pairs developed for *Eucalyptus* was assessed for *Eugenia dysenterica*, which belongs to the same Mirtaceae family but to different genera.

Three hundred and fifty-six primer pairs developed for *Eucalyptus* spp. (*Eucalyptus grandis* x *Eucalyptus urophylla*) by Brondani *et al.* (1998) were used. These pairs of primers were tested for amplification in *Eugenia dysenterica*.

For the reaction of amplification by PCR, 15 ng of genomic DNA were used in 25 μ L volume containing 50 mM KCl, 20 mM Tris-HCl pH 8.8; 1.5 mM MgCl₂; 10 mM dNTPs; 0.2 μ M of each primer (forward and reverse) and 1U of Taq polymerase.

The PCR protocol consisted of a prior initial denaturation at 96 °C for 3 min, followed by 30 cycles at 94 °C for 30 s, 60 °C for 1 min, 72 °C for 1 min, and the last step for extension at 72 °C for 7 min. The amplified fragments were separated in 4% polyacrylamide gel, in a run with 1X TBE at 2000 v for 2 h, and stained with silver nitrate.

Statistical analysis of the data

Allelic and genotypic frequencies for each locus were obtained from the data readings in the gels. These frequencies were submitted to a goodness-of-fit test (Fisher’s exact test) to the proportion of Hardy-Weinberg equilibrium as defined by Weir (1996) using the TFGA program (Miller, 1997). Fisher’s exact test was performed by the conventional Monte Carlo method using 10 batches with 1,000 permutations per batch.

Genetic diversity and F statistics were estimated under a random model according to Weir (1996) where the sampled populations are considered as representatives of the species with a common evolutionary history. The allelic frequencies, the number of alleles per locus (A), the observed (H_o) and expected (H_e) heterozygosities and the F statistics of Wright (F_{IS} , F_{ST} and F_{IT}) were estimated using the GDA program (Lewis and Zaykin, 2000).

Table 1 - Locations in the state of Goiás, number of “cataiga trees” sampled and respective geographic position.

Areas	Site municipal district	Situation of areas	Number of trees sampled	Altitude (m)	Latitude	Longitude
1	Catalão	Intact Cerrado	12	880	18°07'35''	47°54'20''
2	Catalão	Pasture	12	860	18°02'0,3''	48°02'31''
3	Catalão	Pasture	12	800	18°13'39''	47°58'12''
4	Três Ranchos	Pasture	12	820	18°17'15''	47°48'41''
5	Campo Alegre de Goiás	Pasture	12	930	17°39'11''	47°46'37''
6	Campo Alegre de Goiás	Pasture	12	780	17°34'24''	47°42'12''
7	Cristalina	Intact Cerrado	5	890	17°10'47''	47°31'07''
8	Luziânia	Intact Cerrado	11	900	16°28'48''	47°48'40''
9	Goiânia	Urban	16	740	16°40'30''	49°14'42''
10	Senador Canedo	pasture	12	840	16°37'13''	49°04'29''

The mutation process in microsatellite loci is not in line with the expectations under an infinite alleles model with low mutation rates. Therefore the analogue of the F_{ST} statistics, namely the R_{ST} parameter (Slatkin, 1995) developed specifically for microsatellite data, was also used. Parameters R_{ST} and gene flow (Nm) were estimated using the $R_{ST}Cal$ program (Goodman, 1997). The variability structure was visualized using dendrograms constructed from the matrix of Nei's genetic distances and the UPGMA clustering criteria, using the NTSYS program (Rolf, 1989). The stability of the clusters was also tested by a re-sampling procedure with 10,000 bootstraps.

The patterns of spatial variation were analyzed using Pearson's coefficient of correlation (r) between Nei's genetic distance matrix (Nei, 1972) and the geographic distances between populations matrix. The significance of this correlation was tested through Mantel's Z statistic (Mantel, 1967), using 9,999 random permutations.

Results

Transferability of SSR *Eucalyptus* primers to *Eugenia dysenterica*

The 356 primers tested were classified according to the quality obtained in the PCR: 2.8% (ten pairs of primers) amplified clear SSR products, 30.0% presented non specific band amplification and 67.2% did not amplify any band (Zucchi *et al.* 2002).

The selected ten pairs of primers were used for the population genetic structure study. First, all primers were submitted to the basic program with 56 °C for primer annealing. Those that did not amplify satisfactorily were submitted to amplification cycles with lower annealing temperatures. EMBRA 17 and EMBRA 134 amplified satisfactorily at 52 °C.

Table 2 shows the amplification conditions used for the ten SSR loci assessed and their respective allelic amplitudes. The greatest allelic amplitude was 167 base pairs. The allelic frequencies are shown in Table 3.

Genetic Variation

The average number of alleles per polymorphic locus was 10.43, with a range from three alleles (EMBRA 73 locus) to 22 alleles (EMBRA 14 locus).

Table 4 shows that the observed heterozygosity ranged from 0.253 (population 1) to 0.599 (Population 10) with a mean of 0.458. The lowest expected heterozygosity was 0.276 for population 1, whereas population 9 presented the greatest expected heterozygosity (0.670). The mean value obtained was 0.442.

Fisher's exact test (Table 5) showed that some populations were not in Hardy-Weinberg Equilibrium for most of the studied loci. For population 10 this test was significant for six out of seven loci for which the Hardy-Weinberg equilibrium condition was therefore rejected.

Table 2 - Sequence of primer pairs developed for *Eucalyptus** that amplified microsatellite loci in *Eugenia dysenterica* with allele size range, number of alleles per locus (A), expected heterozygosity (H_e), observed heterozygosity (H_o) and annealing temperature (T_a).

SSR locus	Sequence of primers	Allele size range (bp)	A	H_e	H_o	T_a (°C)	GenBank accession number
EMBRA 14	F-5'gCC TCA AAC CAA TTC AAA T3' R-5'CAT gAT TCT CCC ACT CCT C3'	95-170	22,0	0,837	0,514	56	G74881
EMBRA 210	F-5'CgT gTg gTT Atg TgA ACT3' R-5'CCT AAC AAT gCA TAA gCT C3'	88-245	14,0	0,831	0,706	56	G74883
EMBRA 63	F-5'-CAT CTg gAg ATC gAg gAA-3' R-5'-gAg AgA Agg ATC ATg CCA-3'	165-175	4,0	0,497	0,681	56	G74884
EMBRA 122	F-5'TTg CTC CAT CTT TCT TgC3' R-5'AAA ACg ATT AgA ggg TCA Tg3'	290-340	12,0	0,675	0,406	56	G74876
EMBRA 73	F-5'Cgg TCg TTg TCg gAA TCT C3' R-5'AgT Tgg gTA ACg CCA ggT TT3'	153-158	3,0	0,198	0,207	52	G74880
EMBRA 172	F-5'AAA gCg AACggTCACACC3' R-5'gTgCTTCTCCAaggTTCTgATC3'	110-132	5,0	0,669	0,496	56	G74877
EMBRA 72	F-5'CTggTCAACgTCCgAAA3' R-5'AtgCTgCAGAgggCATAA3'	100-190	13,0	0,405	0,200	56	G74879
EMBRA 134	F-5'CTC TgA ggA gTT ggC AgT AgC3' R-5'CAC gTT TAA ATg CgC AA gTg3'	Monomorphic	1,0	-	-	52	G74885
EMBRA 179	F-5'gTCggCTCACAgCATGAA3' R-5'gCCTCCA gTAGTTAACAGACG3'	Monomorphic	1,0	-	-	56	G74878
EMBRA 243	F-5'gCgTACggATCAA gAACA3' R-5'gAAA g gAACgCCA ACTAA3'	Monomorphic	1,0	-	-	56	G74882

*Brondani *et al.* (1998).

Table 3 - Table of allelic frequencies of the seven SSR loci, estimated from 116 individuals of 10 populations of *Eugenia dysenterica*.

Locus	Allele	Pop1	Pop2	Pop3	Pop4	Pop5	Pop6	Pop7	Pop8	Pop9	Pop10
Embra 14	95	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.045
	100	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.063	0.000
	102	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.031	0.000
	110	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.063	0.000
	111	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.045
	113	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.094	0.000
	115	0.042	0.000	0.100	0.000	0.000	0.000	0.800	0.000	0.000	0.000
	118	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.500	0.000	0.000
	120	0.583	0.182	0.150	0.125	0.227	0.227	0.000	0.083	0.031	0.000
	125	0.250	0.500	0.600	0.583	0.500	0.682	0.000	0.000	0.000	0.000
	130	0.042	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
	132	0.042	0.227	0.050	0.292	0.227	0.091	0.100	0.083	0.063	0.000
	135	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.083	0.125	0.000
	137	0.000	0.091	0.100	0.000	0.045	0.000	0.000	0.000	0.000	0.000
	140	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.375	0.273
	142	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.318
	145	0.042	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
148	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.182	
150	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.083	0.094	0.000	
155	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.045	
162	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.083	0.000	0.000	
170	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.100	0.083	0.063	
Embra 210	170	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.182	0.000	0.000
	182	0.042	0.000	0.333	0.000	0.000	0.000	0.000	0.000	0.000	0.000
	188	0.000	0.083	0.042	0.250	0.583	0.200	0.250	0.318	0.000	0.000
	190	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.067	0.000
	194	0.083	0.000	0.125	0.000	0.083	0.000	0.375	0.000	0.000	0.000
	195	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.433	0.083
	204	0.000	0.000	0.000	0.000	0.000	0.150	0.000	0.000	0.000	0.000
	205	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.067	0.333
	210	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.125
	215	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.233	0.292
	220	0.250	0.458	0.167	0.375	0.042	0.150	0.125	0.455	0.167	0.167
225	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.033	0.000	
230	0.625	0.458	0.333	0.333	0.208	0.450	0.250	0.045	0.000	0.000	
243	0.000	0.000	0.000	0.042	0.083	0.050	0.000	0.000	0.000	0.000	
Embra 63	165	0.167	0.333	0.333	0.125	0.500	0.500	0.500	0.364	0.000	0.000
	168	0.000	0.083	0.000	0.333	0.000	0.000	0.000	0.136	0.000	0.000
	170	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.094	0.125
	175	0.833	0.583	0.667	0.542	0.500	0.500	0.500	0.500	0.906	0.875
Embra 122	290	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.056
	296	0.773	0.500	0.438	0.750	0.800	0.875	0.500	0.727	0.000	0.000
	298	0.045	0.000	0.188	0.125	0.000	0.000	0.000	0.000	0.000	0.000
	300	0.182	0.500	0.375	0.125	0.200	0.125	0.500	0.273	0.000	0.000
	305	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.033	0.000
	310	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.233	0.222
	315	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.056
	320	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.200	0.278
	327	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.133	0.333
	330	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.067	0.056
340	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.300	0.000	
343	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.033	0.000	

Table 3 (Cont.)

Locus	Allele	Pop1	Pop2	Pop3	Pop4	Pop5	Pop6	Pop7	Pop8	Pop9	Pop10
Embra 73	153	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.125	0.125
	155	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.281	0.375
	158	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	0.594	0.500
Embra 172	110	0.042	0.292	0.000	0.125	0.208	0.333	0.000	0.045	0.300	0.042
	120	0.000	0.000	0.125	0.208	0.458	0.167	0.300	0.409	0.300	0.292
	123	0.958	0.708	0.875	0.667	0.333	0.500	0.700	0.000	0.133	0.250
	130	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.545	0.233	0.417
	132	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.033	0.000
Embra 72	100	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	0.000	0.000
	107	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.063	0.000
	119	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.031	0.000
	120	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.031	0.000
	130	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.031	0.000
	143	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.063	0.000
	160	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.344	0.227
	169	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.344	0.273
	180	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.031	0.045
	182	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.063	0.182
	190	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.091
	195	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.136
210	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.045	

Table 4 - Estimates of genetic parameters in ten *Eugenia dysenterica* populations N: number of individuals sampled, L: number of microsatellite loci; A: mean number of alleles; H_o: Observed heterozygosity; H_e expected heterozygosity; f: fixation index; t_a: apparent crossing rate.

Pop.	N	L	A	H _o	H _e	f	t _a
1	12	7	2,714	0,253	0,276	0,088	0,838
2	12	7	2,286	0,449	0,403	-0,123	1,280
3	12	7	2,714	0,366	0,394	0,076	0,859
4	12	7	2,572	0,453	0,408	-0,117	1,265
5	12	7	2,572	0,497	0,404	-0,245	1,649
6	12	7	2,428	0,411	0,374	-0,106	1,237
7	5	7	2,143	0,450	0,389	-0,183	1,448
8	11	7	3,000	0,563	0,438	-0,316	1,924
9	16	7	6,000	0,541	0,670	0,197	0,671
10	12	7	4,857	0,599	0,667	0,105	0,810
Average	11	7	3,128	0,458	0,442	-0,037	1,077

Figure 1 shows many exclusive alleles in certain populations, with the presence of 15 exclusive alleles in population 9, 11 in population 10, three in population 8, two in population 2 and one in population 6. Although the populations shared most of the 73 alleles there were an expressive number of alleles that characterized certain populations specifically (such as in the case of populations 9 and 10).

Genetic structure

Table 4 shows the intrapopulation fixation index ($f = F_{IS}$) estimated for each population from the heterozygosity indexes (H_o and H_e). The mean value was $f = -0.037$ with a range of -0.316 to 0.197 . From the analysis of variance the average $f = -0.017$ was obtained (IC 95%: -0.276 to 0.134 ; Table 6). These results, together with Fisher's exact test, indicated that the populations have a rate of cross-pollination compatible with allogamy.

The estimated R_{ST} and F_{ST} values were 0.269 and 0.250, respectively. These estimates are similar and both significantly different from zero, as can be observed in Table 6. The Nm parameter estimate, calculated from the R_{ST} estimate, was 0.680 individuals per generation, indicating an intermediate migration rate among the populations. Considering the $F_{ST} = 0.250$ estimate, $Nm = 0.750$ individuals were obtained per generation.

Nei's genetic distances calculated pairwise between the populations varied from 0.046 to 0.407. These led to the dendrogram presented in Figure 3. The cophenetic correlation of the UPGMA clustering of this matrix was high (0.943). Populations 9 and 10 were shown to be very similar and formed a genetically divergent group. They are located in the extreme west of the sampled region. Populations 1 to 8 formed another group and are located in the eastern part of the studied region. It is important to point out that these populations are separated by the Corumbá river depression, forming two spatially discontinuous groups. This structure was visualized by the dendrogram,

Table 5 - Probabilities of goodness-of-fit test (Fisher's exact test) to Hardy-Weinberg equilibrium for each locus.

Locus	Pop1	Pop2	Pop3	Pop 4	Pop 5	Pop 6	Pop 7	Pop 8	Pop 9	Pop 10
EMBRA 14	0,012	0.084	0.182	0.779	0.657	0.034	0.113	0.408	0.001	0.022
EMBRA 210	0.578	0.001	0.000	0.164	0.436	0.080	0.647	0.637	0.739	0.001
EMBRA 63	1.000	0.136	0.214	0.027	0.0025	0.001	0.121	0.008	1.000	1.000
EMBRA 122	0.427	1.000	0.406	0.139	1.000	1.000	0.128	0.501	0.000	0.011
EMBRA 73	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.414	0.004
EMBRA 172	1.000	1.000	1.000	0.395	0.002	0.842	0.331	0.097	0.000	0.007
EMBRA 72	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.022

Table 6 - Wright's F statistic's, R_{ST} and the number of migrants per generation (N_m) for ten natural *Eugenia dysenterica* populations. Confidence interval of 95% probability based on 10,000 bootstraps.

	F_{IS}	F_{IT}	F_{ST}	R_{ST}	N_{m_A}	N_{m_B}
Under all the loci	-0,017	0,238	0,250	0,269	0,680	0,750
Upper (IC 95%)	0,134	0,388	0,348	0,359	0,766	—
Lower (IC 95%)	-0,276	0,025	0,194	0,245	0,442	—

N_{m_A} : based in R_{ST} .
 N_{m_B} : based in F_{ST} .




Figure 1 - Number of private/exclusive alleles for 73 alleles obtained in seven polymorphic loci of the SSR markers, in ten *Eugenia dysenterica* populations.




Figure 2 - Genetic divergence pattern among ten "cagaita tree" populations, defined by the UPGMA clustering, based on the genetic identity obtained from Nei's genetic distances (1972). Cophenetic correlations equal to 0.943.

where populations 9 and 10 diverged genetically from the other populations. This population structure is congruent with the results obtained by Telles *et al.* (2001) using isoenzymes, with progeny data of the same populations.

The matrix correlation between Nei's genetic distances and the respective geographic distances was high




Figure 3 - Correlation (r) between the geographic distance matrix and the genetic distance matrix obtained by SSR markers.

and positive ($r = 0.872$) and significant at 1% level of probability (Figure 3). This result indicates that the genetic variability pattern among the populations is structured in space. A similar result but with a lower correlation ($r = 0,725$) was reported by Telles *et al.*, 2001, with isoenzyme markers.

Discussion

Knowledge of the genetic variability distribution between and within natural *Eugenia dysenterica* populations is essential to adopt efficient strategies for *ex situ* and *in situ* germplasm conservation.

SSR markers are a powerful tool for this type of study. However, the progress of using markers based on microsatellites has been hindered because of the high cost and time spent for developing species specific primers. The chance of success of heterologous amplification for any DNA sequence by PCR is inversely related to the evolutionary distance between two species. Many studies have shown, however, that there is the possibility of using pairs of primers designed for one species or another species of the same genus (Cipriani *et al.*, 1999) or even of different genera (Roa *et al.*, 2000). The transferability of microsatellites among related species is a consequence of the homology of the DNA sequence in the regions that flank the microsatellites. Other studies on tropical trees have demonstrated a high ratio of SSR loci transferability among taxonomically related tree species, as occurs with

Leguminosae (Dayanandan *et al.*, 1997), Meliaceae (White and Powell, 1997) and among *Eucalyptus* species (Brondani *et al.*, 1998). In this study, only 10 primer pairs of 356 developed for *Eucalyptus* spp. were found to be transferable the two species (Zucchi, *et al.* 2002).

Roa *et al.* (2000) studied transferability in cassava (*Manihot esculenta*) for six different species (all wild) of the *Manihot* genus. Only two in eight of the amplified loci (or two pairs of primers) did not amplify for the two more distant wild cassava species. It has been found that many microsatellite primers can be used to amplify heterology among different genera. These authors showed the possibility of using SSR primers to amplify heterology in different species and genera (Byrne *et al.*, 1996; Katzir *et al.*, 1996; Isagi and Suhandono, 1997; Smulder *et al.*, 1997; Steinkellner *et al.*, 1997).

Dayanandan *et al.* (1997) used pairs of primers developed for a tropical tree, *Pithecellobium elegans*, to detect SSR loci that amplified for other species in the same family (Leguminosae). Thirteen species from the Leguminosae family, 12 from the Mimosidae subfamily and one from the Pappilionoideae family, were used. The six pairs of primers developed for *P. elegans* were successful in amplifying for species of the same genus and of different genera.

Regarding genetic variation, a relatively high level of multiallelism in all the seven polymorphic loci was observed in the present study. The average number of alleles per locus was 10.4 and the mean expected heterozygosity reached 0.442 which is greater than the value found in a similar study with the same “cagaita tree” populations using isoenzyme markers.

The F_{IS} value found here for the *Eugenia dysenterica* populations was negligible. This value suggests that the species is predominantly allogamic. This result contrasted $F_{IS} = -0.017$ value obtained by SSR markers with the $F_{IS} = 0.243$ value obtained by Telles (2000) with isozymes from seedlings of the same population. The conflicting results may be due to the use of data at different stages of development, seedling or adults plants could show different isozymes patterns or even because of differences in the nature of the genetic marker used. Enzymes may be related to adaptive traits and subject to natural selection, while microsatellites are non coding regions of the genome and, therefore, are selectively neutral.

Proença & Gibbs (1994) studied the reproductive biology of *E. dysenterica* and concluded that it is pollinated by large bees. The flowers open in the morning for one day and following the pattern called “big bang”, the plants flower intensely for a relatively short period. Mainly monkeys and humans disperse the seeds, although some other animals also perform this function (Ferreira & Cunha, 1980).

The apparent cross-fertilization rate here estimated was high ($\hat{t} = 1,08$) and greater than that found with isozyme markers ($\hat{t} = 0.83$) and seedling data. Pollinators or

dispersing agents may also be responsible for this difference.

For example, the trees analyzed with SSR markers are probably plants that had been part of the natural Cerrado and were, in most cases, 50 old years or more. On the other hand seedlings (progenies genotyped with isoenzymes) are recent plants from recent pollination events performed by insect populations that probably are different from the insect population of 50 years ago.

Govindajaru (1989) distinguished three levels of gene flow: high $N_m > 1$, intermediate ($0.25 < N_m < 0.99$) and low $N_m < 0.25$. The value found here ($N_m = 0,68$) was therefore intermediate. As this flow was estimated on the basis of the R_{ST} parameter it cannot not be considered contemporaneous, but a consequence of the genetic history of these populations. The restricted gene flow can be also explained by main pollinator, the large bees, and the flowering manner of the “cagaita tree”, that happens fast and abundantly, that does not allow a great number of flights of the pollinator to supply itself with pollen, and is thus restricted to small distances.

Regarding the measure of diversity among populations, it was noted that the R_{ST} and F_{ST} estimates were very similar. It is believed that more results of this nature would be necessary to ascertain the tendencies in the difference between R_{ST} and F_{ST} . These values were higher than those obtained with isozyme markers for the same populations suggesting that microsatellites are more sensitive than isozymes for measuring differentiation among populations than isozymes. This was also evident from the detection of private or exclusive alleles in this study that were not detected in the data reported by Telles (2001).

Although this is a species with a high degree of allogamy, the estimated gene flow among the populations was relatively small and possibly a consequence of human settlement of the Cerrado. An interesting fact to discuss is the high frequency of exclusive alleles in populations 9 and 10 that may have been caused by genetic drift and absence of gene flow. In fact, population 9 is located in an urban region (Table 1) and population 10, although natural, is completely isolated from the others.

Of the localities studied except areas 1, 8 and part of area 7, the others were present in areas with bigger alteration in consequence of human settlement of the Cerrado, either for locating inside the urban area of Goiânia (area 9), or for locating in implanted pastures. Some areas have been fertilized when these grasses were planted, as for example area 2, 4 and 10, that presented higher calcium (Ca), magnesium (Mg) and phosphorus (K) values in comparison with natural areas (Silva, 1999).

This species is widely used by human populations for its wood and fruit, it possesses a reproductive system (preferentially allogamous), and it is pollinated mainly by large bees (*Bombus* sp.). Its demographic and biological characteristics, associated with habitat fragmentation, tend to pro-

duce a relatively high amount of genetic divergence among the local population (Loveless and Hamrick 1984; Proença and Gibbs, 1994). Due to recent expansion of agricultural activities in the Cerrado and high rates of biodiversity loss and endemic species, the region is considered one of the world's hotspots for conservation (Myers *et al.* 2002). The region is highly fragmented by increased agricultural activities, and there is no clear information about the spatial distribution of *Eugenia dysenterica* trees before recent human occupation (Diniz-Filho and Telles, 2002).

Another hypothesis to be considered regarding the high rate of exclusive alleles in population 9 is that its genetic constitution could have been altered by human introductions. Slatkin (1985) described a methodology to assess gene flow from rare (or private/exclusive) alleles. The distribution of the frequencies of these rare alleles (alleles that appear in a single population) is used to estimate the mean number of migrants exchanged among local populations. The logarithm of Nm is approximately linearly related to the logarithm of the mean of the frequency of the private/exclusive alleles (Slatkin, 1985; Slatkin and Barton, 1986).

The high correlation coefficient between the genetic and geographic distance matrices suggested that there is a spatial pattern of genetic variability among the populations. This structure probably originated from a stochastic differentiation process, with higher levels of gene flow among closer populations and decreased flow as distances increased (isolation by distance).

Results indicated a potential of SSR locus transferability developed for *Eucalyptus* to other species of different genera, such as in the case of the "cagaita tree". The high genetic diversity among populations detected with SSR markers indicated that these markers are highly sensitive to detect population structure. Results of the present study were not entirely congruent with those obtained with isozyme markers especially with respect to gene flow and diversity. However estimated Nm being less than 1.0 and the existence of private alleles call attention to damages of the metapopulation structure that may have occurred in these populations.

References

- Almeida SP (1998) Frutas nativas do cerrado: caracterização físico-química e fonte potencial de nutrientes. In: Embrapa-CPAC (ed) Cerrado: ambiente e flora. Plantaltina, pp 556.
- Brondani RPV, Brondani C, Tarchini R and Grattapaglia D (1998) Development, characterization and mapping of microsatellite markers in *Eucalyptus grandis* and *E. urophylla*. *Theor Appl Genet* 97:816-827.
- Byrne M, Marquez-Garcia MI, Uren T, Smith DS and Moran GF (1996) Conservation and genetic diversity of microsatellite loci, in the genus *Eucalyptus*. *Aust J Bot* 44:331-341.
- Cipriani G, Lot G, Huang WG, Marrazzo MT, Peterlunger E and Testolin R (1999) AC/GT and AG/CT microsatellite repeats in peach [*Prunus persica* (L) Batsch]: isolation, characterization and cross-species amplification in *Prunus*. *Theor Appl Genet* 99:65-72.
- Collevatti RG, Brondani RPV and Grattapaglia D (1999) Development and characterization of microsatellite markers for genetic analysis of a Brazilian endangered tree species of *Caryocar brasiliense*. *Heredity* 83:748-756.
- Daynandan S, Bawa KS and Kesseli R (1997) Conservation of microsatellites among tropical tree (Leguminosae). *Am J Bot* 84:1658-1663.
- Diniz-Filho JAF and Telles MCP (2002) Spatial Autocorrelation analysis and the identification of operational units for conservation in continuous populations. *Conservation Biology* 16:924-935.
- Ferreira MB and Cunha LHS (1980) Dispersão de plantas lenhosas de cerrado: germinação e desenvolvimento. *Informe Agropecuário* 16:12-17.
- Gilpin ME and Soulé ME (1986) Minimum viable populations: process of species extinction. In: Soulé, ME (ed) *Conservation Biology, the Science of Scarcity and Diversity*, Sinauer Associates, Sunderland, pp 19-34.
- Goodman SJ (1997) R_{ST}Calc: a collection of computer program for calculating estimates of genetic differentiation from microsatellite and determining their significance. *Mol Ecol* 6:881-885.
- Govindajuru RD (1989) Variation in gene flow levels among predominantly self-pollinated plants. *J Evol Biol* 2:173-181.
- Heringer EP and Ferreira MB (1974) Informações preliminares acerca da floração precoce de vinte espécies arbóreas do Cerrado do Planalto Central. 25 Congresso Nacional de Botânica, Mossoró, Brasil.
- Isagi Y and Suhandono S (1997) PCR primers amplifying microsatellite loci of *Quercus myrsinifolia* Blume and their conservation between oak species. *Mol Ecol* 6:897-899.
- Katzir N, Danin-Poleg Y, Tzuki G, Karchi Z, Lavi V and Cregan PB (1996). Length polymorphism and homology of microsatellites in several Curcubitaceae species. *Theor Appl Genet* 93:1282-1290.
- Lewis PO and Zaykin D (2000) Genetic Data Analysis: Computer program for the analysis of allelic data. Version 1.0 (d15) Free program distributed by authors over the Internet from the GDA Home Page at <http://alleyn.eeb.uconn.edu/gda/2000>.
- Mantel N (1967) The detection of disease clustering and a generalized regression approach. *Cancer Resear* 27:209-220.
- Miller M (1997) Tools For Population Genetic Analyses (TFPGA) 1.3: A windows program for analyses of allozyme and molecular population genetic data.
- Myers N, Mittermeier RA, Mittermeier CG, Fonseca GAB and Kent J (2000) Biodiversity hotspot for conservation priorities. *Nature* 403:853-858.
- Nei M (1972) Genetics distance between populations. *Am Naturalist* 106:283-292.
- Proença CEB and Gibbs PE (1994) Reproductive biology of eight sympatric Myrtaceae from Central Brazil. *New Phytologist* 126:342-354.
- Roa AC, Chavarriaga-Aguirre P, Duque MC, Maya MM, Bonierbale MW, Iglesias C and Thome J (2000) Cross-species amplification of cassava (*Manihot esculenta*) (Euphorbiaceae) microsatellites: allelic polymorphism and degree of relationship *Am J Bot* 87:1647-1655.

- Rolf FJ (1989) NTSYS-Pc: Numerical Taxonomy and Multivariate Analysis System. Exeter publisher, New York.
- Silva RSM (1999) Caracterização de sub-populações de cagaiteira (*Eugenia dysenterica* DC.) da região sudeste do estado de Goiás. Dissertação, Escola de Agronomia, Universidade Federal de Goiás, Goiânia.
- Slatkin M (1985) Gene flow in natural population. *Ann Rev Ecol Syst* 16:393-430.
- Slatkin M (1995) A measure of population subdivision based on microsatellite allele frequencies. *Genetics* 139:457-462.
- Slatkin M and Barton NH (1986) A comparison of three indirect methods for estimating average levels of gene flow. *Evolution* 43:1349-1368.
- Smulder MJM, Bredemeijer G, Rus-Kortekass W, Arens P and Vosman B (1997) Used of short microsatellite from base sequence to generate polymorphism among *Lycopersicum esculentum* cultivars and accession of other *Lycopersicum* species. *Theor Appl Genet* 97:264-272.
- Steinkellner H, Lexer C, Tutresschek E, Glossil J (1997) Conservation of (GA)_n microsatellite loci between *Quercus* species. *Mol Ecol* 6:1189-1194.
- Telles M. Diversidade genética e estrutura genética populacional de cagaiteira (*Eugenia dysenterica* DC.) do sudeste de Goiás. Dissertação, Escola de Agronomia, Universidade Federal de Goiás, Goiânia.
- Telles MPC, Diniz-Filho JAF, Coelho ASG and Chaves LJ (2001) Autocorrelação espacial das frequências alélicas em subpopulações de cagaiteira (*Eugenia dysenterica* DC, Myrtaceae) no Sudeste de Goiás. *Rev Bras Bot* 24:145-154.
- Weir BS (1996) *Genetics data analysis II – Methods for discrete population genetic data*. Sunderland MA, Sinauer Associates, Inc. Publishers.
- White G and Powell W (1997) Cross-species amplification of SSR loci in the Meliaceae family. *Mol Ecol* 6:1195-1197.
- Wright JM and Bentzen P (1994) Microsatellites: genetic markers of the future. *Rev Fish Biol Fisher* 4:384-388.
- Young A, Boyle T and Brown T (1996) The population genetic consequences of habitat fragmentation for plants. *Trends Ecol Evol* 11:413-418.
- Zucchi MI, Brondani RV, Pinheiro JB, Brondani C and Vencovsky R (2002) Transferability of microsatellite markers from *Eucalyptus* spp. to *Eugenia dysenterica* (Myrtaceae family). *Mol Ecol Notes* 2:512-514.

Editor: Natal Antonio Vello