



Chromosome 17 abnormalities and mutation of the TP53 gene: Correlation between cytogenetics, flow cytometry and molecular analysis in three cases of chronic myeloid leukemia

Luize Otero¹, Geraldo Barroso Cavalcanti Júnior^{2,3}, Claudete Esteves Klumb², Marcos Antonio Mauricio Scheiner², Eliane Pereira Simões Magluta², Teresa de Souza Fernandez^{1,4}, Maria Luiza Macedo Silva¹, Virgínia Pires¹, Gabriela Vasconcelos Andrade¹, Raquel Ciuvalschi Maia² and Daniel Tabak¹

¹*Instituto Nacional de Câncer, Centro de Transplante de Medula Óssea, Rio de Janeiro, RJ, Brazil.*

²*Instituto Nacional de Câncer, Serviço de Hematologia, Laboratório de Hematologia Celular e Molecular, Rio de Janeiro, RJ, Brazil.*

³*Universidade Federal do Rio Grande do Norte, Departamento de Análises Clínicas e Toxicológicas, Natal, RN, Brazil.*

⁴*Universidade do Estado do Rio de Janeiro, Departamento de Patologia Geral, Rio de Janeiro, RJ, Brazil.*

Due to a printing error, the image shown in Figure 1 of the article “Chromosome 17 abnormalities and mutation of the TP53 gene: Correlation between cytogenetics, flow cytometry and molecular analysis in three cases of chronic myeloid leukemia” published in v. 28, n. 1, p. 40-43 (2005) was inverted. Please find below the correctly printed Figure.




Figure 1 - Partial karyotypes of cases 1 (A), 2 (B) and 3 (C).