



## Genome-wide identification and phylogenetic analysis of the ERF gene family in cucumbers

Lifang Hu<sup>2</sup> and Shiqiang Liu<sup>1</sup>

<sup>1</sup>*School of Sciences, Jiangxi Agricultural University, Nanchang, China.*

<sup>2</sup>*School of Agriculture, Jiangxi Agricultural University, Nanchang, China.*

### Abstract

Members of the ERF transcription-factor family participate in a number of biological processes, viz., responses to hormones, adaptation to biotic and abiotic stress, metabolism regulation, beneficial symbiotic interactions, cell differentiation and developmental processes. So far, no tissue-expression profile of any cucumber ERF protein has been reported in detail. Recent completion of the cucumber full-genome sequence has come to facilitate, not only genome-wide analysis of ERF family members in cucumbers themselves, but also a comparative analysis with those in *Arabidopsis* and rice. In this study, 103 hypothetical ERF family genes in the cucumber genome were identified, phylogenetic analysis indicating their classification into 10 groups, designated I to X. Motif analysis further indicated that most of the conserved motifs outside the AP2/ERF domain, are selectively distributed among the specific clades in the phylogenetic tree. From chromosomal localization and genome distribution analysis, it appears that tandem-duplication may have contributed to *CsERF* gene expansion. Intron/exon structure analysis indicated that a few *CsERFs* still conserved the former intron-position patterns existent in the common ancestor of monocots and eudicots. Expression analysis revealed the widespread distribution of the cucumber *ERF* gene family within plant tissues, thereby implying the probability of their performing various roles therein. Furthermore, members of some groups presented mutually similar expression patterns that might be related to their phylogenetic groups.

*Key words:* *Cucumis sativus* L., ERF, phylogenetic analysis, transcription factor, genome sequence.

Received: January 21, 2011; Accepted: August 18, 2011.

### Introduction

The AP2/ERF superfamily, one of the largest groups of transcription factors in plants, is characterized by the presence of the AP2/ERF-type DNA-binding domain consisting of from 60 to 70 highly conserved amino acids (Wessler, 2005). Based on sequence similarities and the number of AP2/ERF domains, this superfamily can be classified into three families, viz., AP2, ERF and RAV (Sakuma *et al.*, 2002; Nakano *et al.*, 2006). AP2 family proteins contain two repeated AP2/ERF domains, the ERF, a single AP2/ERF domain, and the RAV one AP2/ERF domain, as well as a B3 domain conserved in other plant-specific transcription factors. The ERF family is usually classified into two major subfamilies, CBF/DREB, and ERF, the latter based on the amino acid sequence of the DNA-binding domain. Both are divisible into I to X groups (Nakano *et al.*, 2006).

ERF family proteins are involved in a series of biological events, such as hormonal signal transduction medi-

ated by ethylene, cytokinin and brassinosteroid (Hu *et al.*, 2004; Rashotte *et al.*, 2006), response to biotic and abiotic stress (Stockinger *et al.*, 1997; Liu *et al.*, 1998), metabolism regulation (van der Fits and Memelink, 2000; Aharoni *et al.*, 2004; Zhang *et al.*, 2005), beneficial symbiotic interaction (Vernie *et al.*, 2008), and cell differentiation (Iwase *et al.*, 2011), as well as developmental processes, such as leaf epidermal cell density (Moose and Sisco, 1996), flower development (Elliott *et al.*, 1996), and embryo development (Boutillier *et al.*, 2002) in various plant species. To date, some ERF family proteins have been identified in various plant species, viz., *Arabidopsis* (*Arabidopsis thaliana*) (Sakuma *et al.*, 2002), soybeans (Li *et al.*, 2005), rice (Cao *et al.*, 2006; Sharoni *et al.*, 2011), cotton (Jin and Liu, 2008), *Populus trichocarpa* (Zhuang *et al.*, 2008), tomato (Sharma *et al.*, 2010) and *Vitis vinifera* (Licausi *et al.*, 2010). The sequenced *Arabidopsis* genome contains 147 postulated genes encoding AP2/ERF-type proteins, 122 of which belonging to the ERF family (Nakano *et al.*, 2006). In *Arabidopsis*, expression of both the *DREB1A* gene and its two homologs in group III is induced by low-temperature stress, but not by drought or high-salt stress, whereas, expression of both the *DREB2A* gene and its single homolog in another group, group IV, is induced by dehydration, but

not by low-temperature stress (Liu *et al.*, 1998; Gilmour *et al.*, 2000), which suggests the functions of members within the same group in the ERF family are likely related to each other, similar to reported MADS-box and bHLH families (Parenicova *et al.*, 2003; Toledo-Ortiz *et al.*, 2003). Thus, the assessment of structural relationships between all the ERF family proteins in plants, as part of each transcription factor function analysis, would provide a guide for predicting the functions of these genes.

Cucumber (*Cucumis sativus* L.), belonging to the *Cucurbitaceae* family, is an economically and nutritionally important vegetable crop cultivated world-wide. Huang *et al.* (2009) proposed the existence of 110 AP2/ERF family genes in the cucumber genome. However, they did not present any specific information regarding individual genes, and no member of the cucumber ERF family has been characterized so far. Furthermore, the expression patterns of this family, as well as details on phylogenetic relationships with ERF members of other plants, remain poorly understood. Thus, the genome-wide identification, and phylogenetic and expression analysis of the family in cucumbers, as well as the comparative analyses with Arabidopsis and rice ERF members, all undertaken here, could be extremely useful in studies on the biological functions of each gene in the cucumber ERF family.

## Material and Methods

### Database search for cucumber *ERF* genes

The AP2/ERF domain of a cucumber ethylene response factor sequence (GenBank number AY792593) was used as a query sequence for TBLASTN (Altschul *et al.*, 1997) searches of AP2/ERF superfamily genes encoded in the cucumber genome. The cucumber genome sequence from Cucumber Genome Initiative (CuGI), obtained and released by The Institute of Vegetables and Flowers, of the Chinese Academy of Agricultural Sciences (IVF-CAAS) was used. Default parameters with the TBLASTN program were wordsize 2 and extension 11. Redundant sequences with the same scaffold or chromosome location were removed from the data set. In addition, we have also obtained the same sequences from the CuGI database using Hidden Markov Model (HMM) analysis with the Pfam number PF00847 containing typical AP2/ERF domain.

To further confirm hypothetical AP2/ERF superfamily genes, the cDNA sequences, first translated into amino-acid sequences, were then searched for the AP2/ERF domain using the Simple Modular Architecture Research Tool (SMART)(Letunic *et al.*, 2004).

### Multiple sequence alignment, tree building and conserved motif prediction

Multiple sequence alignment, using Clustal X (Larkin *et al.*, 2007) with default parameters, was with predicted cucumber *CsERF* protein sequences, with sequential

manual adjustment. Similar amino acids were highlighted using the GeneDoc tool (Nicholas *et al.*, 1997). Multalin software (Corpet, 1988) was also used as a secondary method for aligning sequences and rechecking results. To compare the evolutionary relationships of cucumber, Arabidopsis, and rice ERF family members, multiple sequence alignment was applied, by way of Clustal X, on already obtained *CsERF* protein sequences, and 122 Arabidopsis *AtERF* and 139 rice *OsERF* members predicted by Nakano *et al.* (2006), also with posterior manual adjustment of alignments.

A phylogenetic tree was constructed with aligned *CsERF* protein sequences using MEGA4 (Tamura *et al.*, 2007), and the Neighbor Joining (NJ) method, with Poisson correction, pairwise deletion and bootstrap (1,000 replicates; random seeds), as parameters. Simultaneously, the Maximum Parsimony (MP) method of PHYLIP 3.69 software (Felsenstein., 1989) was employed to create a second phylogenetic tree with a bootstrap of 1,000 replicates, to so validate the results from the NJ method. A combined *CsERF*, *AtERF* and *OsERF* phylogenetic tree was then constructed, also with MEGA4, the NJ method and a bootstrap of 1,000 replicates. The subsequent tree file was visualized by the TreeView1.6.6 tool (Page, 1996).

The MEME tool (Bailey *et al.*, 2003) was used in the search for conserved motifs shared by *CsERF* members, to so identify similar sequences.

### Intron/exon structure, genome distribution, and segmental duplication

For intron/exon structure analysis, the DNA and cDNA sequences corresponding to each predicted gene from BLASTN research and CuGI database annotation, were unloaded, and their intron distribution patterns and splicing phases analyzed, using the GSDS web-based bioinformatics tool.

In order to obtain information on *CsERF* gene location, a map with the distribution of *CsERF* family members throughout the cucumber genome, was drawn with the MapInspect tool. The 100 kb DNA segments flanking each *CsERF* gene were analyzed to detect large segment-duplicated events. Regions on the different linkage groups containing six or more homologous pairs, each with less than 25 nonhomologous intervening genes, were defined as duplicated segments. A gene-pair, separated by less than five intervening genes and sharing  $\geq 40\%$  sequence similarity at the amino acid level, was considered as tandem-duplicated. BioEdit5.0.6 software (Hall., 1999) was used for analyzing homologs for similarity on the NJ phylogenetic tree of these *CsERF* genes.

### Expression analysis of Cucumber *ERF* genes

The Expressed Sequence Tag (EST) was used to detect *CsERF* gene expression patterns. EST data were obtained from 353,941 previously reported high quality EST

sequences (Guo *et al.*, 2010), as well as the ~8,210 cucumber EST sequences available in GenBank. An EST was considered as corresponding to its gene on sharing  $\geq 95\%$  sequence similarity, E values  $\leq 10^{-10}$ , and the length of matching sequences  $\geq 100$  bp. Semi-quantitative RT-PCR was also used to detect the expression patterns of two *CsERF* genes from each group. PCR primers were designed to avoid the conserved region. Information on primer sequences appears in detail in Table S4. Seeds of the 'Chinese long' 9930 inbred line, commonly used in modern cucumber breeding (Huang *et al.*, 2009), were germinated and grown in trays containing a soil mixture (peat: sand: pumice, 1:1:1, v/v/v). Plants were adequately watered and grown at day/night temperatures of 24/18 °C with a 16 h photoperiod. Total RNA of root, stem, leaf, and flower of cucumber at the stage of the 20 main-stem nodes was isolated using the TRIzol Reagent (Invitrogen, USA). RT-PCR was carried out according to manufacturer's recommendations (Tiangen Biotech Co. Ltd, Beijing China). The cucumber *actin* DNA fragment (161 bp) was employed as inner standard for each gene.

## Results and Discussion

### Identification of 103 *CsERF* genes

In order to identify the *CsERF* genes in cucumber genomes, the AP2/ERF domain of a cucumber ethylene response factor sequence (GenBank number AY792593) was used as BLAST query sequence. 131 genes were identified as possibly encoding proteins containing the AP2/ERF domain (Table 1). The same 131 sequences were also obtained from the Cucumber Genome Initiative (CuGI) database, using HMM analysis with PF00847 containing a typical AP2/ERF domain. The individual genes are listed in Table S1. Among these, the 18 predicted to encode proteins containing two AP2/ERF domains, and the 4 to encode one AP2/ERF domain together with one B3 domain, were thus assigned to the AP2 and RAV families, respectively. The remaining 109 genes were all predicted to encode proteins containing a single AP/ERF domain. Among these, 103 were assigned to the ERF family. Of the remaining 6, two, Csa002695 and Csa012456, although also containing a single AP/ERF domain, were distinct from the ERF type and more closely related to the AP2. Hence, they were assigned to the AP2 family. As homology appeared to be quite low in comparison with the others, the remaining 4, viz., Csa020380, Csa005269, Csa013415 and Csa012810, were designated as soloists (Figure S1). Based on the amino acid similarity of AP2/ERF domains, the 103 ERF family members were further classified into two subfamilies, 42 genes encoding CBF/DREB-like proteins were assigned to the CBF/DREB subfamily, and the other 61, encoding ERF-like proteins, to the ERF subfamily. As number designation of the ERF family genes was based on the order of multiple sequence alignments, for study purposes, each was provi-

**Table 1** - Summary of the structure and size of each group of the AP2/ERF superfamily in cucumber, compared with those in Arabidopsis and rice, as classified by Nakano *et al.* (2006). Totals for each family are in bold-type.

Family	Subfamily	Group	Cucumber	Arabidopsis	Rice
AP2			<b>20</b>	<b>18</b>	<b>29</b>
ERF			<b>103</b>	<b>122</b>	<b>139</b>
	DREB		42	57	56
		I	5	10	9
		II	10	16	15
		III	20	22	26
		IV	7	9	6
	ERF		61	65	76
		V	15	12	11
		VI	8	8	6
		VII	3	5	15
		VIII	11	15	13
		IX	16	18	18
		X	8	7	13
		a single group			7
RAV			<b>4</b>	<b>6</b>	<b>5</b>
Soloist			<b>4</b>	<b>1</b>	<b>1</b>
Total			<b>131</b>	<b>147</b>	<b>174</b>


sionally distinguished by a generic name, viz., *CsERF001-CsERF103* (Table S1).

SMART analysis indicated that the AP2/ERF domain of each of the 103 *CsERF* genes was typical, thereby certifying to their reliability.

### Multiple sequence alignments and tree building

To examine sequence features of 103 *CsERF* proteins, we performed a multiple sequence alignment using amino acid sequences of the AP2/ERF domain (Figure S2). The alignment indicated that the residues Gly-4, Arg-6, Arg-8, Trp-38, Gly-40, and Ala-48 were completely conserved (Figure S2). Furthermore, more than 95% of *CsERF*-family members contain Gly-12, Glu-17, Ile-18, Arg-36, Leu-39, Ala-49, Ala-51, and Asp-53 residues (Figure S2).

To determine the evolutionary relationship among *CsERF* proteins, an unrooted NJ phylogenetic tree was constructed, with bootstrap analysis (1000 replicates) based on the multiple sequence alignments of the 103 *CsERF* proteins (Figure 1). The analysis result showed that the 103 *CsERF* members were divided into 10 groups, designated I to X, in accordance with the Arabidopsis ERF gene family classification (Nakano *et al.*, 2006). Detailed information appears in Figure 1 and Table S1. The bootstrapping values for the nodes in this phylogenetic tree were not high in every case, similar to the results of the phylogenetic analysis done on Arabidopsis ERF proteins (Nakano *et al.*, 2006).




**Figure 1** - Phylogenetic analysis of 103 cucumber ERF proteins. An unrooted neighbor-joining phylogenetic tree was constructed using MEGA4 for the multiple sequence alignment of 103 cucumber ERF protein sequences. 10 groups are marked I to X, as described by Nakano *et al.*, (2006).

This is most likely due to the AP2/ERF domain being relatively short, and members within the subfamily highly conserved, with relatively few informative-character positions.

NJ-tree reliability was certified by generating another phylogenetic tree by MP analysis (Figure S3), whereupon it was found that nearly all the *CsERF* members were placed within the same groups.

#### Conserved motifs outside the AP2/ERF domain


Regions outside the DNA-binding domain in transcription factors generally contain either functionally important domains, or motifs associated with transcription-regulation and nuclear localization (Liu *et al.*, 1999). Proteins within a group that share these domains or motifs in a phylogenetic tree are likely to share similar functions. It has been reported that an ERF-associated amphiphilic repression (EAR) motif (DLNxxP) is a repression domain in the C-terminal regions of the repressor-type ERF proteins playing key roles in several biological functions by negatively regulating genes involved in developmental, hormonal and stress signaling pathways (Fujimoto *et al.*, 2000; Ohta *et al.*, 2001). Motif analysis in this study revealed that the EAR motif was only found among proteins within groups II and VIII, as CMII-2 and CMVIII-1 motifs, respectively (Figure 2, Figure 3, Table S2), thus leading us to suspect their involvement in negative-regulation functions. Previous research showed that the Cys repeat sequence-CX<sub>2</sub>CX<sub>4</sub>CX<sub>2-4</sub>C, possibly a zinc-finger motif, plays a part either in DNA binding, or in protein-protein interactions (Nakano *et al.*, 2006). such a consensus sequence was only found within the CMX-2 motif in the N-terminal region of



**Figure 2** - Phylogenetic relationships among *CsERF* proteins from the 10 groups, I to X, in the cucumber ERF family. Only > 50% bootstrap values are shown in this phylogenetic tree. The positions of introns are marked with arrowheads. Each black box represents an AP2/ERF domain. Conserved motifs are marked with different signs, as indicated below the tree.

group X proteins. Liu *et al.* (1999) believed that regions of acidic amino acid-rich, Gln-rich, Pro-rich, and/or Ser/Thr-rich amino acid sequences, can usually be designated as transcriptional-activation domains (Liu *et al.*, 1999). The conserved motifs identified in this study have similar features, such as Gln-rich in group III as the CMIII-7 motif, Pro-rich in group III as the CMIII-2 motif, and/or Ser/Thr-rich in group VIII as the CMVIII-4 motif (Figure 2; Table S2).

In addition, we also found that most of the motifs were selectively distributed among the specific clades in the phylogenetic tree, thereby demonstrating structural similarities among members within the same group (Figure 2, Table S2).



**Figure 3** - The EAR motif-like sequences conserved in group VIII and II ERF proteins. (A) Sequence alignment of C-terminal regions in group VIII proteins. (B) Sequence alignment of C-terminal regions in group II proteins. Conserved motifs are underlined. Black and gray shading indicate identical and conserved amino acid residues present in > 50% of the aligned sequences, respectively. Consensus amino acid residues are given below the alignment. The “x” in the sequence indicates no conservation at this position.

### Structure and evolution of *CsERF* genes

Apparently, most Arabidopsis *ERF* genes do not possess introns (Sakuma *et al.*, 2002). A similar situation also appeared in this study, among the 103 *CsERF* genes, 83 (81%) having no intron, the remaining 20, unevenly distributed in groups I, IV, V, VII and X, having only one or two introns (Figure 2). As shown in Figure 2, 17 of the 20 possess only a single intron, and the other three genes *CsERF002*, *CsERF009* and *CsERF050* possess two. The presence and position of the introns was highly conserved in each group, thus further validating the reliability of the cucumber ERF-family gene classification in this study.

The genomic distribution of the 103 *CsERF* genes was analyzed, in order to acquire an insight into their evolution. With the exception of the seven genes *CsERF008*, *CsERF013*, *CsERF024*, *CsERF052*, *CsERF076*, *CsERF081* and *CsERF085* lying within unassembled scaffold000393, scaffold000131, scaffold000677, scaffold000111, scaffold000379, scaffold000576 and scaffold000118, respectively, the remainder were found to be unevenly distributed among seven chromosomes (Figure 4, Table S1). As indicated in Figure 4, some, clustered

in a large group, within the same small chromosomal region, as, for example, group III members *CsERF087*, *CsERF088* and *CsERF089*, located in a region close to a telomere on chromosome 5, whereas other members of this group were distributed among different chromosomes, *i.e.*, *CsERF083* in chromosome 2 and *CsERF090* in chromosome 4. A similar situation also occurred among *OsERF* and *AtERF* members (Nakano *et al.*, 2006), thereby indicating that *ERF* genes are distributed widely within the genome of the common ancestor of monocots and eudicots.


Although previous research has shown that whole-genome duplication, as a recent event, is not the case with cucumbers, several tandem duplications have in fact actually occurred (Huang *et al.*, 2009), with considerable impact on the increase in the number of family genes in the genome. As the analysis of 100 kb DNA segments flanking each *CsERF* gene indicated that none could have derived from segment duplication, it is most likely that tandem duplication played a crucial role in gene multiplication. According to previously reported results with Arabidopsis, members of groups III and IX play crucial roles in biotic and/or abiotic stress response. Our analyses indicated both to be the two largest groups, gene multiplication in the two possibly having arisen from the higher frequency of tandem duplication, as a means of adapting to various environment changes.

### *In silico* identification of *ERF* genes in Arabidopsis and rice, and a comparative analysis of cucumbers

As a previous report indicated there to be 122 and 139 *ERF* family members distributed within Arabidopsis and rice genomes, respectively (Nakano *et al.*, 2006), Arabidopsis and rice genomes were re-screened for *ERF* sequences, with the subsequent discovery of a further four *AtERF* and six *OsERF* genes, designated as *AtERF123~AtERF126* and *OsERF140~OsERF145* (Table S3), respectively, based on the names of the previous 122 *AtERFs* and 139 *OsERFs*. This discrepancy is probably owing to fresh information on Arabidopsis and rice genome sequences.

To define the evolutionary relationship of cucumber *ERF* family proteins with those of Arabidopsis and rice, an unrooted neighbor-joining (NJ) phylogenetic tree was generated, based on bootstrap analysis (1000 replicates) of multiple sequence alignment of their respective *ERF* members. In addition to the ten groups I to X in Arabidopsis and rice, described by Nakano *et al.* (2006), another group was found, containing four new *AtERF* members clustering with three new *OsERF* members, *viz.*, *OsERF140*, *OsERF142* and *OsERF144* (Figure S4). A further three new *OsERF* members, *viz.*, *OsERF141*, *OsERF143* and *OsERF145*, were placed in groups I, II and VIII, respectively.

Phylogenetic analysis revealed that most *CsERF* members were closer to eudicot *AtERF* members than



**Figure 4** - Chromosomal localization of 103 cucumber *ERF* genes. The scale is in megabases (Mb). The ovals in the middle of the seven chromosomes show the centromeric positions according to the sequencing results of the cucumber genome (Huang *et al.*, 2009). Seven genes *CsERF008*, *CsERF013*, *CsERF024*, *CsERF052*, *CsERF076*, *CsERF081*, and *CsERF085* on scaffold000393, scaffold000131, scaffold000677, scaffold000111, scaffold000379, scaffold000576, and scaffold000118, respectively, could not be anchored onto a specific chromosome.

monocot *OsERFs* in this classification. For example, based on bootstrap values, three group III *CsERF* members (*CsERF071*~*CsERF073*) clustered with six *AtERF* (*AtERF028*~*AtERF033*), whereas another nine *OsERFs* (*OsERF024*~*OsERF031*, *OsERF133*) were branched into a single clade (Figure S4). On closely examining group IV, it was found that only one member, *CsERF070*, was clustered with *OsERF117* and *AtERF052*. As previous studies have shown that *AtERF052* mainly mediates the effects of exogenous trehalose on *Arabidopsis* growth and starch breakdown, and vegetative development by sugar, besides repressing endosperm induced seed germination, *CsERF070* may also participate in these plant-development processes, the detailed function needs further researched and confirmation. At the same time, *CsERF078* was close to *AtERF024* in group III and *CsERF017* to *AtERF081* in group VIII, although the functions of these genes have not, as yet, been rigorously demonstrated.

Usually, the intron/exon position pattern provides clues on evolutionary relationships. Research by Nakano *et al.* (2006) indicated that the position of the intron was conserved in *Arabidopsis ERF* groups V, VII, X, with Xb-L containing only one intron. As with *Arabidopsis* and rice *ERF*, in the present study, it was revealed that both the presence and position of *CsERF* introns in groups IV, V, VII and X were highly conserved, with only one or two exceptions (Figure 2). Thus, besides further validating the classi-

fication of *CsERF* family genes, this was an indication of the conservation of intron position patterns that existed in the common ancestor of monocots and eudicots. On the other hand, and in the same group, introns were observed in only one species, but not in others. For example, two *CsERF* members in group I, namely *CsERF028* and *CsERF031*, possessed one intron at the N-terminal region, although *AtERF* and *OsERF* members in the same group possessed none. A similar situation also occurred in group II, two *OsERF* members, *OsERF015* and *OsERF016*, having one intron, and *CsERF* and *AtERF* members none, thus possibly indicating intron insertion after the divergence of monocots and eudicots.

As described above, proteins within a group that share conserved domains or motifs outside the DNA-binding domain in transcription factors in a phylogenetic tree, are likely to share similar functions. Cheong *et al.* (2003) believed that the CMVII-4 motif, as a putative MAP kinase phosphorylation site in *OsEREBP1*, and the phosphorylation of *OsEREBP1*, resulted in the enhancement of its binding to the GCC box and GCC box-mediated transcriptional activation. Conserved motif analysis showed that the very CMVII-4 motif (CMVII-3 in this study) has been observed in *CsERF025* and *CsERF026* in group VII, *OsERF070*~*OsERF072* in rice, and *AtERF074* and *AtERF075* in *Arabidopsis*. Whether *CsERF025* and

*CsERF026* share similar functions in transcriptional activation regulation, needs to be confirmed.


### Expression analysis of 103 cucumber *ERF* genes

As gene expression patterns are often correlated with their functions, the ESTs, created by partially sequencing randomly isolated gene transcripts, have proved to be invaluable in discovery through expression pattern analysis.

By using data originating from both 353,941 previously reported high quality ESTs (Guo *et al.* 2010) and ~8,210 cucumber ESTs available in GenBank, 41 *CsERF* genes were discovered in at least one tissue among the four investigated, *i.e.*, root, shoot, leaf and flower, the remaining 62 having presented no expression signal. Whereas among the 41 expressed genes, 35, with a high 85% ratio, were found in flowers, two, *CsERF067* and *CsERF072*, of the remaining six were expressed in roots, one, *CsERF067*, in shoots, and, three, *CsERF026*, *CsERF036*, and *CsERF045*, in leaves. The fact that most sequence tag expression corresponded to *CsERF* genes in flowers and little in the other tissues, might be due to the 353,941 EST sequences (98%) having all originated from one flower tissue (Guo *et al.* 2010).

For further study of *CsERF* gene expression patterns, two members of each group were selected for RT-PCR analysis with RNA from roots, stems, leaves and flowers. In general, patterns were conserved within subfamilies, although expression levels of specific members could change in different organs. Similar expression patterns were observed among members belonging to 6 groups of *CsERF* genes (I, III, IV, VII, VIII, X). Among these, the members of 4 groups (I, VII, VIII, X) were expressed wherever investigated, thereby implying that these genes could play regulatory roles in various cucumber tissues. As regards the other two groups, *CsERF067* and *CsERF068* from group IV presented transcript signals in three plant tissues, whereas in group III, *CsERF072* and *CsERF073* transcript signals were detected only in the root and stem, a possible indication of their taking part in specific biological processes in cucumber vegetative development. Given that similar expression patterns were observed in two members in each group, it is speculated that this similarity might also extend to other group-members, pending corroboration by further experiments.

On the other hand, expression patterns in members of the other 4 groups (II, V, VI and IX) were varied. As indicated in Figure 5, in *CsERF02* and *CsERF03* in group V, the high transcript signals detected in three of the four vegetative tissues were different from those expressed in flowers. A similar situation was also found in two members, *CsERF057* and *CsERF058*, in group VI. More obvious variation in gene expression among group members could be observed in the remaining groups, II and IX. Group II member *CsERF087* presented transcripts in the stem, leaf and flower, whereas for the other member, *CsERF089*, no



**Figure 5** - Expression analysis of 20 cucumber *ERF* genes in different tissues using RT-PCR. RT-PCR was with primers specific for the 20 *CsERF* genes. PCR products were run on 1.5% agarose gels. *CsACTIN* primers giving a 161 bp product were used as inner standard for each gene. Sample identities are as follows: root (R), stem (S), leaf (L) and flower (F).

detectable signal was observed anywhere. In group IX, *CsERF051* was highly expressed in all the tissues, whereas the other member, *CsERF048*, was expressed only in the root and stem. The different expression patterns among these 4 groups could imply the existence of a probable intragroup functional divergence.

In summary, after extensive analysis, 103 *CsERF* genes were compared with 126 Arabidopsis and 145 rice *ERF* genes. The 103 *CsERF* genes were divided into 10 groups, I to X, thus in general accordance with previous studies (Nakano *et al.*, 2006). This classification was based on the presence and position of the introns and the conserved amino acid sequence motifs outside the AP2/ERF domain. Chromosomal localization and genome distribution revealed that tandem duplication may have contributed to *CsERF*-gene expansion. Expression data revealed the widespread distribution of this gene family within cucumber plant tissues. Furthermore, in most of the groups, two different members presented similar expression patterns, whereby a possible basis for functional analysis to discover the role of *CsERF* genes in cucumber development.

### Acknowledgments

The authors wish to thank Zhonghua Zhang for help with statistical analysis. This work was supported by the National Natural Science Foundation of China (31060262), the Natural Science Foundation of Jiangxi, China (2009GQN0034) and the Foundation of Jiangxi Agricultural University, Jiangxi, China (2009).

### References

- Aharoni A, Dixit S, Jetter R, Thoenes E, van Arkel G and Pereira A (2004) The SHINE clade of AP2 domain transcription factors activates wax biosynthesis, alters cuticle properties,

- and confers drought tolerance when overexpressed in *Arabidopsis*. *Plant Cell* 16:2463-2480.
- Altschul SF, Madden TL, Schaffer AA, Zhang J, Zhang Z, Miller W and Lipman DJ (1997) Gapped BLAST and PSI-BLAST: A new generation of protein database search programs. *Nucleic Acids Res* 25:3389-3402.
- Aukerman MJ and Sakai H (2003) Regulation of flowering time and floral organ identity by a MicroRNA and its APE-TALA2-like target genes. *Plant Cell* 15:2730-2741.
- Bailey PC, Martin C, Toledo-Ortiz G, Quail PH, Huq E, Heim MA, Jakoby M, Werber M and Weisshaar B (2003) Update on the basic helix-loop-helix transcription factor gene family in *Arabidopsis thaliana*. *Plant Cell* 15:2497-2502.
- Boutillier K, Offringa R, Sharma VK, Kieft H, Ouellet T, Zhang L, Hattori J, Liu CM, van Lammeren AA, Miki BL, *et al.* (2002) Ectopic expression of BABY BOOM triggers a conversion from vegetative to embryonic growth. *Plant Cell* 14:1737-1749.
- Broun P, Poindexter P, Osborne E, Jiang CZ and Riechmann JL (2004) WIN1, a transcriptional activator of epidermal wax accumulation in *Arabidopsis*. *Proc Natl Acad Sci USA* 101:4706-4711.
- Brown RL, Kazan K, McGrath KC, Maclean DJ and Manners JM (2003) A role for the GCC-box in jasmonate-mediated activation of the PDF1.2 gene of *Arabidopsis*. *Plant Physiol* 132:1020-1032.
- Cao Y, Song F, Goodman RM and Zheng Z (2006) Molecular characterization of four rice genes encoding ethylene-responsive transcriptional factors and their expressions in response to biotic and abiotic stress. *J Plant Physiol* 163:1167-1178.
- Che P, Lall S, Nettleton D and Howell SH (2006) Gene expression programs during shoot, root, and callus development in *Arabidopsis* tissue culture. *Plant Physiol* 141:620-637.
- Cheong YH, Moon BC, Kim JK, Kim CY, Kim MC, Kim IH, Park CY, Kim JC, Park BO, Koo SC, *et al.* (2003) BWMK1, a rice mitogen-activated protein kinase, locates in the nucleus and mediates pathogenesis-related gene expression by activation of a transcription factor. *Plant Physiol* 132:1961-1972.
- Corpet F (1988) Multiple sequence alignment with hierarchical clustering. *Nucleic Acids Res* 16:10881-10890.
- Dong CJ and Liu JY (2010) The *Arabidopsis* EAR-motif-containing protein RAP2.1 functions as an active transcriptional repressor to keep stress responses under tight control. *BMC Plant Biol* 10:e47.
- Elliott RC, Betzner AS, Huttner E, Oakes MP, Tucker WQ, Gerentes D, Perez P and Smyth DR (1996) *AINTEGUMENTA*, an *APETALA2*-like gene of *Arabidopsis* with pleiotropic roles in ovule development and floral organ growth. *Plant Cell* 8:155-168.
- Felsenstein J (1989) PHYLIP: Phylogeny Inference Package. *Cladistics* 5:164-166.
- Fujimoto SY, Ohta M, Usui A, Shinshi H and Ohme-Takagi M (2000) *Arabidopsis* ethylene-responsive element binding factors act as transcriptional activators or repressors of GCC box-mediated gene expression. *Plant Cell* 12:393-404.
- Gilmour SJ, Sebolt AM, Salazar MP, Everard JD and Thomashow MF (2000) Overexpression of the *Arabidopsis* CBF3 transcriptional activator mimics multiple biochemical changes associated with cold acclimation. *Plant Physiol* 124:1854-1865.
- Guo S, Zheng Y, Joung JG, Liu S, Zhang Z, Crasta OR, Sobral BW, Xu Y, Huang S and Fei Z (2010) Transcriptome sequencing and comparative analysis of cucumber flowers with different sex types. *BMC Genomics* 11:e384.
- Hall TA (1999) BioEdit: A user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symp Ser* 41:95-98.
- Hinz M, Wilson IW, Yang J, Buerstenbinder K, Llewellyn D, Dennis ES, Sauter M and Dolferus R (2010) *Arabidopsis* RAP2.2: An ethylene response transcription factor that is important for hypoxia survival. *Plant Physiol* 153:757-772.
- Hu YX, Wang YX, Liu XF and Li JY (2004) *Arabidopsis* RAV1 is down-regulated by brassinosteroid and may act as a negative regulator during plant development. *Cell Res* 14:8-15.
- Huang S, Li R, Zhang Z, Li L, Gu X, Fan W, Lucas WJ, Wang X, Xie B, Ni P, *et al.* (2009) The genome of the cucumber, *Cucumis sativus* L. *Nat Genet* 41:1275-1281.
- Ikeda Y, Banno H, Niu QW, Howell SH and Chua NH (2006) The ENHANCER OF SHOOT REGENERATION 2 gene in *Arabidopsis* regulates CUP-SHAPED COTYLEDON 1 at the transcriptional level and controls cotyledon development. *Plant Cell Physiol* 47:1443-1456.
- Iwase A, Mitsuda N, Koyama T, Hiratsu K, Kojima M, Arai T, Inoue Y, Seki M, Sakakibara H, Sugimoto K, *et al.* (2011) The AP2/ERF transcription factor WIND1 controls cell dedifferentiation in *Arabidopsis*. *Curr Biol* 21:508-514.
- Jin LG and Liu JY (2008) Molecular cloning, expression profile and promoter analysis of a novel ethylene responsive transcription factor gene GhERF4 from cotton (*Gossypium hirsutum*). *Plant Physiol Biochem* 46:46-53.
- Larkin MA, Blackshields G, Brown NP, Chenna R, McGettigan PA, McWilliam H, Valentin F, Wallace IM, Wilm A, Lopez R, *et al.* (2007) Clustal W and Clustal X v. 2.0. *Bioinformatics* 23:2947-2948.
- Letunic I, Copley RR, Schmidt S, Ciccarelli FD, Doerks T, Schultz J, Ponting CP and Bork P (2004) SMART 4.0: Towards genomic data integration. *Nucleic Acids Res* 32:D142-D144.
- Li XP, Tian AG, Luo GZ, Gong ZZ, Zhang JS and Chen SY (2005) Soybean DRE-binding transcription factors that are responsive to abiotic stresses. *Theor Appl Genet* 110:1355-1362.
- Licausi F, Giorgi FM, Zenoni S, Osti F, Pezzotti M and Perata P (2010) Genomic and transcriptomic analysis of the AP2/ERF superfamily in *Vitis vinifera*. *BMC Genomics* 11:e719.
- Lim CJ, Hwang JE, Chen H, Hong JK, Yang KA, Choi MS, Lee KO, Chung WS, Lee SY and Lim CO (2007) Overexpression of the *Arabidopsis* DRE/CRT-binding transcription factor DREB2C enhances thermotolerance. *Biochem Biophys Res Commun* 362:431-436.
- Lin RC, Park HJ and Wang HY (2008) Role of *Arabidopsis* RAP2.4 in regulating light- and ethylene-mediated developmental processes and drought stress tolerance. *Mol Plant* 1:42-57.
- Liu L, White MJ and MacRae TH (1999) Transcription factors and their genes in higher plants functional domains, evolution and regulation. *Eur J Biochem* 262:247-257.



- Liu Q, Kasuga M, Sakuma Y, Abe H, Miura S, Yamaguchi-Shinozaki K and Shinozaki K (1998) Two transcription factors, DREB1 and DREB2, with an EREBP/AP2 DNA binding domain separate two cellular signal transduction pathways in drought- and low-temperature-responsive gene expression, respectively, in *Arabidopsis*. *Plant Cell* 10:1391-1406.
- Moose SP and Sisco PH (1996) Glossy15, an APETALA2-like gene from maize that regulates leaf epidermal cell identity. *Genes Dev* 10:3018-3027.
- Nag A and Yang Y and Jack T (2007) DORNROSCHEN-LIKE, an AP2 gene, is necessary for stamen emergence in *Arabidopsis*. *Plant Mol Biol* 65:219-232.
- Nakano T, Suzuki K, Fujimura T and Shinshi H (2006) Genome-wide analysis of the ERF gene family in *Arabidopsis* and rice. *Plant Physiol* 140:411-432.
- Nakashima K, Shinwari ZK, Sakuma Y, Seki M, Miura S, Shinozaki K and Yamaguchi-Shinozaki K (2000) Organization and expression of two *Arabidopsis* DREB2 genes encoding DRE-binding proteins involved in dehydration- and high-salinity-responsive gene expression. *Plant Mol Biol* 42:657-665.
- Nicholas KB, Nicholas Jr HB and Deerfield II DW (1997) GeneDoc: Analysis and visualization of genetic variation. *Embnew News* 4:14.
- Ogawa T, Pan L, Kawai-Yamada M, Yu LH, Yamamura S, Koyama T, Kitajima S, Ohme-Takagi M, Sato F and Uchimiya H (2005) Functional analysis of *Arabidopsis* ethylene-responsive element binding protein conferring resistance to Bax and abiotic stress-induced plant cell death. *Plant Physiol* 138:1436-1445.
- Ohta M, Matsui K, Hiratsu K, Shinshi H and Ohme-Takagi M (2001) Repression domains of class II ERF transcriptional repressors share an essential motif for active repression. *Plant Cell* 13:1959-1968.
- Page RD (1996) TreeView: An application to display phylogenetic trees on personal computers. *Comput Appl Biosci* 12:357-358.
- Parenticova L, de Folter S, Kieffer M, Horner DS, Favalli C, Busscher J, Cook HE, Ingram RM, Kater MM, Davies B, *et al.* (2003) Molecular and phylogenetic analyses of the complete MADS-box transcription factor family in *Arabidopsis*: New openings to the MADS world. *Plant Cell* 15:1538-1551.
- Pre M, Atallah M, Champion A, De Vos M, Pieterse CM and Memelink J (2008) The AP2/ERF domain transcription factor ORA59 integrates jasmonic acid and ethylene signals in plant defense. *Plant Physiol* 147:1347-1357.
- Qin F, Sakuma Y, Tran LS, Maruyama K, Kidokoro S, Fujita Y, Fujita M, Umezawa T, Sawano Y, Miyazono K, *et al.* (2008) *Arabidopsis* DREB2A-interacting proteins function as RING E3 ligases and negatively regulate plant drought stress-responsive gene expression. *Plant Cell* 20:1693-1707.
- Rashotte AM, Mason MG, Hutchison CE, Ferreira FJ, Schaller GE and Kieber JJ (2006) A subset of *Arabidopsis* AP2 transcription factors mediates cytokinin responses in concert with a two-component pathway. *Proc Natl Acad Sci USA* 103:11081-11085.
- Riechmann JL, Heard J, Martin G, Reuber L, Jiang C, Keddie J, Adam L, Pineda O, Ratcliffe OJ, Samaha RR, *et al.* (2000) *Arabidopsis* transcription factors: Genome-wide comparative analysis among eukaryotes. *Science* 290:2105-2110.
- Sakuma Y, Liu Q, Dubouzet JG, Abe H, Shinozaki K and Yamaguchi-Shinozaki K (2002) DNA-binding specificity of the ERF/AP2 domain of *Arabidopsis* DREBs, transcription factors involved in dehydration- and cold-inducible gene expression. *Biochem Biophys Res Commun* 290:998-1009.
- Schmid M, Uhlenhaut NH, Godard F, Demar M, Bressan R, Weigel D and Lohmann JU (2003) Dissection of floral induction pathways using global expression analysis. *Development* 130:6001-6012.
- Shaikhali J, Heiber I, Seidel T, Stroher E, Hiltcher H, Birkmann S, Dietz KJ and Baier M (2008) The redox-sensitive transcription factor Rap2.4a controls nuclear expression of 2-Cys peroxiredoxin A and other chloroplast antioxidant enzymes. *BMC Plant Biol* 8:e48.
- Sharma MK, Kumar R, Solanke AU, Sharma R, Tyagi AK and Sharma AK (2010) Identification, phylogeny, and transcript profiling of ERF family genes during development and abiotic stress treatments in tomato. *Mol Genet Genomics* 284:455-475.
- Sharoni AM, Nuruzzaman M, Satoh K, Shimizu T, Kondoh H, Sasaya T, Choi IR, Omura T and Kikuchi S (2011) Gene structures, classification and expression models of the AP2/EREBP transcription factor family in rice. *Plant Cell Physiol* 52:344-360.
- Solano R, Stepanova A, Chao Q and Ecker JR (1998) Nuclear events in ethylene signaling: A transcriptional cascade mediated by ETHYLENE-INSENSITIVE3 and ETHYLENE-RESPONSE-FACTOR1. *Genes Dev* 12:3703-3714.
- Stockinger EJ, Gilmour SJ and Thomashow MF (1997) *Arabidopsis thaliana* CBF1 encodes an AP2 domain-containing transcriptional activator that binds to the C-repeat/DRE, a cis-acting DNA regulatory element that stimulates transcription in response to low temperature and water deficit. *Proc Natl Acad Sci USA* 94:1035-1040.
- Tamura K, Dudley J, Nei M and Kumar S (2007) MEGA4: Molecular Evolutionary Genetics Analysis (MEGA) software v. 4.0. *Mol Biol Evol* 24:1596-1599.
- Toledo-Ortiz G, Huq E and Quail PH (2003) The *Arabidopsis* basic/helix-loop-helix transcription factor family. *Plant Cell* 15:1749-1770.
- van der Fits L and Memelink J (2000) ORCA3, a jasmonate-responsive transcriptional regulator of plant primary and secondary metabolism. *Science* 289:295-297.
- Vernie T, Moreau S, de Billy F, Plet J, Combier JP, Rogers C, Oldroyd G, Frugier F, Niebel A and Gamas P (2008) EFD Is an ERF transcription factor involved in the control of nodule number and differentiation in *Medicago truncatula*. *Plant Cell* 20:2696-2713.
- Welsch R, Maass D, Voegel T, Dellapenna D and Beyer P (2007) Transcription factor RAP2.2 and its interacting partner SINAT2: Stable elements in the carotenogenesis of *Arabidopsis* leaves. *Plant Physiol* 145:1073-1085.
- Wessler SR (2005) Homing into the origin of the AP2 DNA binding domain. *Trends Plant Sci* 10:54-56.
- Xu H, Wang X and Chen J (2010) Overexpression of the Rap2.4f transcriptional factor in *Arabidopsis* promotes leaf senescence. *Sci China Life Sci* 53:1221-1226.
- Zhang JY, Broeckling CD, Blancaflor EB, Sledge MK, Sumner LW and Wang ZY (2005) Overexpression of WXP1, a puta-

tive *Medicago truncatula* AP2 domain-containing transcription factor gene, increases cuticular wax accumulation and enhances drought tolerance in transgenic alfalfa (*Medicago sativa*). *Plant J* 42:689-707.

Zhuang J, Cai B, Peng RH, Zhu B, Jin XF, Xue Y, Gao F, Fu XY, Tian YS, Zhao W, *et al.* (2008) Genome-wide analysis of the AP2/ERF gene family in *Populus trichocarpa*. *Biochem Biophys Res Commun* 371:468-474.

Zhu Q, Zhang J, Gao X, Tong J, Xiao L, Li W and Zhang H (2010) The Arabidopsis AP2/ERF transcription factor RAP2.6 participates in ABA, salt and osmotic stress responses. *Gene* 457:1-12.

## Internet Resources

Cucumber Genome Initiative (CuGI), <http://cucumber.genomics.org.cn> (October 15, 2010).

Clustal X, <http://www.clustal.org> (October 20, 2010).

GeneDoc tool, <http://www.nrbsc.org/gfx/genedoc/> (October 20, 2010).

Multalin software, <http://multalin.toulouse.inra.fr/multalin/multalin.html> (October 20, 2010).

MEGA4, <http://www.megasoftware.net/index.html> (October 20, 2010).

PHYMLIP 3.69, <http://evolution.genetics.washington.edu/phymlip.html> (October 20, 2010).

TreeView1.6.6, <http://taxonomy.zoology.gla.ac.uk/rod/treeview.html> (October 20, 2010).

MEME, [http://meme.sdsc.edu/meme4\\_4\\_0/cgi-bin/meme.cgi](http://meme.sdsc.edu/meme4_4_0/cgi-bin/meme.cgi) (October 20, 2010).

GSDS, <http://gsds.cbi.pku.edu.cn/> (October 20, 2010).

MapInspect, [http://www.plantbreeding.wur.nl/UK/software\\_mapinspect.html](http://www.plantbreeding.wur.nl/UK/software_mapinspect.html) (October 20, 2010).

BioEdit5.0.6, <http://www.mbio.ncsu.edu/BioEdit/bioedit.html> (October 20, 2010).

TIGR Arabidopsis annotation deduced protein database, <http://www.tigr.org/tdb/e2k1/ath1/> (October 15, 2010).

Rice genome, release version 5 of TIGR pseudomolecules, <http://rice.plantbiology.msu.edu/> (October 15, 2010).

## Supplementary Material

The following material is available for this article:

Figure S1 - Multiple sequence alignment of the AP2/ERF domain.

Figure S2 - Multiple sequence alignment of the AP2/ERF domains.

Figure S3 - Phylogenetic analysis of 103 cucumber ERF proteins.

Figure S4 - Comparative phylogenetic analysis of cucumber ERFs with those of Arabidopsis and rice.

Table S1 - The *CsERF* genes identified in this study.

Table S2 - Summary of conserved motifs (CMs) within the *CsERF* family.

Table S3 - New *AtERF* and *OsERF* genes identified in this study.

Table S4 - Information on the primers used in RT-PCR reactions.


This material is available as part of the online article from <http://www.scielo.br/gmb>.

*Associate Editor: Adriana Silva Hemerly*


License information: This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.







**Figure S3** - Phylogenetic analysis of 103 cucumber ERF proteins. A Maximum Parsimony (MP) tree was constructed using PHYLIP3.69, by multiple sequence alignment of 103 cucumber ERF protein sequences. 10 groups are marked I to X, as described by Nakano *et al.*, (2006).



**Figure S4** - Comparative phylogenetic analysis of cucumber ERFs with those of Arabidopsis and rice. An unrooted neighbor-joining phylogenetic tree was constructed using MEGA4, by multiple sequence alignment. The 10 major groups are marked I to X, as described by Nakano *et al.*, (2006).

**Table S1** The *CsERF* genes identified in this study are listed according to their *CsERF* numbers

defined by multiple sequence alignment (Figure S2).

Serial No.	Gene Name	Gene ID	CuGI(5'-3')	Length of chr (bp)	Chromosomal location	Length (a.a.)	No. of Introns	Group name
1	<i>CsERF001</i>	Csa002836	1350153-1349175	22434855	4	180	1	V
2	<i>CsERF002</i>	Csa014841	15208734-15209357	22434855	4	138	2	V
3	<i>CsERF003</i>	Csa005862	18494975-18495575	22000590	2	164	1	V
4	<i>CsERF004</i>	Csa001957	15271088-15271957	17370973	7	196	1	V
5	<i>CsERF005</i>	Csa005861	18488382-18489011	22000590	2	175	1	V
6	<i>CsERF006</i>	Csa006854	631812-630472	22000590	2	246	1	V
7	<i>CsERF007</i>	Csa002035	31922812-31923895	34445220	3	194	1	V
8	<i>CsERF008</i>	Csa022657	8462-9248	13703	Scaffold000393	223	1	V
9	<i>CsERF009</i>	Csa003049	1527385-1528304	22434855	4	190	2	V
10	<i>CsERF010</i>	Csa009694	26544766-26545851	28091742	5	361	0	V
11	<i>CsERF011</i>	Csa015404	19399419-19400496	28091742	5	212	0	V
12	<i>CsERF012</i>	Csa020206	8143974-8143339	22434855	4	211	0	V
13	<i>CsERF013</i>	Csa019531	316141-316866	319939	Scaffold000131	241	0	V
14	<i>CsERF014</i>	Csa007982	17124858-17125502	26547709	6	214	0	VIII
15	<i>CsERF015</i>	Csa011614	9915259-9915918	26547709	6	219	0	VIII
16	<i>CsERF016</i>	Csa019805	3390113-3389469	34445220	3	214	0	VIII
17	<i>CsERF017</i>	Csa016336	7651209-7651703	25430643	1	164	0	VIII
18	<i>CsERF018</i>	Csa015626	19286010-19286591	22434855	4	193	0	VIII
19	<i>CsERF019</i>	Csa003390	2180880-2180251	17370973	7	209	0	VIII
20	<i>CsERF020</i>	Csa016229	7666682-7666047	25430643	1	211	0	VIII
21	<i>CsERF021</i>	Csa000861	21263555-21264580	28091742	5	341	0	VIII
22	<i>CsERF022</i>	Csa011398	14429466-14430542	22000590	2	358	0	VIII
23	<i>CsERF023</i>	Csa003016	1119240-1119923	22434855	4	227	0	VIII
24	<i>CsERF024</i>	Csa023463	3481-4167	6853	Scaffold000677	228	0	VIII
25	<i>CsERF025</i>	Csa009872	16743027-16740950	17370973	7	389	1	VII
26	<i>CsERF026</i>	Csa013360	22272823-22271143	34445220	3	370	1	VII
27	<i>CsERF027</i>	Csa002799	1832463-1831545	22434855	4	272	1	VII
28	<i>CsERF028</i>	Csa007704	7871002-7869906	26547709	6	337	1	I
29	<i>CsERF029</i>	Csa003368	2420250-2419057	17370973	7	397	0	I
30	<i>CsERF030</i>	Csa000642	24948986-24949513	26547709	6	175	0	IX
31	<i>CsERF031</i>	Csa012067	22481978-22481499	26547709	6	250	1	I
32	<i>CsERF032</i>	Csa011099	14386960-14385884	22434855	4	358	0	I
33	<i>CsERF033</i>	Csa006177	9209327-9208125	22000590	2	400	0	I
34	<i>CsERF034</i>	Csa016672	9166496-9167080	22434855	4	194	0	IX
35	<i>CsERF035</i>	Csa012537	1252601-1253083	28091742	5	160	0	IX
36	<i>CsERF036</i>	Csa012536	1247100-1247813	28091742	5	237	0	IX
37	<i>CsERF037</i>	Csa017104	1586163-1585129	34445220	3	344	0	IX
38	<i>CsERF038</i>	Csa002749	2287694-2286942	22434855	4	250	0	IX
39	<i>CsERF039</i>	Csa017144	1669749-1670588	34445220	3	279	0	IX

40	<i>CsERF040</i>	Csa000407	6006431-6005958	34445220	3	157	0	IX
41	<i>CsERF041</i>	Csa015122	20933066-20933446	34445220	3	126	0	IX
42	<i>CsERF042</i>	Csa000150	8965136-8964735	34445220	3	133	0	IX
43	<i>CsERF043</i>	Csa019346	11809343-11809789	17370973	7	148	0	IX
44	<i>CsERF044</i>	Csa022480	10161945-10162631	22000590	2	228	0	IX
45	<i>CsERF045</i>	Csa000151	8945006-8944320	34445220	3	228	0	IX
46	<i>CsERF046</i>	Csa001354	18506357-18505683	34445220	3	224	0	IX
47	<i>CsERF047</i>	Csa019306	11796883-11796188	17370973	7	231	0	IX
48	<i>CsERF048</i>	Csa010078	20115512-20116789	26547709	6	338	1	X
49	<i>CsERF049</i>	Csa000408	6003134-6002700	34445220	3	144	0	IX
50	<i>CsERF050</i>	Csa005593	1951916-1955219	26547709	6	364	2	X
51	<i>CsERF051</i>	Csa008584	4931216-4930683	26547709	6	177	0	X
52	<i>CsERF052</i>	Csa018661	326047-327680	411159	Scaffold000111	285	1	X
53	<i>CsERF053</i>	Csa008878	4066091-4064303	26547709	6	404	1	X
54	<i>CsERF054</i>	Csa010237	20921657-20921076	26547709	6	193	0	VII
55	<i>CsERF055</i>	Csa003542	358765-357989	17370973	7	258	0	VII
56	<i>CsERF056</i>	Csa008749	13733610-13732786	34445220	3	274	0	VII
57	<i>CsERF057</i>	Csa008966	2537686-2538642	26547709	6	318	0	VI
58	<i>CsERF058</i>	Csa001015	23085965-23086894	28091742	5	309	0	VI
59	<i>CsERF059</i>	Csa020336	11118158-11119108	22434855	4	316	0	VI
60	<i>CsERF060</i>	Csa012752	15983553-15984509	28091742	5	318	0	VI
61	<i>CsERF061</i>	Csa019969	16219410-16220390	34445220	3	326	0	VI
62	<i>CsERF062</i>	Csa015455	19016343-19015537	28091742	5	268	0	VI
63	<i>CsERF063</i>	Csa007656	8048779-8049393	26547709	6	204	0	VI
64	<i>CsERF064</i>	Csa017883	30847250-30846696	34445220	3	184	0	IV
65	<i>CsERF065</i>	Csa003622	16252055-16252693	22000590	2	212	0	IV
66	<i>CsERF066</i>	Csa005696	1334610-1333345	26547709	6	421	0	IV
67	<i>CsERF067</i>	Csa007621	7427734-7428273	26547709	6	179	0	IV
68	<i>CsERF068</i>	Csa018854	12152362-12151250	22434855	4	370	0	IV
69	<i>CsERF069</i>	Csa002451	27503953-27503030	34445220	3	230	1	IV
70	<i>CsERF070</i>	Csa005700	1276290-1275373	26547709	6	305	0	IV
71	<i>CsERF071</i>	Csa012414	522938-522336	28091742	5	200	0	III
72	<i>CsERF072</i>	Csa008861	11874380-11873739	34445220	3	213	0	III
73	<i>CsERF073</i>	Csa012870	15262990-15262313	28091742	5	225	0	III
74	<i>CsERF074</i>	Csa004675	25841918-25841304	34445220	3	204	0	III
75	<i>CsERF075</i>	Csa012869	15277531-15276773	28091742	5	252	0	III
76	<i>CsERF076</i>	Csa022610	8225-7716	15472	Scaffold000379	169	0	III
77	<i>CsERF077</i>	Csa008860	11887175-11886597	34445220	3	192	0	III
78	<i>CsERF078</i>	Csa005091	25834056-25834607	34445220	3	183	0	III
79	<i>CsERF079</i>	Csa000388	6224581-6224039	34445220	3	180	0	III
80	<i>CsERF080</i>	Csa008546	5675974-5675333	26547709	6	213	0	III
81	<i>CsERF081</i>	Csa023237	1739-1134	8810	Scaffold000576	201	0	III
82	<i>CsERF082</i>	Csa009843	17099285-17098680	17370973	7	201	0	III
83	<i>CsERF083</i>	Csa021087	17482290-17481601	22000590	2	229	0	III



84	<i>CsERF084</i>	Csa002191	33648340-33647657	34445220	3	227	0	III
85	<i>CsERF085</i>	Csa018998	559702-560208	651424	Scaffold000118	168	0	III
86	<i>CsERF086</i>	Csa017112	1509830-1509186	34445220	3	214	0	III
87	<i>CsERF087</i>	Csa009589	27747151-27746573	28091742	5	192	0	II
88	<i>CsERF088</i>	Csa009590	27726472-27725981	28091742	5	163	0	II
89	<i>CsERF089</i>	Csa009622	27355787-27355296	28091742	5	163	0	II
90	<i>CsERF090</i>	Csa003012	1044217-1044711	22434855	4	164	0	II
91	<i>CsERF091</i>	Csa012919	19140452-19139841	25430643	1	203	0	II
92	<i>CsERF092</i>	Csa013079	14191691-14192332	26547709	6	213	0	II
93	<i>CsERF093</i>	Csa009304	22349007-22348447	22434855	4	186	0	II
94	<i>CsERF094</i>	Csa008952	2719822-2719217	26547709	6	201	0	II
95	<i>CsERF095</i>	Csa011091	14097006-14097704	22434855	4	232	0	II
96	<i>CsERF096</i>	Csa011471	15315130-15315558	22000590	2	142	0	II
97	<i>CsERF097</i>	Csa009478	21425457-21425906	22434855	4	149	0	II
98	<i>CsERF098</i>	Csa020278	12479634-12480113	22000590	2	159	0	II
99	<i>CsERF099</i>	Csa010224	21063174-21062653	26547709	6	173	0	II
100	<i>CsERF100</i>	Csa005900	19222095-19222541	22000590	2	148	0	II
101	<i>CsERF101</i>	Csa005450	2122879-2123478	25430643	1	199	0	V
102	<i>CsERF102</i>	Csa000299	7196619-7195888	34445220	3	243	0	V
103	<i>CsERF103</i>	Csa004614	4193219-4194205	25430643	1	328	0	VI

**Table S2** Summary of conserved motifs (CMs) within the CsERF family

Serial No.	Gene Name	Gene ID	CuGI(5'-3')	Length of chr (bp)	Chromosomal location	Length(a.a.)	No. of Introns
1	<i>AtERF123</i>	At2g39250	16398009-16396740	19705359	2	222	4
2	<i>AtERF124</i>	At2g41710	17409981-17407352	19705359	2	423	7
3	<i>AtERF125</i>	At3g54990	20387499-20386013	23470805	3	247	4
4	<i>AtERF126</i>	At5g60120	24225379-24228293	26992728	5	485	8

**Table S3** New *AtERF* and *OsERF* genes identified in this study are listed according to their *CsERF*

numbers.

Serial No.	Gene Name	Gene ID	CuGI(5'-3')	Length of chr (bp)	Chromosomal location	Length(a.a.)	No. of Introns
1	<i>OsERF140</i>	LOC_Os02g29550	17571123-17576025	35930381	2	222	5
2	<i>OsERF141</i>	LOC_Os02g42585	25591508-25590087	35930381	2	473	0
3	<i>OsERF142</i>	LOC_Os05g32270	18749964-18754112	29894789	5	348	7
4	<i>OsERF143</i>	LOC_Os06g09717	4959148-4959771	31246789	6	207	0
5	<i>OsERF144</i>	LOC_Os10g26590	13791722-13792405	23134759	10	227	0
6	<i>OsERF145</i>	LOC_Os12g40960	25326352-25326768	27497214	12	138	0

**Table S4** Information on the primers used in RT-PCR reactions.

Primer name	Sequence (5' to 3')	Size (bp)
<i>CsERF002_1F</i>	GGGTTTCGAAATTCGTCAT	215
<i>CsERF002_1R</i>	CACCGCCGGTATAATCTCCT	
<i>CsERF003_1F</i>	GAGGCGTTCGTCAAAGACAT	239
<i>CsERF003_1R</i>	ACGGTGGAGTTTAGCAGTCA	
<i>CsERF014_1F</i>	ATTCCATTCTCGCAGCAGTT	361
<i>CsERF014_1R</i>	ACAGAGCAGTGGCATGAAGA	
<i>CsERF015_1F</i>	TGAGTGCGTTCAGGCTAATA	256
<i>CsERF015_1R</i>	GTCCATCGGAGGTGGTAAGT	
<i>CsERF025_1F</i>	TTCAAGGATGATTCCGATTT	216
<i>CsERF025_1R</i>	GATTCTGCTGCCCATTTAC	
<i>CsERF027_1F</i>	TCCGACATTGGCCTAACTC	240
<i>CsERF027_1R</i>	TTCCTCAGCCGTATTGAAAG	
<i>CsERF028_1F</i>	GAAGAAGAAGCCTCACAAGC	270
<i>CsERF028_1R</i>	TGGGCAAGATGAAGAAGAAG	
<i>CsERF030_1F</i>	GGGGTAAATACGCTGCCGAGAT	216
<i>CsERF030_1R</i>	GCTTCCTCTTCGACGGTGGG	
<i>CsERF031_1F</i>	AGACTCTGGCTTGGGACCTT	223
<i>CsERF031_1R</i>	ACTCCTTCACCCATTGAGA	
<i>CsERF034_1F</i>	GAAAGTCGCCGTGGTAGAGG	183
<i>CsERF034_1R</i>	AACGCCGCACAGTCATAAGC	
<i>CsERF048_1F</i>	TCTTCCTCTGTTCTGCTCCCG	182
<i>CsERF048_1R</i>	ACCGTTGCCGACCCTGTTTG	
<i>CsERF051_1F</i>	CCTGCCAGTATTGCTGCTCC	254
<i>CsERF051_1R</i>	TCACCGTAACGATCCTGATCTTCT	
<i>CsERF057_1F</i>	ATCAATGGATGTA ACTACCAGCAC	247
<i>CsERF057_1R</i>	GTAGGCTTGGGAGGCTTCTT	
<i>CsERF058_1F</i>	GAAGAAGCAAAGGGAGTGG	337
<i>CsERF058_1R</i>	TGGGTTTCATCATCAGCGAGT	
<i>CsERF067_1F</i>	AAATTCGAGCACCGAATAGA	287
<i>CsERF067_1R</i>	AACTGCACAATCCTCAGACAG	
<i>CsERF068_1F</i>	AGAAAGGCACTCGTTTATGG	200
<i>CsERF068_1R</i>	CGAAGAAGAAGAGGAGGAAG	
<i>CsERF072_1F</i>	TGTCTCAATTCGCCGATTC	242
<i>CsERF072_1R</i>	GCCATATCCTCCAACAATCC	
<i>CsERF073_1F</i>	GAGCCGAATAAGAAGACGAG	352
<i>CsERF073_1R</i>	AAGATGGAGGAGATAGCAGTAG	
<i>CsERF087_1F</i>	GAACCAGGAAAGAAGACGAGG	120
<i>CsERF087_1R</i>	TGGGAAGTTGAGGCGAGCAT	
<i>CsERF089_1F</i>	ATTCTCAACTCCCTGACTCTGT	225
<i>CsERF089_1R</i>	GCCAAGTAGACCCAACCTCA	
<i>CsACTIN_F</i>	GACATTCAATGTGCCTGCTATG	161
<i>CsACTIN_R</i>	CATACCGATGAGAGATGGCTG	