

RISK ANALYSIS OF NOISE IN INDUSTRY MAKING CLOTHES

Análise do risco ruído em indústria de confecção de roupa

Daniela Dalapicula Barcelos⁽¹⁾, Soraya Gama de Ataíde⁽²⁾

ABSTRACT

Purpose: to get into an underexplored work environment and to identify if this environment might bring hearing losses to the workers. **Methods:** the environment chosen is a garment factory located in Colatina, Espírito Santo/Brazil. The two noisiest areas of the factory were selected to be analyzed, as well as six employees that have been working there for five years or so. The employees were asked to fill in a questionnaire on the hearing symptoms and then the audiometric tests taken for the last five years were compared. The analyzed areas produce noise from 83,5 to 97,8 dB. **Results:** from the audiometric tests we could conclude that 83% of the workers did not have any hearing loss, although 16% had occupational loss, but this was classified as stable. From the questionnaire we can conclude that 33% of the workers feel some kind of fatigue when exposed to high level sound for some time, 50% feel themselves stressed after their work shift, but none of them stated to hear any buzz nor insomnia, nor difficulty hearing other people. **Conclusion:** this work environment presents some risks to the hearing, once the employees are exposed to sound pressure levels during their work shifts, and the occupational hearing loss is evident although to a lesser extent – considering that the sample employees have been working there for at least five years. Therefore, it is plain to understand the importance of studying a good variety of work environments to identify where prevention programs, like the HLPP (Hearing Loss Prevention Program) should be applied.

KEYWORDS: Hearing Loss; Hearing Loss, Noise-Induced; Occupational Risks; Clothing

■ INTRODUCTION

Hearing is one of the noblest senses; its main function is linked to the acquisition and development of oral, essential language in interpersonal relations and the environment¹. Due to this importance, is that scholars increasingly seek ways to prevent the occurrence of hearing loss, especially those caused by noise.

“Noise threatens hearing, health, learning and behavior,” says Nancy Nadler, spokesman for the Center for Hearing and Communication (CHC), responsible for the institution of the International Day of Awareness about Noise, affecting humans in physical, psychological and social level². At high intensity and with continued exposure, noise can cause structural changes in the inner ear, which

determine the occurrence of Noise Induced Hearing Loss (NIHL), and is the most frequent cause of injury to the health of workers³, present in many⁴ lines of business, such as in industries, textiles, construction works, metallurgy, printing, sewing workshops, among other work environments⁵.

The garment industry in the country has emerged from the process of national industrialization with textile production. Brazilian Association of Clothing⁶ – With the evolution of the textile industry, the modernization of the manufacturing sector, which currently has twenty-one different articles of clothing and accessories segments, according to the classification of Abravest occurred. This is one of the sectors that soon developed in the country by requiring little technological level and small capital investment, in many regions, as occurred in the early industrialization of Brazil, this industry is the precursor of the industrialization process.

With efforts to increase the amount of production and maintain the products competitiveness in the market, companies invest and modernize faster and faster their machines. The great need of skilled

⁽¹⁾ University of Vila Velha – UVV, Vitória, ES, Brazil.

⁽²⁾ Specialization Course in Audiology CEFAC, Vitória, ES, Brazil.

Conflict of interest: non-existent

manpower in diverse industry, due to the variety of productive stages in the industry, such as: warehouse of fabrics and trims, creating, modeling, enfesto, cutting, sewing, crafts, laundry, ironing, finishing, labeling, packaging, final inspection and shipment⁶; attracts people who search for a paid service.

This niche market that has been growing steadily, it is underexplored in scientific research as working conditions and occupational health. There is little explanation about this environment in the speech, and he presents some risk, especially hearing the employee.

Because of this question, the aim of this study is to verify the presence of noise in garment manufacturing industry, since this risk is not widespread in research related to this work environment.

METHODS

The research, retrospective and documental study conducted in a company Tailoring located in Colatina-ES/Brazil.

The chosen industry has a total population of 368 employees. Through the Program of Environmental Risk Prevention supplied by them, noise measurements were analyzed in each sector, and thus raised the two sectors of higher risk noise, identified as Ironing (six employees) and area USED (with four employees), totaling ten workers. After applying the exclusion criteria, only six workers were selected to participate in the study (four are workers Ironing and two workers are Used).

Exclusion criteria were: employees with less than five years of service to the participating undertaking; workers in other sectors with less intensity noise or no risk and workers who declined to participate voluntarily in the study. With selected subjects, we analyzed the audiometric tests performed since the entrance exam to the current exam (2011) and can not therefore have fewer than five tests for each worker (at least one test for each year), available at medical records filed in the sector responsible for this function. Audiograms were analyzed for normality and hearing loss, which are subdivided into occupational and non-occupational. Occupational hearing loss found were analyzed for stability, worsening or triggering.

The selected officials responded in a single day and individually questionnaire approved by the ethics committee – Protocol 071/11 (Figure 1) of the closed type of information and auditory symptoms containing four questions, developed by the researcher based on descriptions of symptoms in cases NIHL in the literature.

With the questionnaires, they were collected and discharged workers. The responses obtained through the questionnaire and analysis of audiometry were transformed into percentages and presented in this work by means of graphs.

RESULTS

1. Description Field of research: Synthesis of the infrastructure of the selected company

The analyzed company is located in the town of Colac in the state of Espírito Santo, and acts for clothes in the jeans, shirts, knitwear and social segment. It employs 368 staff, with workday of 9 hours / day (7am entry ace out ace 17:15 pm with 1 hour 20 minute stop for lunch). The measurements of each sector were provided by the company resulting from PPRA conducted by a Technical Work Safety, the company contracted to carry out the reviews: Cutting Sector I – 80.8 to 89.7 dB (A) ; Sector Cut II – 80.8 to 82.2 dB (A) ; Sector Sewing I – 79.9 to 89.2 dB (A) , Sector Sewing II – 79.8 to 88.4 dB (a) ; Sewing Sector III – 80.5 to 84.4 dB (a) ; Sewing Sector IV – 79.8 to 84.1 dB (A) ; Sector Sewing Pilot – 81.1 to 86.4 dB (a) ; Salon Embroidery – 86.4 to 90.3 dB (a) ; Limpadeiras industry – from 80.1 to 82.7 dB (a) , Maintenance Sector I – 82.0 to 89.1 dB (a) ; industry Corrosion / sanding – 81.9 to 86.1 dB (a) ; Sector Boiler – 81.4 to 82.7 dB (A) ; Finishing Sector I – 76.8 to 79.7 dB (a) , Sector II Finish – 80.4 to 89.4 dB (a) ; industry Washing / Drying – 80.4 to 87.7 dB (a) ; Sector Montage / Collage – 79.7 to 82.3 dB (a) ; Sector area USED – from 88.4 to 98.2 dB (a) ; Sector Ironing – 83.5 to 93.5 dB (a) ; Sector General Administrative Services – 84.2 to 86.1 dB (a) . The other sectors as Snap , Modeling , Styling, Effluent Treatment , Storage of Chemicals , Warehouse , Shipping , Chair , Human Resources , Billing , Purchasing , Accounts payable / receivable , Sales , Production Planning and Control , Guaritas , Canteen , Interfacing and Accounting , show measurements below 80 dB (A) with no risk noise , however the PPRA other risks is described as gifts , heat, dust and solvents in their respective sectors , which were not listed because they are not subject of this study.

The company makes mandatory the use of individual protective shield sound, performing and training vouchers for their use, however, it can be observed that most workers do not use or even know the basic instructions on individual protectors. The company has an occupational physician who acts in certain on company time, and archiving in place of audiometric tests performed by specialized clinics, chosen for each contract.

QUESTIONNAIRE

1. Personal data:

1.1 Age : _____ years

1.2 You are exposed to noise outside the work environment? () YES () NO

1.2.1 Name the places: _____

2. Professional data:

2.1 How long have you worked at this company? _____

2.2 Have you worked in another company, you was exposed to noise? () YES () NO

2.2.1 If yes: How long? _____

2.2 Current Work Day:

2.2.1 Entry: _____ hours

2.2.2: Output _____ hours

2.2.3 Lunch: _____ hours

3. In your current job, you have contact with chemicals (solvents) () YES () NO

3.1 Make use of Personal Protective Equipment? () YES () NO _____

3.2 If yes: Is there training to use? () YES () NO

3.3 Do you think that effective training? () YES () NO

4. Have any of the symptoms below:

4.1 Tinnitus? () YES () NO

4.2 Difficulty in understanding what people say? () YES () NO

4.3 Insomnia? () YES () NO

4.4 Sense of hearing loss? () YES () NO

4.4.1 If yes: How long have you been noticing this change? _____

4.5 Stress or irritation at the end of the workday? () YES () NO

4.5.1 If yes: What do you attribute this stress ? _____

4.6 Do you feel uncomfortable when exposed to loud sounds? () YES () NO

Figure 1 – Questionnaire for company employees participating in the research*1.1. Description of the sectors analyzed:*

The sector identified with higher noise level was the Sector Area USED, where it performs the operation through the dyeing process that describes Used by putting the garment in place to be dyed, apply the dye through the air pistol compressed, remove clothes once dyed, deposit it in a pre-determined location, and other related services. The consequences outlined in the Prevention of Environmental Risks for this sector were: fatigue, irritability, headaches, decreased hearing, and so on. The equipment used in this industry as well as the noise generated by them, are listed in Table 1.

The second area examined was the Ironing industry, identified as the second sector presents

the greatest risk noise. The function of this sector is performing crossing clothing services, using the process of the steam press, noting the temperature and pressure of steam used and other related services. The consequences outlined in the Program for Prevention of Environmental Risks were: fatigue, irritability, headaches and decreased hearing. Table 2 lists the equipment used in this industry and their measurements.

The data found in measurements of the equipment used in sectors surveyed (Table 1 and 2) were analyzed and compared with the limits established by Norm 15 (NR15) and are available in Table 3.

Table 1 – Description of the equipment used in industry and their measurements Used

Machinery	Spot noise average / dB(A)	Spot noise maximum / dB(A)	Dosimetry real dose %
Used (cabin) Set Hood	96,5	97,8	353,08
Used (external) Hoods and Compressed Air - 1st job	96,1	98,2	
2nd job	93,2	94,1	
3rd job	88,4	92,7	

Table 2 – Description of the equipment used in industry and Ironing your measurements

Machinery	Spot noise average / dB(A)	Spot noise maximum / dB(A)	Dosimetry real dose %
Steam press I	88,1	91,2	143,40
Steam Presses II	88,9	92,7	
Press (turn pants / dummy)	87,8	90,5	
Ironing	84,5	86,3	

Table 3 – Noise Limits of Tolerance (NR-15)

Noise Level dB(A)	Maximum Permitted Daily Exposure
85	8 hours
86	7 hours
87	6 hours
88	5 hours
89	4 hours e 30 minutes
90	4 hours
91	3 hours e 30 minutes
92	3 hours
93	2 hours e 40 minutes
94	2 hours e 15 minutes
95	2 hours
96	1 hours e 45 minutes
98	1 hours e 15 minutes
100	1 hours
102	45 minutes
104	35 minutes
105	30 minutes
106	25 minutes
108	20 minutes
110	15 minutes
112	10 minutes
114	8 minutes
115	7 minutes

Of the six workers interviewed three complained of stress or irritation at the end of the workday, attributing the noise as the main factor. Two people reported discomfort when exposed to loud sounds and none of the respondents reported having tinnitus, insomnia or feeling of hearing loss / difficulty in understanding what people say (Figure 2).

The sample analyzed, five workers have a history of normality and has hearing loss. Hearing loss was found lodged in the chart above, specified as occupational hearing loss, this loss is stable (Figure 3).

The age range of respondents was 28-51 years old, and this item was not used as a criterion for participation in research. The analyzed group has 5-24 years of service to the company, in general, the same industry. Only 16% of subjects reported exposure to noise outside the work environment, as justifying another job, the waiter service at parties at least three times a week. Two subjects make use of ear and reported that training for the same is efficient, an employee reports using the shield, but thinks ineffective training, while three officials do not make use of ear training and complain.



Figure 2 – with responses obtained through the questionnaire




Figure 3 – Chart showing the results of analysis of Audiometries

DISCUSSION

During the 1980s, the noise was the main disease among metalworkers and affected more than 60% of category 7, and today is still rated as the most frequent agent of occupational exposure⁸⁻¹¹, being described as a major cause of sensorineural hearing loss in adults^{12,13}, usually bilateral, irreversible^{4,8,11,14-16}, affecting the inner ear and / or auditory nerve¹⁷, insidiously, slowly progressive changes and producing symptoms only after several years¹⁸.

This exposure can be very harmful, totally harmless or a measure of limits. The key issue is the duration of exposure, ie, the level of exposure determines the extent of the auditory effects of noise. All those working exposed to noise exceeding

85dBNPS and prolonged exposure should be monitored, which can be accomplished through pure tone audiometry^{8,19}.

The higher noise level of the equipment found in the analyzed sectors, generates noise of 98.2 dB (A) and considering the range of intensity to which employees are exposed (84.5 to 98.2 dB (A)), there that the exposure time, in a daily working hours exceeds the thresholds described in NR15 allowed.

Statistically, it is estimated that 15% of the exposed population constant noise of 90dB, eight hours per day, five days per week, 50 weeks a year develop hearing damage after ten years²⁰. This fact becomes even more important to analyze the various work sectors, often exceeding the exposure time allowed or do without proper protection.

Exposure during working hours to noise above permissible limits entitlement to punitive damages. Understanding is the Regional Labor Court of the 15th Region (Campinas, Brazil) ²¹, as in the case of the seamstress who won damages from garment factory that operated in the laundry industry ²².

When the noise stops being just a causative agent of discomfort in the workplace to become a potentially harmful agent to the worker's health, he will take a more specific approach within the field of occupational hygiene. Stands out in Annex 1 of NR 15 (Table 3), which describes the noise levels present in the work environment considered harmful to workers.

Auditory cause a major impact on people's lives, which can lead them to social isolation ²³, by difficulties in participating in conversations, making frustrating any family gathering or social event ^{24,25}.

Tinnitus is a symptom that often accompanies occupational hearing loss ²⁶, difficulties in being able to provide out of work contexts and negative influence on the quality of life of workers and the people who surround ^{27,28}. While none of the candidates analyzed have complained of tinnitus, this symptom should not be disregarded because to discard it, one should first examine the complaints of a larger population of workers exposed to noise and with more than five years of exposure.

Besides the frequent auditory symptoms – namely hearing loss, difficulty understanding speech, tinnitus ^{29,30} and intolerance to loud sounds – the worker carries PAIR also has complaints such as headache, dizziness and vertigo – because of the important changes that may occur in the labyrinthine structures, making the use of ear not only important to protect your hearing, but also for preservation of vestibular function ³¹ – irritability and digestive problems ⁴. Are also listed in the literature, symptoms such as hearing loss, ear fullness, earache, transient changes in blood pressure, stress ³², disturbances of vision, memory, sleep and mood directly related to the exposure time, with sound pressure levels (SPL) and individual susceptibility ^{33,34}, besides the reduction of speech perception in noisy environments, television, radio, film, theater, sound warning signs, music and environmental sounds ³⁵ and inability to relax ³⁶.

Stress was reported by participants in this study who reported feeling irritation / stress at the end of the workday, showing parity with the symptoms presented in other research that also still reported insomnia and lack of attention to consequences of noise exposure ¹⁵.

The volunteers showed another symptom, which is also described in the literature as a consequence of Noise Induced Hearing Loss: nuisance when

exposed to loud sounds. None of the respondents complained of hearing loss / difficulty understanding what people say. The age of the participants was not taken into account in this study, although it is relevant to correlate with complaints and investigate whether the absence of symptoms on exposure, is not beyond the short time working (most with only five years of service in this industry), the playfulness of the individual as well as the entire auditory system.

The diagnosis of noise-induced hearing loss of occupational origin depends on the audiogram and the proof of the existence of noise exposure in the workplace, considering the intensity and characteristics of the agent, and the manner and time of exposure, the general conditions health, age of the worker ³⁷, and the type of noise (continuous or intermittent) ¹¹.

The causes and consequences of NIHL are already quite widespread general knowledge, although many do not follow guidelines for preventing this loss. However, some work environments still present dearth of information, because the current concern is facing the risk of grade 3 or 4, where the presence of noise is notorious companies, not taking into account other sectors, which even if not noticeable feature risks to workers' health.

Although only one of the six analyzed workers present Noise Induced Hearing Loss, is adamant the literature the effects and consequences of hearing people exposed to noise. Therefore, considering that the site poses risks to hearing, for better measurement of health problems, new research may be carried out using as target workers with ten or more years of services with noise exposure. Even though this study has shown low levels of claims or losses, described and analyzed in the participating workers would not rule out preventive workers presented here with normal hearing, and that may present a possible trigger NIHL is only visible after a few more years of continuous exposure. The environment selected for analysis (two sectors of clothing firm) exposes workers to thresholds above the tolerance limit. Considering this fact, the company should seek to prevent and resolve problems relevant to excessive exposure to the worker of this site is not affected.

According to the regulations of the Ministry of Labour, companies must maintain a Program Environmental Risk Prevention (NR9), in which the various risks involved in the work must be identified and quantified, from that information, direct the actions of the the Medical Control of Occupational Health (NR7), which shall carry out reviews of health workers ⁴.

Regarding the noise risk, there is a specific program for its management, which consist of steps

as Assessment , Risk Management and Control , Management Audiometric ; Training and Educational Programs , among others . The objectives of this program are : To improve the quality of life of workers avoiding deafness and reducing extra- auditory effects caused by exposure to high noise levels and other risk agents for the hearing ; Early diagnosis for cases of occupational hearing loss , establishing effective measures , preserving the health of the worker; adapt the company to the requirements of the law (Regulatory Standards Nos. 7 and 9.) to be in compliance with occupational programs : Professional Profile for Social Security , the Medical Control of occupational health ; Prevention Program Environmental Risk ; Hearing Conservation Program ³⁸. The Hearing Conservation Program also considerably reduces the risk of accidents, since the risk of accidents is about twice as high among workers exposed to noise ³⁵.

The reflecting the implementation of the Hearing Conservation Program goes far beyond compliance. Develop educational programs is an act of intelligence, because it develops awareness, resulting in direct benefits for both the company and for the employee ³. For the viability of this program the involvement of professionals in health and safety, industrial management and human resources of the company and the health of workers is necessary ³⁹.

One way to make room for employees exposed themselves participate in programs, is through the Internal Accident Prevention Commission, which was created with the intention of staff to use as a tool for organization and better conditions of work and health as the risk analysis must rely on the experience, knowledge and participation of those undertaking the daily work and suffer its effects ⁷. Therefore knowing the difficulties of individuals towards noise can be able to deploy an educational process within a Program Hearing Loss Prevention ²³.

The Hearing Conservation Program is a coordinated set of actions that aims to prevent or stabilize occupational hearing loss. These, featuring a continuous and dynamic process of implementation of routines in companies seeking to reduce or eliminate risks to hearing in the present processes, described as one of the goals actions: improving the quality of life of workers avoiding deafness and reducing extra-auditory effects caused by exposure to high levels of sound pressure and other risk

agents for the hearing, which probably decreases discrimination against them with varying degrees of hearing loss, that when unemployed, face the difficulty of achieving new employment ⁴⁰, thus initiating a process of social and psychological devaluation generated by unemployment.

Audiometric evaluation of workers is still the most effective way to determine the success of a Hearing Conservation Program. Compared with the annual audiometric results can detect changes or reinforcements in the proposed interventions, finding areas in which the hearing (collective and / or individual) protection has flaws ³⁸. The speech therapist is a trained professional that making use of the knowledge that agents have about risk, the affected organ and audiological tests it performs, can develop or enhance important preventers' practices in promoting hearing health of the population¹³.

One of the most talked about topics and worked by professional speech therapy and training is the use of Personal Protective Equipment that must be carefully considered, because most workers do not like to use them and end up creating some resistance . Most people exposed to noisy environments, do not know the risks that noise brings health , either for lack of information or as a result of the unpreparedness of the company or in bad faith when it intends to gain something from the company. In this study specifically , we note that of the six subjects selected for study , only three are used ear plugs , one of which relates find ineffective training , which leads us to think that the shield may be being used incorrectly and it disable its effect . However , even if this data is analyzed without questions still must be taken of the fact that three employees – half of the sample – not a daily use of the shield.

The hearing conservation programs in industry are an important way to prevent hearing loss. Hearing loss caused by noise is totally incurable today, but it is also absolutely avoidable ^{26,40}.

■ CONCLUSION

This workplace noise risk has since recorded the sound pressure levels at which workers are exposed during the workday, and the presence of occupational hearing loss, even on a smaller scale – awarded to the minimum of five years of services and the limited number of participants in research.

RESUMO

Objetivo: verificar a presença do ruído em indústria de confecção de roupas, visto que este risco é pouco difundido nas pesquisas relacionadas a este ambiente de trabalho. **Métodos:** a empresa escolhida foi uma fábrica de confecção de roupa localizada em Colatina no Espírito Santo/Brasil. Foram analisados os dois setores com maior risco ruído na empresa, após a exclusão dos funcionários com menos de cinco anos de trabalho neste local, totalizamos uma amostra de seis funcionários. Os trabalhadores selecionados responderam questionário sobre informações e sintomas auditivos e posteriormente, foram analisadas as audiometrias realizadas nos últimos cinco anos. **Resultados:** por meio de medições realizadas nos setores analisados, constatou-se produção de ruído de 83,5 a 97,8 dB(A). Após observar o histórico das audiometrias, obteve-se 83% dos trabalhadores, destes setores, sem perda auditiva, 16% com perda ocupacional, sendo classificada como estável. Os dados levantados no questionário revelaram que 33% dos trabalhadores sentem-se irritados quando expostos a sons elevados, 50% sentem-se estressados após a jornada de trabalho, e nenhum trabalhador relatou queixa de zumbido, insônia ou dificuldade em entender as pessoas. **Conclusão:** este ambiente de trabalho apresenta risco ruído, uma vez constatados os níveis de pressão sonora no qual os trabalhadores são expostos durante a jornada de trabalho, bem como a presença de perda auditiva ocupacional, mesmo em menor escala – atribuído ao tempo mínimo de cinco anos de serviços prestados utilizado na pesquisa.

DESCRIPTORIOS: Perda Auditiva; Perda Auditiva Provocada por Ruído; Riscos Ocupacionais; Vestuário

■ REFERENCES

1. Botelho CT, Paz APML, Gonçalves AM, Frota S. Estudo comparativo de exames audiométricos de metalúrgicos expostos a ruído e ruído associado a produtos químicos. *Rev. Bras. Otorrinolaringol.* [periódico na internet]. 2009 [acesso em 2011 Fev 04];75(1):[51-7]. Disponível em: <http://www.scielo.com.br>
2. Farias VHV, Buriti AKL, Rosa MRD. Ocorrência de perda auditiva induzida pelo ruído em carpinteiros. *Rev CEFAC* [periódico na internet]. 2012 Mai [acesso em 2012 Jul 20];14(3):413-22. Disponível em: <http://dx.doi.org/10.1590/S1516-18462011005000119>
3. Bramatti L, Morata TC, Marques JM. Ações educativas com enfoque positivo em programa de conservação auditiva e sua avaliação. *Rev CEFAC* [periódico na internet]. 2008 [acesso em 2012 Ago 02];10(3):398-408. Disponível em: <http://dx.doi.org/10.1590/S1516-18462008000300016>
4. Brasil. Ministério da Saúde. Perda auditiva induzida por ruído (Pair). Secretaria de Atenção à Saúde. Departamento de Ações Programáticas Estratégicas. Brasília: Editora do Ministério da Saúde. 2006.
5. Lima CG. Dossiê Técnico: Prevenção de perdas auditivas ocupacionais causadas por ruídos. Fundação Centro Tecnológico de Minas Gerais – CETEC. 2007 [acesso em 2011 Abri 12]. Disponível em: <http://www.sbrt.ibict.br>
6. Assis SF, Souza JF, Nascimento LC. Diagnóstico dos resíduos sólidos gerados pelas indústrias de confecção de Colatina/ES. Colatina: Centro Federal de Educação Tecnológica do Espírito Santo. 2008 Out 23 [acesso em 18 Out 2012]. Disponível em: http://pse.ifes.edu.br/prppg/pesquisa/jornadas/jornada_2007_2008/pibiti/engenharias/pibiti%20-%20eng%20-%20jonio%20-%20sarina%20-%20146%20147.pdf
7. Mazzeu FJC, Demarco DJ, Kalil L. Segurança e saúde no trabalho. São Paulo: Unitrabalho-Fundação Interuniversitária de Estudos e Pesquisas sobre o Trabalho. Brasília – DF: Ministério da Educação. SECAD-Secretaria de Educação Continuada, Alfabetização e Diversidade, (Coleção Cadernos de EJA); 2007 [acesso em 2012 Jan 12]; [aproximadamente 64 p.]. Disponível em: http://portal.mec.gov.br/secad/arquivos/pdf/10_cd_al.pdf
8. Santos CCS, Juchem LS, Rossi AG. Processamento auditivo de militares expostos a ruído ocupacional. *Rev CEFAC* [periódico na internet]. 2008 Mar [acesso em 2012 Ago 02];10(1):92-103. Disponível em: <http://dx.doi.org/10.1590/S1516-18462008000100013>
9. Dias A, Cordeiro R. Fração atribuível de acidentes do trabalho decorrentes da exposição ao ruído ocupacional em cidade do Sudeste do Brasil. *Cad Saúde Pública* [periódico na internet]. 2007 Jul [acesso em 2012 Jul 23];23(7):1649-55.

Disponível em: <http://dx.doi.org/10.1590/S0102-311X2007000700016>

10. Holanda WTG, Lima MLC, Figueiroa JN. Adaptação transcultural de um instrumento de avaliação do handicap auditivo para portadores de perda auditiva induzida pelo ruído ocupacional. *Ciênc Saúde Coletiva* [periódico na internet]. 2011 [acesso em 2012 Ago 02];16(1):755-67. Disponível em: <http://dx.doi.org/10.1590/S1413-81232011000700006>

11. Cézar MRV. Atuação do fonoaudiólogo na prevenção da perda auditiva induzida por ruído. CEFAC Recife [monografia na internet]; 2008 [acesso em 2012 Jul 15]; [aproximadamente 42 páginas]. Especialização em Audiologia. Disponível em: <http://www.cefac.br/library/teses/6c632bfa6a1536f760aad180eb36afae.pdf>

12. Prudente MF, Costa EA. Exposição ao ruído ocupacional: alterações no exame de emissões otoacústicas. *Rev. Bras. Otorrinolaringol.* [periódico na internet]. 2006 Jun [acesso em 2012 Jul 20];72(3):362-6. Disponível em: <http://dx.doi.org/10.1590/S0034-72992006000300011>

13. Morata TC, Zucki F. Promoção da saúde auditiva diante de riscos ambientais: uma reflexão da atuação e da produção científica na área. In: Morata TC, Zucki F. Caminhos para a saúde auditiva: ambiental – ocupacional. São Paulo: Plexus Editora; 2005. p.11-27.

14. Teles RM, Medeiros MPH. Perfil audiométrico de trabalhadores do distrito industrial de Maracanau – CE. *Rev. Soc. Bras. Fonoaudiol.* [periódico na internet]. 2007 [acesso em 2012 Jul 20];12(3):233-9. Disponível em: <http://www.scielo.br/pdf/rsbf/v12n3/a11v12n3.pdf>

15. Faria CAR, Suzuki FA. Avaliação dos limiares auditivos com e sem equipamento de proteção individual. *Rev. Bras. Otorrinolaringol.* [periódico na internet]. 2008 [acesso em 2012 Jul 20];74(3):417-22. Disponível em: <http://dx.doi.org/10.1590/S0034-72992008000300017>

16. Riveiro VFM. Ruído e suas consequências para o adoecimento: uma revisão integrativa de literatura. Universidade Federal do Rio Grande do Sul [monografia na internet]; 2010 [acesso em 2012 Ago 02]. [aproximadamente 45 páginas]. Disponível em: <http://www.lume.ufrgs.br/bitstream/handle/10183/28226/000770079.pdf?sequence=1>

17. Fernandes LCBC, Gil D, Maria SLS, Azevedo MF. Potencial evocado auditivo de tronco encefálico por via óssea em indivíduos com perda auditiva sensorioneural. *Rev CEFAC* [periódico na internet]. 2012 Mar [acesso em 2012 Ago 02], pp.0-0. Disponível em: <http://dx.doi.org/10.1590/S1516-18462012005000018>

18. Pereira RT, Minette LJ, Magalhães BL, Abrantes BA, Silva EP, Souza AP. Avaliação ergonômica dos fatores ambientais de uma indústria de confecção na zona da mata mineira. UFV: III SAEPRO; 2007 [acesso em 2012 Jan 13]. Disponível em: <http://www.saepru.ufv.br/Image/artigos/SA04.pdf>

19. Rocha EB, Azevedo MF, Filho JAX. Estudo da audição de crianças de gestantes expostas ao ruído ocupacional: avaliação por emissões otoacústicas – produto de distorção. *Rev. Bras. Otorrinolaringol.* [periódico na internet]. 2007 Jun [acesso em 2012 Jul 20];73(3):359-69. Disponível em: <http://dx.doi.org/10.1590/S0034-72992007000300011>

20. Rodrigues MAG, Dezan AA, Marchiori LLM. Eficácia da escolha do protetor auditivo pequeno, médio e grande em programa de conservação auditiva. *Rev CEFAC* [periódico na internet]. 2006 Dez [acesso em 2012 Jul 20];8(4):543-7. Disponível em: <http://dx.doi.org/10.1590/S1516-18462006000400016>

21. Costa JL. Perda auditiva por excesso de ruído no trabalho gera dano moral. *Revista Consultor Jurídico.* [base de dados na internet]. 2005 Jun 17 [acesso em 2012 Jul 20]. [7 páginas]. Disponível em: http://www.conjur.com.br/2005-jun-17/perda_auditiva_ruido_trabalho_gera_dano_moral

22. Revista Consultor Jurídico. [base de dados na internet]. 2011 Mar 1 [acesso em 2012 Jan 13]. Costureira deve ser indenizada por perda auditiva; [1 página.]. c2011. Disponível em: <http://www.conjur.com.br/2011-mar-01/costureira-indenizada-perda-auditiva-decide-trt-rs>

23. Bramatti L, Morata TC, Marques JM, Martini UG. Versão e adaptação para o português brasileiro do questionário: crenças e atitudes sobre prevenção de perda auditiva. *Rev CEFAC* [periódico na internet]. 2012 Fev [acesso em 2012 Ago 02]; 14(1):65-78. Disponível em: <http://dx.doi.org/10.1590/S1516-18462011005000082>

24. Signorini TA. Deficiência auditiva do idoso e sua implicação na comunicação. In: Gonçalves CGO, Mota PHM, Marques JM. Ruído e idade: análise da influência na audição em indivíduos com 50 – 70 anos. *Pró-Fono R. Atual. Cient.* [periódico na Internet]. 2009 Mar [acesso 2011 Jul 02];21(1):57-62. Disponível em: http://www.scielo.br/scielo.php?script=sci_arttext&pid=SOQO456872009000100010&lng=pt

25. Mondelli MFCG, Rocha AB. Correlação entre os achados audiológicos e incômodo com zumbido. *Arquivos Int. Otorrinolaringol. (Impr.)* [periódico na internet]. 2011 Jun [acesso em 2012 Ago 02];15(2):172-80. Disponível em: <http://dx.doi.org/10.1590/S1809-48722011000200009>

26. Pfeiffer M, Rocha RLO, Oliveira FR, Frota S. Intercorrência audiológica em músicas após um

show de rock. Rev CEFAC [periódico na Internet]. 2007 [acesso em 2012 Jul 23];9(3):423-9. Disponível em: <http://redalyc.uaemex.mx/src/inicio/ArtPdfRed.jsp?iCve=169313371017>

27. Dias A, Cordeiro R. Interação entre grau de perda auditiva e o incômodo com zumbidos em trabalhadores com história de exposição ao ruído. Rev. Bras. Otorrinolaringol. [periódico na Internet]. 2008 Dez [acesso em 2011 Jul 01];74(6): 876-83. Disponível em: http://www.scielo.br/scielo.php?script=sci_arttext&pid=S00347299200800060010&lng=pt

28. Ganime JF, Silva LA, Robazzi MLCC, Sauzo SV, Faleiro SA. El ruido como riesgo laboral: una revisión de la literatura. Revista eletrônica de enfermagem [periódico na internet]. 2010 [acesso em 2012 Jul 17];19 [aproximadamente 10 p]. Disponível em: http://scielo.isciii.es/pdf/eg/n19/pt_revision1.pdf

29. National Institute for Occupational Safety and Health. Preventing occupational hearing loss: a practical guide. In: Bramatti L, Morata TC, Marques JM. Ações educativas com enfoque positivo em programa de conservação auditiva e sua avaliação. Rev CEFAC [periódico na internet]. 2008 [acesso em 2012 Jul 20];10(3): 398-408. Disponível em: <http://dx.doi.org/10.1590/S1516-18462008000300016>

30. Steinmetz LG, Zeigelboim BS, Lacerda AB, Morata TC, Marques JM. Características do zumbido em trabalhadores expostos a ruído. Rev. Bras. Otorrinolaringol. [periódico na internet]. 2009 Fev [acesso em 2012 Jul 20];75(1):7-14. Disponível em: <http://dx.doi.org/10.1590/S0034-72992009000100002>

31. Teixeira CS, Körbes D, Rossi AG. Ruído e equilíbrio: aplicação da posturografia dinâmica em indústria gráfica. Rev CEFAC [periódico na internet]. 2011 Fev [acesso em 2012 Ago 02];13(1):92-101. Disponível em: <http://dx.doi.org/10.1590/S1516-18462010005000016>

32. Boger ME, Branco AB, Ottoni AC. A influência do espectro de ruído na prevalência de Perda Auditiva Induzida por Ruído em trabalhadores. Braz. j. otorhinolaryngol. (Impr.) [periódico na internet]. 2009 Jun [acesso em 2012 Ago 02];75(3):328-34. Disponível em: <http://dx.doi.org/10.1590/S1808-86942009000300003>

Received on: April 20, 2012

Accepted on: December 10, 2012

Mailing address:

Daniela Dalapicula Barcelos

Rua Dulce Brito Espíndula, 21, Ed. Praia de Capuba, apto 601

Jardim Camburi – Vitória – ES

CEP: 29090-340

E-mail: danidalapicula@yahoo.com.br

33. Ribeiro BL, Machado AA, Back DC. Perda auditiva induzida por níveis de pressão sonora elevados na indústria. CEFAC [monografia na internet]. 2009 [acesso em 2011 Jul 2]; [aproximadamente 12 p.]. Disponível em: <http://www.cefac.br/library/artigos/4994382af80a4b40a3fdd9dea3e405e9.pdf>

34. Ribeiro AMD, Câmara VM. Perda auditiva neurossensorial por exposição continuada a níveis elevados de pressão sonora em trabalhadores de manutenção de aeronaves de asas rotativas. Cad. Saúde Pública [periódico na Internet]. 2006 [acesso em 2012 Jul 20];22(6):1217-24. Disponível em: <http://www.scielo.br/scielo>

35. Lopes AC, Almeida AC, Mello ADP, Otubo KA, Lauris JRP, Santos CC, et al. Characterization of the Audiologic Thresholds in Workers of Funeral Urns. Int Arch Otorhinolaryngol. [periódico na internet]. 2009 [acesso em 2012 Ago 02];13(3):244-51. Disponível em: <http://www.arquivosdeorl.org.br/conteudo/pdfForl/13-03-02.pdf>

36. Weber SR, Périco E. Zumbido no trabalhador exposto ao ruído. Rev. Soc. Bras. Fonoaudiol. [periódico na internet]; 2011 Dez [acesso em 2012 Ago 02]. 16(4): 459-65. Disponível em: <http://dx.doi.org/10.1590/S1516-80342011000400016>

37. Guida HL, Morini RG, Cardoso ACV. Avaliação audiológica em trabalhadores expostos a ruído e praguicida. Braz. j. otorhinolaryngol. (Impr.) [periódico na internet]. 2010 Ago [acesso em 2012 Ago 02];76(4):423-7. Disponível em: <http://dx.doi.org/10.1590/S1808-86942010000400003>

38. Bernardi e Saldanha. 2003. In: Lima CG. Dossiê Técnico: Prevenção de perdas auditivas ocupacionais causadas por ruídos. Fundação Centro Tecnológico de Minas Gerais – CETEC. 2007 [acesso em 2011 Abri 12]. Disponível em: <http://www.sbrt.ibict.br>

39. Ministério da Previdência e Assistência Social. OS/INSS nº 608, 05/08/1998 – Aprova Norma Técnica sobre Perda Auditiva Neurossensorial por Exposição Continuada a Níveis Elevados de Pressão Sonora de Origem Ocupacional. In: Souza MHP, Mercuri AAS, Azevedo MF, Bernardi APA. Análise de emissões otoacústicas produtos de distorção em trabalhadores de uma indústria metalúrgica na cidade de Birigui. Rev CEFAC. [periódico na internet]. 2010 [acesso em 2012 Jul 20]. Disponível em: <http://www.cefac.br/library/artigos/f18097d018dd48c1f9743c47e7835710.pdf>

40. Ogido R, Costa EA, Machado HC. Prevalência de sintomas auditivos e vestibulares em trabalhadores expostos a ruído ocupacional. Rev. Saúde Pública [periódico na internet]. 2009 [acesso em 2012 Jul 20];43(2):377-80. Disponível em: <http://www.scielo.br/pdf/rsp/v43n2/7239.pdf>