



Initial Identification and Sensitivity to Antimicrobial Agents of *Salmonella* sp. Isolated from Poultry Products in the State of Ceara, Brazil

Author(s)

Oliveira WF¹
Cardoso WM^{2*}
Salles RPR¹
Romão JM³
Teixeira RSC¹
Câmara SR¹
Siqueira AA³
Marques LCL⁴

- ¹ Postgraduate students in Veterinary Sciences of Ceara State University
- ² Professor Doctor of Veterinary Medicine School of Ceara State University
- ³ Undergraduate students of Veterinary Medicine School of Ceara State University
- ⁴ Veterinarian of the Laboratory of Animal Health of the Brazilian Ministry of Agriculture

Mail Address

William Maciel Cardoso
Av. Rogaciano Leite, 200 - Aptº 1303
Bl. Tulipe, Bairro Salinas
60810-000. Fortaleza, CE, Brazil
Phone/Fax: 55 (085) 241 1307 or 299 2748
or 9989 4742

E-mail: william.maciel@uol.com.br
josueromao@yahoo.com.br

Keywords

Salmonella sp, identification, sensitivity, poultry products.

ABSTRACT

The objective of this research was to isolate and to verify the sensitivity to antimicrobial agents of strains of *Salmonella* sp. isolated from poultry products in the state of Ceara, Brazil. A total number of 114 samples was collected from 63 broiler carcasses derived from two processing plants and two supermarkets, and 51 excreta samples were collected in broiler farms located in the state of Ceara, which used three live production stages. Each excreta sample consisted of a fresh excreta pool from 100 birds. Samples were submitted to microbiological analyses, and the isolated *Salmonella* strains were tested for antimicrobial sensitivity. No *Salmonella* was isolated from excreta samples, while broiler carcass samples showed a high contamination rate of 11.8%. Three serotypes were identified: *Salmonella enterica* serovar Enteritidis, 50%; *Salmonella enterica* serovar Panama 33%, and *Salmonella enterica* serovar Newport, 17%. As to the susceptibility tests to antimicrobial agents, 100% of the isolated *Salmonella* strains showed resistance to Ampicillin and Tetracycline, and sensitivity to Gentamycin, Netilmycin, Carbenicillin, Chloramphenicol.

INTRODUCTION

Brazil is one of the largest broiler producers in the world. In order to maintain this annual productivity, rational management emphasizing the quality of poultry products is necessary. For this purpose, the government has implemented health programs aiming at controlling diseases that cause economic and health problems. General prophylactic procedures and biological security practices applied in every stage of broiler production (broiler grand-parent farms, broiler parent farms, commercial broiler chick farms, and hatcheries) decrease, but do not prevent the presence of bacteria. Among the pathogenic microorganisms in industrial poultry production, the genus *Salmonella* is of paramount importance.

In terms of public health, there are *Salmonella* strains that may cause paratyphoid infection, as well as *Salmonella enteritidis*, *Salmonella typhimurium*, *Salmonella infantis*, and *Salmonella agona*, which are important sources of food-borne illnesses. These bacteria are not related to specific diseases, and are capable of infecting indistinctly several animal species, including humans (Lax *et al.*, 1995). The importance of *Salmonella* in the public health is very significant; for instance, it accounted for nearly 84% of food-borne human illnesses in Scotland from 1980 to 1989 (Oboegbulem *et al.*, 1993), and 81% in Italy from 1991 to 1994 (Scuderi *et al.*, 1996). According to the Centers for Disease Control and Prevention in United States, *Salmonella* is responsible for 1.34 million of cases of disease, 16.430 hospitalizations, and 582 deaths each year (Mead *et al.*, 1999). The total annual cost resulting from food-



borne *Salmonella* infections in United States is estimated in about US\$3.5 billion (Pathogen, 1995).

The epidemiology of *Salmonella* infections in birds is very complex (Hinton, 1988), and it is difficult to determine how a flock was infected, or how the dissemination occurred in the flock. Hong'ombe *et al.* (1999) identified *S. enteritidis* in broiler carcasses ready for market in several different situations. Many authors identified poultry products as sources of infection of *Salmonella* that causes enteritis in humans (Dhillon *et al.*, 2001). Approximately 10% of salmonellosis cases are caused by poultry meat, and in the United States, there are between 15 and 20 cases of salmonellosis for each 100,000 people (Bryan & Doyle, 1995). According to Skov *et al.* (2002), the number of diagnosed cases of salmonellosis in humans in Denmark increased during the last decade, and the serotypes *S. enteritidis* and *S. typhimurium* were the most commonly isolated strains. Major outbreaks of human salmonellosis were caused by the consumption of incorrectly manipulated foods in restaurants and institutional kitchens. Undercooked food, slow freezing, and keeping foods for many hours under no refrigeration are considered as contributing factors for the emergence of this disease in humans (Costa, 1996).

In spite of the programs of Brazilian Agriculture Ministry to control poultry pathogens, there are reports of outbreaks of salmonellosis, and later detection of the pathogen in broiler carcasses and foods containing eggs in Brazil. The main factors of this contamination are improper handling of these products in some companies, the epidemiologic complexity of *Salmonella*, and deficient inspection during production, processing, and trade of animal products. However, studies on the contamination of broilers with *Salmonella* are practically null in our region.

Due to the need to constantly monitor of the health quality of birds for human consumption, this study aimed at isolating and identifying *Salmonella* serotypes present during live production, and trading of chicken meat, as well as the behavior of *Salmonella* spp. strains relative to common antimicrobial drugs.

MATERIALS AND METHODS

Collection of excreta samples during live production

A total number of 63 excreta samples were collected from 21 broiler flocks in three management stages: starter (1 day of age), grower (20 days of age), and finisher (45 days of age), from broiler companies

located in the state of Ceara. Each sample consisted of a fresh excreta pool from 100 birds, randomly collected in commercial broiler houses. The collected samples were placed in sterile plastic bags, and were submitted to the laboratory, following the criteria recommended by the National Poultry Health Program – PNSA (Brazil/MAARA, 2002).

Collection of samples from broilers carcasses at retail

Broiler carcasses were collected from sales points and processing plants, and were divided into the following categories: fresh carcasses (14 samples), collected in two processing plants with no federal sanitary inspection (7 carcasses in each plant); refrigerated carcasses (18 samples), ten in one supermarket and eight in another; and frozen carcasses (19 samples), ten samples collected in one supermarket, and nine from another.

Microbiological analyses

The excreta samples (live production phase) and the carcasses were randomly collected during typical management, slaughter, and sales days.

Feces samples weighing 25 g were diluted in 225 mL of buffered peptone water at 0.1%, and carcass samples were submitted to the "carcass wash" method, through the addition of 300 mL of buffered peptone water at 0.1%, as described by Cox *et al.* (1978), and then placed in Erlenmeyer flasks.

Bacterial culture followed the procedures established by National Poultry Health Program – PNSA (Brasil/MAARA, 2002) with some modifications. The protocol started with pre-enrichment: the solutions resulting from the process "carcass wash" were incubated at 37 °C for 24 hours. Aliquots from pre-enrichment were inoculated into selective enrichment liquid media at a ratio of 1/100 in Rappaport-Vassiliadis broth and at 1/10 in Selenite-Cysteine broth. A loopful of each broth was streaked on plates of Brilliant Green agar, MacConkey agar, and *Salmonella-Shigella* agar. The temperature and the period of incubation were standardized at 37 °C for 24 hours, respectively, for both feces and carcass samples. Two or three suspected colonies of *Salmonella* from each plate were collected for presumptive identification by biochemistry tests. The media utilized for presumptive identification were: Triple Sugar Iron slant agar (TSI); Lysine-Iron slant agar (LIA); Sulfur Indol Motility agar (SIM), and the Citrate test. Tubes were incubated at 37 °C for 24 hours. Colonies with biochemistry profile of *Salmonella*



were submitted to serologic tests by the use of polyvalent serum against O and H *Salmonella* antigens. The colonies that agglutinated during the period of one to two minutes were considered as positive for *Salmonella*, and were preserved in Nutrient agar. Isolates were submitted to Adolfo Lutz Institute in São Paulo, Brazil, for complete identification and serotyping.

Antimicrobial sensitivity test

The behavior of all isolated of *Salmonella* sp. strains was checked as to the action of antimicrobial drugs. Strains were submitted to sensitivity tests according to the BAUER-KIRBY method (Bauer *et al.*, 1966). Each strain was inoculated in BHI (brain and heart infusion) broth, and after 24 hours of incubation at 37 °C they were streaked using sterile swabs on Mueller-Hinton agar plates. Plates were kept in environmental temperature for a period of five minutes, and then diffusion disks with antimicrobial drugs were distributed on the plates and incubated for 24 hours at 37 °C. The antimicrobial drugs were: Ampicillin (10 mg); Amoxicillin (10 mg); Amikacin (30 mg); Norfloxacin (10 mg); Tetracycline (30 mg); Chloramphenicol (30 mg); Netilmycin (30 mg); Gentamycin (10 mg); Sulfonamide (300 mg), and Carbenicillin (100 mg). Results were interpreted by measuring inhibition zones with the use of a millimeter scale rule. The results were presented as resistant or sensitive according to *National Committee for Clinical Laboratory Standards* (NCCLS, 2000).

Statistical Analysis


Companies and categories of carcasses were compared using the Chi-square (χ^2) non-parametric test for qualitative analysis at a probability of 5% ($p < 0.05$).

RESULTS

The presence of *Salmonella* spp. was investigated in 63 excreta samples of 21 flocks in three different live production stages. No *Salmonella* isolated from any of the samples, thereby characterizing the analyzed flocks as free from *Salmonella* infection.

Salmonella was isolated in all categories of analyzed broiler carcasses, as shown in Table 1.

In fresh carcass samples, *Salmonella panama* (14.3%) was isolated. Refrigerated carcasses showed a high rate of contamination with *Salmonella enteritidis* and *Salmonella newport* (16.7%). Frozen carcasses showed the lowest contamination rate, with the isolation of *Salmonella enteritidis* (5.26%). *Salmonella* contamination among carcass samples varied; however, these differences were not statistically significant ($p < 0.05\%$).



Graph 1 - Percentage of *Salmonella* strains isolated in broiler carcasses.

Among all the isolated strains, *Salmonella enteritidis* presented the highest rate of isolation (50%), followed by *Salmonella panama* (33.3%) and *Salmonella newport* (16.6%).

The results of antimicrobial sensibility tests are presented in Table 2. The behavior of the isolated strains from broiler carcasses relative to the action of antibiotics showed that all the isolated strains were resistant to Ampicillin and Tetracyclin. Four isolates (*Salmonella panama* -1 and *Salmonella enteritidis* -3) were resistant to Amoxicillin. Three isolates (*Salmonella panama* -1 and *Salmonella enteritidis* -2)

Table 1 – Isolation and characterization of *Salmonella* in broiler carcasses.

Carcasses	Isolation (%)	Samples (n ¹)	Isolated strains (n ²)
Fresh	2 (14.3)	14	<i>S. panama</i> (2)
Refrigerated	3 (16.7)	18	<i>S. newport</i> (1); <i>S. enteritidis</i> (2)
Frozen	1 (5.26)	19	<i>S. enteritidis</i> (1)
Total	6 (11.8)	51	<i>S. enteritidis</i> (3); <i>S. panama</i> (2); <i>S. newport</i> (1);

n - number of samples. Not statistically different ($p < 0.05\%$).



Table 2 - Antimicrobial sensitivity test of *Salmonella* strains isolated from broiler carcasses.

<i>Salmonella</i>	N. of samples	Resistance to antibiotics (µg)									
		AMP(10)	TET(30)	AMO(10)	AMI(30)	GEN(10)	NET(30)	SUF(300)	NOR(10)	CLO(30)	CAR(100)
Panama	2	2	2	1	-	-	-	1	1	-	-
Newport	1	1	1	-	-	-	-	-	-	-	-
Enteritidis	3	3	3	3	1	-	-	2	-	-	-
Total (%)	100,0	100,0	100,0	66,7	16,7	-	-	50,0	16,7	-	-

Legend: AMP – Ampicillin, TET – Tetracycline, AMO – Amoxicillin, AMI – Amikacin, GEN – Gentamycin, NET – Netilmycin, SUF – Sulfonamide, NOR – Norfloxacin, CLO – Chloramphenicol, CAR – Carbenicillin.

were resistant to Sulfonamide, while only one isolate of *Salmonella enteritidis* presented resistance to Amikacin, and one isolate of *Salmonella panama* presented resistance to Norfloxacin.

DISCUSSION

Salmonella is one of the most important sources of food-borne illness in humans, mainly due to the complexity of Salmonellosis epidemiology (Zancan *et al.* 2000). Broiler meat is a high risk product, according to Berchieri (2000), and production systems and processing facilitate the presence of *Salmonella* in the final product.

There are several methods to verify the contamination of *Salmonella* in samples (Read *et al.* 1994). The bacterial analysis of excreta is a sensitive test (Aho, 1992), and present better results as compared to the methods using cloacal swabs and antibody research in avian serum. Other advantages are that it does not cause stress, and is more representative of bird contamination (Giessen *et al.* 1991). The absence of *Salmonella* in the 63 excreta samples from broiler flocks produced in the state of Ceará implies that management and hygiene are good in these farms. These results are in agreement with Moreira (2002), who did not find *Salmonella* in day old broiler chicks reared by poultry companies in the same region. However, Gama (2001) found *Salmonella enteritidis* contamination in commercial layer chicks, and Zancan *et al.* (2000) verified 44.45% of *Salmonella* contamination in the transport boxes of commercial layer chicks.

On the other hand, all carcass categories (fresh, refrigerated, and frozen) presented some kind of contamination, as shown in Table 1.

Salmonella panama was isolated in 14.3% of the fresh carcasses. In refrigerated carcasses, there was a high level of contamination (16.7%) with *Salmonella enteritidis* and *Salmonella newport*. The lower rate of contamination occurred in frozen carcasses (5.26%) with the isolation of *Salmonella enteritidis*. The total

rate of *Salmonella* contamination of broiler carcasses was 11.8%, which is considered high, The Codex Alimentarius established 0% contamination in 25 g of analyzed food, including poultry meat and eggs (Santos *et al.*, 2000). This percentage is lower than that found by Santos *et al.* (2000), with 32% of contamination; by Machado & Bernardo (1990), with 57% of contamination; by Plummer *et al.* (1995), with 23%; and Arvanitidou *et al.* (1998), with 69%. Our results are consistent with the percentage verified by Sharma (1992), with 9.21% of contamination of broiler carcasses. However, our findings are higher than those reported by Verde *et al.* (2003), who found 3.6% of *Salmonella* contamination in broiler carcasses, and of the USDA, which reported that the national pathogen-reduction program reduced the incidence of *Salmonella* contamination in broiler carcasses to less than 10% in 2000 (Food, 2000).

The presence of *Salmonella panama* (14.3%) in fresh broiler carcasses and the high rate of *Salmonella enteritidis* and *Salmonella newport* in refrigerated broiler carcasses indicates that processing must be better controlled. Navarro (1995) asserts that the number of food-borne disease increased considerably in Central and South America, mainly due to *Salmonella enteritidis* (Santos *et al.*, 2000). The importance of the workers in the processing line was studied by several authors, as the lack of hygiene favors the dissemination of contamination of broiler surfaces. Pether & Gilbert (1971) analyzed the level of dissemination of *Salmonella anatum* by the extremity of fingers from previously infected human volunteers. They verified that the isolation of *Salmonella* from the finger extremity of these people was directly proportional to the inoculated bacterial population; the results were 100% positive for 10⁶ inoculated bacteria, and 30% for the interval of 10³ to 10⁴ inoculated bacteria.

The isolated strains in refrigerated carcasses were *Salmonella newport* in one case, and *Salmonella enteritidis* in the two other cases. This represents a significant contamination risk, as many authors report that *Salmonella enteritidis*, among the genus



Salmonella, is the main source of food-borne diseases (Banatvala *et al.*, 1999; Ferris *et al.*, 1999; Plummer *et al.*, 1995; Rodrigues *et al.*, 1990). According to Costa *et al.* (1996), refrigerated carcasses present high levels of contamination, whereas there are few reports of *Salmonella* in frozen carcasses. Santos *et al.* (2003) verified that, out of 272 isolates of *Salmonella*, 111 were found in frozen carcasses, 126 in food and biological human material involved with cases of food-borne diseases, and 35 in different poultry materials, with the phage type 4 as the most prevalent bacterium. Roberts (1982) observed 80% contamination in frozen carcasses. This result and our experiment indicate the possibility that broiler carcasses, including frozen carcasses, to disseminate *Salmonella* to humans; however, Foster & Mead (1976) assert that the carcass freezing decreases or impairs the survival of the *Enterobacteriaceae* family.

Among isolated and characterized strains (Graph 1), *Salmonella newport* presented the lowest isolation rate in broiler carcasses. According to Uyttendaele *et al.* (1998), *Salmonella newport* was the most prevalent *Salmonella* isolated in turkeys, and this serotype is not commonly isolated from chickens (Bokanyi *et al.*, 1990; Ferris *et al.*, 1999; Poppe *et al.*, 1998; Read *et al.*, 1994). *Salmonella enteritidis* was the most prevalent isolate in the present experiment (50%). This result is consistent with the findings of Santos *et al.* (2003), and Rodrigues *et al.* (1990). Kinde *et al.* (1997) found seven isolates of *Salmonella enteritidis* out of 683 isolated strains in waste water from processing plants located in different districts of California. Other contamination sources were described by other authors, such Cortinez *et al.* (1995), who isolated *Salmonella newport* and *Salmonella panama* from river water samples in Argentina, and Hofer *et al.* (2000), who isolated these serotypes in horse meat in the Northeast Region of Brazil.

Salmonella strains resistant to antibiotics are commonly found, and this may be due to the comprehensive use of antibiotics included in feeds as growth promoters. The results of the antimicrobial sensitivity tests carried out in the present experiment are alarming, as all isolates showed 100% resistance to Ampicillin and Tetracycline. These results agree with the findings of Cortinez *et al.* (1995), who found strains sensitive to Gentamicin and Chloramphenicol, and resistant to Tetracycline. Berchieri *et al.* (1983) verified 77% resistance to Tetracycline of *Salmonella* in poultry feed, and Antunes *et al.* (2003) found 36% of *Salmonella* strains resistant to Tetracycline in broiler

carcasses. Bokanyi *et al.* (1990), Lee *et al.* (1993), and Nascimento *et al.* (1997) showed similar results, with 100% of sensitivity to Chloramphenicol in broiler carcasses. Lee *et al.* (1993) and Santos *et al.* (2000) found 100% of resistance to Ampicillin of *Salmonella* in broiler carcasses. Our results showed that 66.7% of isolates were resistant to Amoxicillin, while Antunes *et al.* (2003) found 19%, 19%, and 3% of resistance to Amoxicillin, Carbenicillin and Chloramphenicol, respectively. The low isolation rates of *Salmonella* strains resistant to sulfonamide is consistent with Arvanitidou *et al.* (1998), who found 19.35% of resistance to Ampicillin, and few strains were resistant to sulfafurazole in chicken carcasses.

According to our results, broilers evaluated during live production in farms located in the state of Ceará presented good microbiological quality in terms of *Salmonella* infection; however, there was a marked decrease of the microbiological quality of poultry products analyzed in this experiment. As to antimicrobial sensitivity tests, the results were alarming, because 100% of the isolates were resistant to Ampicillin and Tetracycline.

REFERENCES

- Aho M. Problems of *Salmonella* sampling. International Journal of Food Microbiology 1992; 15:225–235.
- Antunes P, Cristina R, Sousa JC, Peixe L, Pestana N. Incidence of *Salmonella* from poultry products and their susceptibility to antimicrobial agents. International Journal of Food Microbiology 2003; 82:97–103.
- Arvanitidou M, Tsakris A, Sofianou D, Katsouyannopoulos V. Antimicrobial resistance and R-factor transfer of salmonellae isolated from chicken carcasses in Greek hospitals. International Journal of Food Microbiology 1998; 40:197-201.
- Banatvala N, Cramp A, Jones IR, Feldman RA. Salmonellosis in North Thames (East), UK: Associated risk factors. Epidemiology and Infection 1999;122:201-207.
- Bauer AW, Kirby WMM, Sherris JC, Turck M. Antibiotic susceptibility testing by a standardized single disk method. American Journal of Clinical Pathology 1966; 45:493–496.
- Berchieri Jr A. Contaminação por *Salmonella* em farinhas de origem animal utilizadas no preparo de ração. São Paulo (SP): USP; 1983, [Master Dissertation]. São Paulo (SP): Instituição de Ciências Biomédicas, USP; 1983.
- Berchieri Jr A. Salmoneloses aviárias. In: Berchieri JrA, Macari M editor. Doenças das aves. Campinas: FACTA; 2000. p.185–196.
- Bokanyi RP, Stephens JF, Foster DN. Isolation and characterization



- of *Salmonella* from broiler carcasses or parts. Poultry Science 1990; 69:592–598.
- Bokanyi RP Jr, Stephens JF, Foster DN. Isolation and characterization of *Salmonella* from broiler carcasses or parts. Poultry Science 1990; 69:592–598.
- Brasil. Portaria SDA nº 126, de 03 de setembro de 1995. Aprova as normas de credenciamento e monitoramento de Laboratórios de Diagnóstico das *Salmonellas* aviárias (*S. enteritidis*, *S. gallinarum*, *S. pullorum* e *S. typhimurium*). Diário Oficial da União, Brasília, n.212, p.17694-17698, de 6 de novembro, seção I.
- Bryan FL, Doyle MP. Health risks and consequences of *Salmonella* and *Campylobacter jejuni* in raw poultry. Journal of Food Protection 1995; 58:326–344.
- Cortinez IJM, Velasquez LDC, Escudero ME, Caffer MI, Cobo MF, Guzman AMS. *Salmonella* serotypes from surface waters in San Luis, Argentina. Revista Microbiologia São Paulo 1995; 26:180–185.
- Costa FN. Sorotipos de *Salmonella* em carcaças e cortes de frango obtidos na indústria e no comércio e comportamento das cepas isoladas frente à ação de antimicrobianos. [Master dissertation]. Jaboticabal (SP): Faculdade de Ciências Agrárias e Veterinárias; UNESP; 1996. 82 p.
- Cox NA, Mercuri AJ, Tanner DA, Carson MO, Thomson JE, Bailey JS. Effectiveness of sampling methods for *Salmonella* detection on processed broilers. Journal Food Protection, Des Moines 1978; 41:341–343.
- Dhillon AS, Shivaprasad HL, Roy P, Alisantosa B, Schaberg D, Johnson S. Pathogenicity of Environmental Origin *Salmonella* in Specific Pathogen – Free Chicks. Poultry Science 2001; 80:1323–1328.
- Ferris KE, Fisher SD, Flugrad BR, Timm JM. *Salmonella* serotypes from animals and related sources reported during July 1998. In: Proceedings of the 103rd Annual Meeting of the United States Animal Health Association; 1999; San Diego, CA. p.488–507.
- Food Safety and Inspection Service – FSIS. Interim progress report on *Salmonella* testing of raw meat and poultry products. Washington; 2000.
- Forster RD, Mead GC. Effect of temperature and added polyphosphate on the survival of salmonellae in poultry meat during cold storage. Journal of Applied Bacteriology 1976; 41:504–510.
- Gama NMSQ. *Salmonella* spp em aves de postura comercial [Master dissertation]. Jaboticabal (SP): Faculdade de Ciências Agrárias e Veterinárias, UNESP; 2001.
- Giessen AW, Peters R, Berkers PA, Jansen WH, Notermans SH. *Salmonella* contamination of poultry flocks in the Netherlands. Veterinary Quarterly 1991; 13:41–46.
- Hinton M. *Salmonella* infection in Chicks Following the Consumption of Artificially Contaminated Feed. Epidemiology and Infection 1988; 100:247–256.
- Hofer E, Zamora MRN, Lopes AE, Moura AMC, Araújo HL, Leite MDD, Filho SJS. Sorovares de *Salmonella* em carne de equídeos abatidos no nordeste do Brasil. Pesquisa Veterinária Brasileira 2000; 20:80–84.
- Hong'ombe BM, Sharma RN, Skjerve E, Tuchili LM. Occurrence of *Salmonella enteritidis* in Pooled Table Eggs and Market Ready Chicken Carcasses in Zambia. Avian Diseases 1999; 43:597–599.
- Kinde A, Adelson A, Ardans A, Little EH, Willoughby D, Berchtold D, Read DH, Breitmeuer R, Kerr D, Tarbell R, Hughes E. Prevalence of *Salmonella* in Municipal Sewage Treatment Plant Effluents in Southern California. Avian Disease 1997; 41:392–398.
- Lax AJ, Barrow PA, Jones PW, Wallis TS. Current Perspectives in Salmonellosis. British Veterinary Journal 1995; 151:351–377.
- Lee LA, Threatt VL, Puhr ND, Levine P, Ferris K, Tauxe RV. Antimicrobial-resistant *Salmonella* spp. isolated from healthy broiler chickens after slaughter. Journal of the American Veterinary Medical Association 1993; 202:752–755.
- Machado J, Bernardo F. Prevalence of *Salmonella* in chicken carcasses in Portugal. Journal of Applied Bacteriology 1990; 69:477–480.
- Mead PS, Slutsker L, Diet V, McCaig LF, Bresee JS, Shapiro C, Griffin PM, Tauxe RV. Food-related illness and death in the United States. Emerging Infectious Diseases 1999; 5:607–625.
- Moreira APO. Pesquisa de *Salmonella* sp. em frangos de corte de um dia de idade da Região Metropolitana de Fortaleza [Master dissertation]. Fortaleza (CE): UECE; 2002.
- Nascimento VP, Cardoso MO, Ribeiro AR, Santos LR, Silva AB, Pontes AP, Oliveira SD. Prevalência e perfis de resistência de *Salmonella* isoladas de carcaças de frango frente a antimicrobianos e desinfetantes selecionados. In: XIX Congresso Brasileiro de Microbiologia; 1997; Rio de Janeiro, RJ, Brasil. p.291.
- National Committee for Clinical Laboratory Standards – NCCLS. MIC testing supplemental tables M100 – S10 (M7) 2002. Weyne (Pa); 2002.
- Navarro MP. Infecção por *Salmonella enteritidis* em reprodutoras pesadas na América Latina. In: Conferência APINCO de Ciência e Tecnologia Avícolas; 1995; Curitiba, PR, Brasil. p.7–16.
- Oboegbulem SI, Collier PW, Sharp JCM, Reilly WJ. Epidemiological aspects of outbreaks of food-borne salmonellosis in Scotland between 1980 and 1989. Revue scientifique et technique 1993; 12:957–967.
- PATHOGEN reduction; hazard analysis and critical control point (HACCP) systems. Federal Register, 1995; 60(143):6774–6889.
- Pether JVS, Gilbert RJ. The survivals of salmonellas on fingertips and transfer of the organisms to food. The Journal of Hygiene 1971; 69:673–681.
- Plummer RAS, Blissett SJ, Dodd CER. *Salmonella* contamination of retail chicken products sold in UK. Journal of Food Protection 1995; 58:843–846.



Poppe C, Duncan CL, Mazzocco A. Salmonella contamination of hatching and table eggs: a comparison. Canadian Journal of Veterinary Research 1998; 62:191–198.

Read SC, Irwin RJ, Poppe C, Harris J. A comparison of two methods for isolation of Salmonella from poultry litter samples. Poultry Science. 1994; 73:1617–1621.

Roberts T. Factors contributing to outbreaks of food poisoning in England and Wales 1970-1979. The Journal of Hygiene 1982; 89:491–498.

Rodrigues DC, Taxue RV, Rowe B. International increase in *Salmonella enteritidis*: A new pandemic? Epidemiology and Infection 1990; 105:21–27.

Santos DMS, Berchieri Jr A, Fernandes SA, Tavechio AT, Amaral LA. *Salmonella* em Carcaças de Frango Congeladas. Pesquisa Veterinária Brasileira 2000; 20:39-42.

Santos LR, Nascimento VP, Oliveira SD, Rodrigues DP, Reis EMF, Ribeiro AR, Fernandes SA. Phage types of *Salmonella enteritidis* isolated from clinical end food samples, and from broiler carcasses in southern Brazil. Revista do Instituto de Medicina Tropical de São Paulo 2003; 45:1-4.

Scuderi G, Fantasia M, Filetici E, Anastásio MP. Foodborne outbreaks caused by salmonella in Italy, 1991-1994. Epidemiology and Infection 1996; 116:257–265.

Sharma VD. *Salmonella* contamination of foods of animal origin. In: *Salmonella* and Salmonellosis Symposium; 1992; Ploufragan, FR. p137–138.

Skov MN, Feld NC, Carstensen B, Madsen M. The Serologic Response to *Salmonella* typhimurium in Experimentally Infected Chickens, Followed by an Indirect Lipopolysaccharide Enzyme – Linked Immunosorbent Assay and Bacteriologic Examinations Through a One – Year Period. Avian Diseases 2002; 46:265–273.

Uyttendaele MR, Debevere JM, Lips RM, Neyts KD. Prevalence of *Salmonella* in poultry carcasses and their products in Belgium. International Journal of Food Microbiology 1998; 40:1–8.

Verde JCL, Bezerra FSBC, Soares MIM, Brito ECO, Santos CD. Monitoramento de *Salmonella* em carcaça de frango coletados em diversos estabelecimentos comerciais da cidade de Fortaleza. In: XIII Encontro Nacional de Analistas de Alimentos; 2003 jun. 22-25; Rio de Janeiro, RJ, Brasil, 2003. p.164.

Zancan FT, Berchieri Jr A, Fernandes SA, Gama NMSQ. *Salmonella* investigation in transport boxes of day-old birds. Brazilian Journal of Microbiology 2000; 31:230–232.