

*Letter to Editor*

## Prevalence of glucose-6-phosphate dehydrogenase deficiency in blood donors of Mossoró, Rio Grande do Norte

Ulysses Madureira Maia<sup>1</sup>

Danielle Cristiny de Azevedo Batista<sup>1</sup>

Wogelsanger Oliveira Pereira<sup>2</sup>

Thales Allyrio Araújo de Medeiros Fernandes<sup>3</sup>

Universidade do Estado do Rio Grande do Norte – Mossoró (RN), Brasil.

<sup>1</sup>Biological Sciences Course, Universidade do Estado do Rio Grande do Norte, Mossoró, Brazil

<sup>2</sup>Cytology and Bimolecular Organization Service, Biomedical Sciences Department, Universidade do Estado do Rio Grande do Norte, Mossoró, Brazil

<sup>3</sup>Human Genetics Service, Biomedical Sciences Department, Universidade do Estado do Rio Grande do Norte, Mossoró, Brazil

*Conflict of interest:* none

Dear Sir,

Glucose-6-phosphate dehydrogenase (G6PD) deficiency, affecting about 330 million people worldwide, is the most common hereditary enzymopathy.<sup>(1)</sup> Although most individuals are asymptomatic, this condition can lead to the development of neonatal jaundice, chronic nonspherocytic hemolytic anemia and acute hemolytic anemia induced by certain types of drugs, infections or the consumption of broad beans.<sup>(2)</sup>

Moreover, blood transfusions from G6PD deficient donors are less effective as the red blood cells of these individuals are deficient in antioxidant defense mechanisms; they are more vulnerable to extra- and intra-vascular hemolysis due to oxidative stress and can have excessive storage degradation with decreased survival.<sup>(3)</sup>

The overall prevalence of G6PD deficiency is geographically correlated to endemic malaria, hence higher in sub-Saharan Africa, the Middle East, Southeast Asia, Mediterranean Europe and some areas of Latin America.<sup>(1)</sup> Studies on the Brazilian population have reported a prevalence of G6PD deficiency in about 1% to 10% of the population with the highest rates being found among men of African descent.<sup>(4)</sup> In the particular case of Rio Grande do Norte, previous studies have shown a prevalence of 3.5% in patients seen at a referral hospital in the city of Natal.<sup>(5)</sup> However, no study on G6PD deficiency in a provincial population of Rio Grande Norte had been performed previously. Hence, this cross-sectional

study was conducted during the period from August 2006 to August 2008, to determine the prevalence of G6PD deficiency in blood donors from the city of Mossoró, the largest provincial city of Rio Grande do Norte.

A total of 714 individuals (aged 18-62 years, mean 30 years) who voluntarily went to the local blood center to donate blood were analyzed; 576 (80.7%) were men and 138 (19.3%) women, 343 (48.0%) said they were white and 371 (52.0%) stated they were mulatto, brown, yellow or black (grouped as nonwhites).

Donors were informed about the nature and importance of the work and were asked about their willingness to participate in the study. Those who agreed were asked to sign informed consent forms and a standard questionnaire designed to obtain demographic data was applied. Subsequently, 5-mL samples of peripheral venous blood were collected in tubes containing 50 µL 10% EDTA for laboratory tests.

The G6PD activity was initially measured by the methaemoglobin reduction test<sup>(6)</sup> and with confirmation of positive results by the methaemoglobin reduction test modified for semiquantitative evaluation.<sup>(7)</sup> Individuals were considered deficient if the enzyme activity was below 70%. Statistical analyses of the results were carried out employing the Pearson  $\chi^2$  test using the PEPI program. Statistical significance was set for p-values <0.05. The protocol of this study was approved by the Research Ethics committee of the Hospital Universitário Onofre Lopes (CAAE # 0395.0.000.294-07).

Twenty-seven (3.8%) cases of G6PD deficiency were identified, a percentage similar to those described for the population of Natal (3.5%) and Bahia (3.2%). Among the G6PD deficient individuals, 21 (77.8%) were male and 6 (22.2%) were female. There were no statistically significant differences in the prevalence in enzyme deficiency between genders ( $\chi^2 = 0.151$ , p = 0.698) (Table 01), even though it has characteristically X-linked inheritance. This finding may be attributed to the fact that the deficiency does not cause significant harm and thus reproduction between affected individuals and carriers of G6PD deficiency is relatively common.

Table 1. Cases of G6PD deficiency in blood donors from Mossoró, Brazil in respect to gender

Gender	Total samples	G6PD activity			
		Normal		Deficient	
		n	(%)	n	(%)
Male	576	555	(96.4)	21	(3.6)
Female	138	132	(95.7)	6	(4.3)
Total	714	687	(96.2)	27	(3.8)

(Pearson  $\chi^2$  test):  $\chi^2 = 0.151$ , p = 0.698

Thirteen (48.1%) of the G6PD deficient individuals stated they were white and 14 (51.9%) stated they were non-whites; no statistical differences were observed between the ethnical groups ( $\chi^2 = 0.145$ ; p = 0.703) (Table 2).

These relatively high prevalence rates of G6PD deficiency, together with the reduced effectiveness of blood transfusions from

Table 2. Cases of G6PD deficiency in blood donors from Mossoró, Brazil in respect to ethnical background

Gender	Total samples	G6PD activity			
		Normal		Deficient	
		n	(%)	n	(%)
Whites	343	331	(96.5)	12	(3.5)
Non-whites	371	356	(96.0)	15	(4.0)
Total	714	687	(96.2)	27	(3.8)

(Pearson  $\chi^2$  test):  $\chi^2 = 0.145$ , p = 0.703

individuals with this genetic condition, reinforces the need of G6PD deficiency screening of donors, especially when blood is transfused in neonates.

### Abstract

*Glucose-6-phosphate dehydrogenase (G6PD) deficiency is the most common human enzymopathy. It affects as many as 330 million individuals worldwide. This deficiency may determine neonatal jaundice, chronic nonspherocytic hemolytic anemia and acute hemolytic anemia induced by drugs, infections and broad bean ingestion. The efficacy of blood transfusion is decreased when the donor is G6PD deficient. In this study, we aimed at determining the prevalence of G6PD deficiency in blood donors of Mossoro, Brazil. Samples of 714 blood donors (576 men and 138 women; 343 white and 371 non-white) with ages ranging from 18 to 62 years and that accepted to participate in the study were analyzed. All participants answered a standard questionnaire. G6PD activity was analyzed by the methemoglobin reduction test with deficiency being confirmed by the semiquantitative test. The overall prevalence of G6PD deficiency in blood donors was 3.8%, similar to the rate described for others regions of Brazil. There was no significant statistical difference in the frequency of G6PD deficiency between men and women, nor between white and non-white blood donors. This relatively high frequency of G6PD deficiency highlights a need to screen blood donors for this condition.*

**Keywords:** Glucosephosphate dehydrogenase deficiency; Blood donors; Anemia

### References

- Nkhowa AT, Poole C, Vannappagari V, Hall AS, Beutler E. The global prevalence of glucose-6-phosphate dehydrogenase deficiency: a systematic review and meta-analysis. *Blood Cells Mol Dis.* 2009; 42(3):267-78.
- Minucci A, Giardina B, Zuppi C, Capoluongo E. Glucose-6-phosphate dehydrogenase laboratory assay: how, when and why. *IUBMB Life.* 2009;61(1):27-34.
- Samanta S, Kumar P, Kishore SS, Garewal G, Narang A. Donor blood glucose-6-phosphate dehydrogenase deficiency reduces the efficacy of exchange transfusion in neonatal hyperbilirubinemia. *Pediatrics.* 2009;123(1):96-100.
- Silva RT, Iglessias MAC, Medeiros ID, Bezerra IM, Medeiros TMD. Deficiências de glicose-6-fosfato desidrogenase em adultos. Newslab [Internet]. 2005 [citado 2009 Jul 27];79:96-102. Disponível em: [http://www.newslab.com.br/ed\\_anteriores/79/art03/art03.pdf](http://www.newslab.com.br/ed_anteriores/79/art03/art03.pdf)
- Mauricio CRF, Maia RD, Queiroz SMV, Araújo MGM, Miranda RGC, Medeiros TMD. Deficiência de glicose-6-fosfato desidrogenase: dados de prevalência em pacientes atendidos no Hospital Universitário Onofre Lopes, Natal - RN. RBAC [Internet]. 2006[citado 2009 Maio 24];38(1):57-9. Disponível em: [http://www.sbac.org.br/pt/pdfs/rbac/rbac\\_38\\_01/rbac3801\\_14.pdf](http://www.sbac.org.br/pt/pdfs/rbac/rbac_38_01/rbac3801_14.pdf)
- Brewer GJ, Tarlov AR, Alving AS. Methaemoglobin reduction test : a new simple in vitro test for identifying primaquine-sensitivity. *Bul World Health Organ.* 1960;22:633-40.
- Severo LG, Nogueira DM, Hoxter G. Determinação da atividade da G-6-PD em eritrócitos humanos. *Laes & Haes.* 1985;1:22-30.

Submitted: 5/28/2010

Accepted: 6/28/2010

**Correspondence:** Thales Allyrio Araújo de Medeiros Fernandes  
Faculdade de Ciências da Saúde – UERN  
Rua Miguel Antônio da Silva Neto, S/N, Aeroporto  
59607-360 – Mossoró (RN), Brasil  
Phone: 55 84 3315-2248  
E-mail: [thalesallyrio@uern.br](mailto:thalesallyrio@uern.br)