

The following table replaces the Table incorrectly published in the Original Article “Reticulocytes indices in β thalassemia trait individuals” published in Volume 33 Issue n° 5 of the *Revista Brasileira de Hematologia e Hemoterapia*.

Table 1- Hematological indices in β thalassemic individuals and controls

Index	β -thalassemia trait (median \pm standard deviation)	Controls (median \pm standard deviation)	p-value
Age (years)	44.3 \pm 19.4	43.0 \pm 16.5	0.717
White blood cells ($\times 10^9/L$)	7.0 \pm 2.1	6,0 \pm 1.4	0.018
Red blood cells ($\times 10^{12}/L$)	5.53 \pm 0.53	4.66 \pm 0.44	< 0.001
Hemoglobin (g/dL)	11.5 \pm 1.2	14.1 \pm 1.4	< 0.001
Hematocrit (%)	37.0 \pm 3.8	42.4 \pm 3.7	< 0.001
mean cell volume (fL)	66.8 \pm 4.4	91.2 \pm 4.3	< 0.001
mean cell hemoglobin (pg)	20.9 \pm 1.3	30.2 \pm 1.5	< 0.001
mean cell hemoglobin concentration (g/L)	312 \pm 8	331 \pm 8	< 0.001
Red cell distribution width (%)	15.9 \pm 1.0	13.0 \pm 0.6	< 0.001
Reticulocytes (%)	1.58 \pm 0.59	1.05 \pm 0.31	< 0.001
Reticulocytes ($\times 10^9/L$)	87.0 \pm 31.2	49.2 \pm 14.4	< 0.001
Immature reticulocyte fraction	0.1347 \pm 0.062	0.0995 \pm 0.039	0.007
Low fluorescence reticulocytes (%)	89.9 \pm 17.5	90.0 \pm 3.9	0.97
Medium fluorescence reticulocytes (%)	12.3 \pm 0.5	9.2 \pm 3.5	0.007
High fluorescence reticulocytes (%)	1.2 \pm 1.1	0.7 \pm 0.7	0.037