



Revista Brasileira de Hematologia e Hemoterapia Brazilian Journal of Hematology and Hemotherapy

www.rbhh.org



Original article

Pilot randomized controlled trial to evaluate the effect of aquatic and land physical therapy on musculoskeletal dysfunction of sickle cell disease patients



Camila Tatiana Zanoni, Fábio Galvão, Alberto Cliquet Junior, Sara Teresinha Olalla Saad*

Universidade Estadual de Campinas (UNICAMP), Campinas, SP, Brazil

ARTICLE INFO

Article history:

Received 11 June 2014

Accepted 25 July 2014

Available online 21 November 2014

Keywords:

Anemia, sickle cell

Hydrotherapy

Physical therapy modalities

ABSTRACT

Objective: To compare the effect of aquatic and land-based physiotherapy in reducing musculoskeletal hip and lower back pain and increasing overall physical capabilities of sickle cell disease patients.

Methods: Informed written consent was obtained from all volunteers who were submitted to evaluations using different functional scales: Lequesne's Algofunctional Questionnaire and Oswestry Disability Index, trunk and hip range of motion, goniometry, trunk and hip muscle strength assessment using load cell, and surface electromyography of the iliocostalis, long dorsal (longissimus), gluteus maximus, gluteus medius and tensor fasciae latae muscles. Ten patients were randomized into two groups: aquatic physiotherapy with a mean age of 42 years (range: 25–67) and conventional physiotherapy with a mean age of 49 years (range: 43–59). Both groups were submitted to a twelve-week program of two sessions weekly.

Results: After the intervention, significant improvements were observed regarding the Lequesne index (p -value = 0.0217), Oswestry Disability Index (p -value = 0.0112), range of motion of trunk extension (p -value = 0.0320), trunk flexion muscle strength (p -value = 0.0459), hip extension and abduction muscle strength (p -value = 0.0062 and p -value = 0.0257, respectively). Range of motion of trunk and hip flexion, extension, adduction and abduction, trunk extensor muscle strength and all surface electromyography variables showed no significant statistical difference.

Conclusion: Physical therapy is efficient to treat musculoskeletal dysfunctions in sickle cell disease patients, irrespective of the technique; however, aquatic therapy showed a trend toward improvement in muscle strength. Further studies with a larger patient sample and longer periods of therapy are necessary to confirm these results.

© 2014 Associação Brasileira de Hematologia, Hemoterapia e Terapia Celular. Published by Elsevier Editora Ltda. All rights reserved.

* Corresponding author at: Hemocentro Unicamp, Instituto Nacional de Ciência e Tecnologia do Sangue, Rua Carlos Chagas, 480, Cidade Universitária Zeferino Vaz, 13083-878 Campinas, SP, Brazil.

E-mail address: sara@unicamp.br (S.T.O. Saad).

<http://dx.doi.org/10.1016/j.bjhh.2014.11.010>

1516-8484/© 2014 Associação Brasileira de Hematologia, Hemoterapia e Terapia Celular. Published by Elsevier Editora Ltda. All rights reserved.

Introduction

Sickle cell disease (SCD) is a genetic disorder that results in the sickling of red blood cells, triggering vaso-occlusion episodes which lead to pain and organ damage. This inherited disorder is caused by a point mutation in the beta-globin gene. The mutant form of hemoglobin in SCD (Hb S) is capable of polymerization and complex molecular and structural changes within the red cell. Occurring in homozygotes (Hb SS) and in compound heterozygotes, such as those carrying Hb S and beta-thalassemia (Hb SB⁺ or Hb SB⁰) and Hb S and hemoglobin C (Hb SC), hemolytic anemia and vaso-occlusion crises are the main complications of SCD. The illness tends to gradually develop toward multisystem organ failure.¹ Bone involvement, frequently causing painful vaso-occlusive crises, is the most common clinical manifestation. Furthermore, bone involvement is a source of chronic, progressive disability, with long-term effects upon bone mass density, growth, and bone damage such as avascular necrosis and osteomyelitis. Osteopenia and osteoporosis are often asymptomatic; however, pain, fractures, deformities, and vertebral collapse may occur and require chronic analgesia, mechanical support, and surgical interventions.^{2,3}

Chronic and progressive damage such as, for example, avascular necrosis of the femoral head, the leading cause of hip deformity in these patients, commonly results in gait disturbances, pain, and activity and functional limitations in adult patients. Lower back pain is one of the main complaints among SCD patients and occurs due to the flattening and widening of the vertebral bodies with biconcave depressions of the endplates, probably caused by infarction of the central portion of the vertebral body.^{4,5} There are few studies in the literature on the role of physiotherapy as a resource to prevent and treat locomotor system disorders in SCD individuals.

According to recent studies, the life expectancy of SCD patients' has improved dramatically over the last century.⁶ However this longer life span has, as an unfortunate consequence, the development of progressive organ damage which includes osteoarticular lesions.⁷

Chronic pain is considered a serious public health problem which negatively affects the quality of life of individuals. Therefore, a multi-action therapeutic plan, specifically physiotherapy, could help decrease pain, and improve mobility and the rehabilitation of osteoarticular disorders, positively impacting on the quality of life.⁸

Despite this fact, there are few studies in the literature on the role of physiotherapy as a resource to prevent and treat locomotor system disorders in SCD patients. One study⁹ compared the efficacy of physiotherapy alone with physiotherapy associated with surgical femur decompression in SCD patients with osteonecrosis of the femoral head. The results showed no significant difference between these two approaches, suggesting that physical therapy alone appeared to be as effective as surgical decompression to improve hip function, thus deferring the need for surgery.

Within the existing physiotherapy resources, aquatic physiotherapy used in rehabilitation has demonstrated positive effects against pain, in regaining physical function and in improving quality of life in adults with musculoskeletal

conditions.¹⁰ Movements performed in the water are facilitated by the elimination of the effects of gravity, resulting in increased muscle strength (MS) and flexibility. The benefits of water are mainly explained by the physiological effects of immersion and by the hydrodynamic principles of exercise, such as buoyancy, in this environment thereby enabling functional exercises with a reduced gravitational load. Furthermore, the immersion in thermo-neutral water (34 °C) in combination with the effects of hydrostatic pressure reduces the perception of pain. The physical properties and heated water play an important role in improving and maintaining the range of joint motions, reducing muscular tension and promoting relaxation, as well as preparing the muscle for stretching. The buoyancy induces muscle relaxation and the decrease in impact enables increased mobility and flexibility.^{11,12}

This study aimed to evaluate the efficacy of aquatic and land-based physical therapy in decreasing hip and lower back musculoskeletal pain and increasing overall physical well-being in SCD patients.

Methods

Adult SCD patients who regularly attended (at least three times a year during the previous three years) the Outpatient Clinic of the Hemocentro of the Universidade Estadual de Campinas (UNICAMP) with chronic hip and lumbar spine pain, and who had not participated in a physical therapy program during the previous 12 months, were invited to participate in this study. Patients with acute episodes, absence of over three physical therapy sessions without justification, or any dermatological issue which would prevent them from entering a therapeutic pool, were excluded from the study. The National Ethics Board approved this study, and all patients provided written informed consent.

Study design

Initially, the volunteers were evaluated according to functional scales, including the Lequesne's Algofunctional Questionnaire and Oswestry Disability Index (ODI), range of motion (RoM) measurements of trunk flexion and extension, hip adduction and abduction, assessment of MS of the trunk flexors and extensors, and the flexors, extensors, adductors and abductors of the hip through load cell and surface electromyography (SEMG) of the iliocostalis, long dorsal (longissimus), gluteus maximus, gluteus medius and tensor fasciae latae muscles. Volunteers were then randomized by an investigator not involved in data collection, using the blind allocation method of sequentially numbered, opaque sealed envelopes,^{13,14} into two different program groups: aquatic physiotherapy (AP) and conventional or land physiotherapy (CP). A total of 24 sessions over a twelve-week period (two sessions per week) were administered. Patients were assessed after the intervention comparing the results before and after the sessions according to data obtained for the dominant side of each patient.^{15,16}

Questionnaires

The Lequesne's Algofunctional Questionnaire was developed for patients with osteoarthritis and evaluates symptoms and functional capacity of the hip and knee. This index is composed of 11 questions that evaluate pain, discomfort and function. The ODI is a self-administered questionnaire used to measure the degree of lumbar spine disability, and contains topics concerning intensity of pain and physical activity.¹⁷ The ODI has been used in scientific research to evaluate patients with nonspecific or specific low-back pain after surgical procedures, medication and rehabilitation.

Range of motion

The RoM was evaluated by a single examiner using a conventional 360° free shaft goniometer. The following movements were assessed: trunk flexion and extension and flexion, extension, adduction and abduction of the hip joint, according to the standardization of the goniometry manual of Marques.¹⁸

Muscle strength

MS was analyzed by maximal voluntary isometric contraction (MVIC) using a load cell (MIOTEC®, Porto Alegre, Brazil). The load cell was connected to a Miotool 400® apparatus (MIOTEC®) using a SDS1000® sensor connected via a USB cable to a notebook. During movements, the force generated by traction on the load cell was transmitted to the Miograph® software which produces a plot of MS in kilograms-force (kgf). Volunteers were submitted to isometric MS tests of the trunk flexors and extensors and hip flexors, extensors, adductors and abductors.

Surface electromyography

Myoelectric signals of the gluteus maximus, gluteus medius, tensor fascia lata, long dorsal (longissimus) and iliocostalis muscles were sampled at 2000 Hz in single differential mode

from each muscle through a four channel electromyography system (MIOTEC®, Porto Alegre, Brazil) using disposable Ag/AgCl circular bipolar electrodes (3M®). The 10 mm diameter electrodes with adhesive conducting gel were positioned on the skin overlying the muscles at an inter-electrode distance of 20 mm. Abrasion of the skin was achieved at the fixation sites with gauze soaked in alcohol to reduce impedance and the electrodes were then fixed at the muscular belly, distant from the motor point, and fixed with transparent tape and elastic band wrapping to avoid movement artifacts. The data acquisition Miograph USB® software system with windowing 32 (RMS – Root Mean Square) and gain of 2000 for each channel was used to capture the electrical potentials of the muscles evaluated in microvolts (µV). Butterworth filters were used: order 4 and band pass 20–500 Hz. The four channels were connected to active SDS500® sensors by clamps. Signal analysis was performed using Miograph USB® system software. The sensors were calibrated before data collection. The electrical potentials of the muscles were collected in accordance to international standardization of SENIAM.¹⁹

Aquatic physiotherapy

The 9 m² pool in a 16 m² room was warmed to 34°C; the patients changed their clothes in this temperature-controlled room. Each session consisted of lower limb muscle stretching, jogging in the pool (forward, backward and sideways), suspended bicycle exercises in the vertical position, stair climbing exercises, active exercises in the supine position using floats, and finally relaxation exercises.

Conventional physiotherapy

Each session consisted of lower limb stretches, hip exercises to strengthen hip adductors and abductors, supine bridge, exercises using ankle-weights to strengthen the quadriceps and when necessary, transcutaneous electrical nerve stimulation was used for pain relief.

Table 1 – Clinical and laboratory data of sickle cell disease patients submitted to two physiotherapy programs.

Patient	1	2	3	4	5	6	7	8	9	10
Group	Aquatic	Aquatic	Aquatic	Aquatic	Aquatic	Land	Land	Land	Land	Land
Age (years)	42	25	67	28	53	59	49	45	58	43
Gender	Female	Female	Male	Male	Female	Female	Female	Male	Male	Female
Type of disease	Hb SS	Hb SS	Hb SC	Hb SS	Hb SC	Hb SC	Hb SS	Hb SC	Hb SC	Hb SC
Transfusions	No	Yes	No	No	Yes	No	No	No	No	No
α-Thalassemia	Neg	Neg	Neg	Neg	Neg	Neg	Neg	Neg	Neg	H
Leukocytes (×10 ⁹ /L)	6.13	8.41	6.80	6.04	7.49	10.88	13.61	4.44	5.97	6.34
Hemoglobin (g/dL)	7.4	7.1	10.4	10.1	9.6	10.2	10.6	11.7	12.7	10.0
MCV (fL)	113.7	101.0	67.0	104.7	91.9	82.4	85.4	76.9	77.2	69.3
Reticulocytes (%)	12.68	3.51	3.59	11.81	7.48	7.56	2.70	4.83	5.55	4.90
Platelet count (×10 ⁹ /L)	331	335	508	304	392	163	175	94	275	114
Fetal hemoglobin (%)	7.3	4.0	0.2	23.7	2.8	1.4	28.1	0.5	1.0	1.2
LDH (U/L)	1680	944	405	1000	535	445	862	403	529	431
HU dose (mg/kg)	10	11	0	26	0	0	21	0	0	0

MCV: mean corpuscular volume; LDH: lactate dehydrogenase; HU: hydroxyurea; Neg: negative.




Figure 1 – Median, minimum and maximum values of the Lequesne index (Lequesne’s Algofunctional Questionnaire) and Oswestry Disability Index evaluated before and after interventions for the aquatic physiotherapy (AP) and conventional physiotherapy (CP) groups. The p₁-value refers to the comparison of numerical values over time with repeated measurements with transformation stations (ANOVA test). The p₂-value refers to the comparison between time points for each group (Wilcoxon test).

Statistical analyses

The statistical analysis system (SAS) computer program for Windows (version 9.2) and GraphPad Prism (version 5.00 – Trial) were used for statistical analysis. A p-value of 0.05 or less was considered statistically significant. The following tests were then performed.

Fisher’s exact test to compare proportions, the Mann-Whitney test to compare numerical measurements

between the two groups, ANOVA to compare numerical values over time with repeated measurements with transformation stations and the Wilcoxon test for paired samples before and after the intervention.

Results

The final sample comprised ten volunteers randomized into two groups: AP and CP. Median age was 42 years old (range:




Figure 2 – Median, minimum and maximum values of Range of Motion (RoM) of truck flexion and extension and hip flexion, extension, adduction and abduction measured by goniometry. Variables evaluated before and after interventions for the aquatic physiotherapy (AP) and conventional physiotherapy (CP) groups. The p₁-value refers to the comparison of numerical values over time with repeated measurements with transformation stations (ANOVA test). The p₂-value refers to the comparison between time points for each group (Wilcoxon test).




Figure 3 – Median, minimum and maximum values of muscle strength (MS) of the trunk flexors and extensors, and hip flexors, extensors, adductors and abductors by load cell. Variables evaluated before and after the interventions for the aquatic physiotherapy (AP) and conventional physiotherapy (CP) groups. The p_1 -value refers to the comparison of numerical values over time with repeated measurements with transformation stations (ANOVA test). The p_2 -value refers to the comparison between time points for each group (Wilcoxon test).

25–67) for the AP group and 49 years old (range: 43–59) for the CP group. The clinical and laboratory data of the participants are shown in Table 1.

Comparison of numerical values over time between the two groups showed a statistically significant difference after the intervention in respect to the Lequesne index (p -value=0.0217), ODI (p -value=0.0112), RoM of trunk extension (p -value=0.0320), trunk flexion MS (p -value=0.0459), and hip extension and abduction MS (p -value=0.0062 and p -value=0.0257, respectively). There were no significant statistical differences in the RoM of trunk and hip flexion, extension, adduction and abduction, trunk extensor MS and hip flexion and adduction MS and all SEMG variables (Figures 1–4).

Discussion

The present study aimed to evaluate two types of physiotherapy intervention for hip and lumbar spine functionality of

adult SCD patients. The dominant side of each patient was considered in the results.^{15,16}

The major limitation of this study was the recruitment of volunteers, as most of the patients live far from the center and find it very difficult to attend the clinic twice every week. Therefore, only ten patients, six compound heterozygous for Hb S and Hb C and four Hb S homozygotes, completed the physiotherapeutic program.

In view of the high frequency of avascular necrosis of the femoral head in SCD patients, the intervention was focused on the hip joint. The Lequesne questionnaire was used to evaluate the functionality of this joint. This scale assesses pain and hip function for daily activities. The results of the Lequesne Algorithmic questionnaire showed a statistically significant improvement after the intervention for both the groups. The AP group had very severe impairment (13.5 points) in the first assessment and moderate impairment (5.5 points) in the second. The CP group also had very severe impairment (11.5 points) in the first assessment and severe impairment (8 points) in the second. These results suggest that aquatic physiotherapy may lead to a greater improvement




Figure 4 – Median, minimum and maximum values of surface electromyography (SEMG) of iliocostalis, long dorsal (longissimus), gluteus maximus, gluteus medius and tensor fasciae latae muscles. Variables evaluated before and after interventions for the aquatic physiotherapy (AP) and conventional physiotherapy (CP) groups. The p_1 -value refers to the comparison of numerical values over time with repeated measurements with transformation stations (ANOVA test). The p_2 -value refers to the comparison between time points for each group (Wilcoxon test).

in hip functionality and are in agreement with Wang et al.²⁰ and Hinman et al.²¹ who observed improvement in physical function after a program of aquatic physical therapy for patients with hip and knee osteoarthritis. Other studies carried out in individuals with hip and/or knee disorders however, showed no significant differences between the two rehabilitation strategies, suggesting that both techniques are equally effective.²²⁻²⁴

Lower back pain is one of the main complaints of SCD patients. The ODI was herein used to assess lower back pain and function during daily activities. This index also showed statistically significant improvements in both study groups after the intervention. The CP group improved from moderate disability (26.5%) to minimal disability (18%); the AP group, however, despite some significant improvement in the scores of the second assessment (from 35% to 22%) showed no change in the severity of the disability caused by back pain. Longer or more frequent sessions may render better results, as has been described by others.^{24,25} These studies showed that patients who performed AP two or more times weekly had greater improvement in physical assessment scores than those who exercised only once a week.^{24,25} Nevertheless, Ariyoshi et al.²⁵

extended the program for six months and concluded that water therapy exercises were useful for patients with back pain as they provide pain relief.

In this study, a significant improvement in RoM of the trunk extension and a trend toward an improvement in trunk flexion goniometry were observed in both groups, especially in the AP group. However, no significant change was detected in either group regarding motion amplitude, probably due to the inflammatory phenomena and bone infarctions which may have caused permanent limitations.

Furthermore, late interventions may not be sufficient to improve joint RoM in this age group in which chronic degenerative hip injuries may have reached a level of severity that precludes greater joint flexibility.

Moreover, the fact that there was no significant improvement in goniometry may be a consequence of the techniques used in both groups which may favor strength gain. Thus, perhaps the program should increase the time devoted to stretching certain target muscles during therapy, thereby promoting improved muscle flexibility.

Regarding trunk MS, a significant improvement in flexion was observed in both groups after the interventions. Despite

the trend toward improvement in trunk extensor MS in the AP group, there were no significant differences between the two groups. These results are in agreement with other studies that showed improved MS after specific land-based and water-based trunk exercises.^{24,26,27}

There was a statistically significant increase in hip extension and abduction MS in both groups after the interventions. Although, hip flexion MS was unchanged in the CP group, there was a trend toward improvement in the AP group, and hip adduction MS showed a slight trend toward improvement in both groups.

Thus, albeit slight, the results of this study showed improvement of all MS variables in the AP group, in accordance with Wang et al.²⁰ who also observed improved flexibility and lower limb strength after 12 weeks of aquatic therapy, and Cochrane et al.²⁸ who observed significant improvement in pain and physical function after aquatic exercises in adults with hip and knee osteoarthritis. In another study, Rahmann²⁹ demonstrated a positive effect of a specific program of aquatic physical therapy on early recovery of strength after hip and knee surgeries. Furthermore, Hinman et al.²¹ observed a slight improvement in pain, physical function, quality of life and MS after aquatic therapy for patients with hip and knee osteoarthritis in a protocol of two sessions per week for six weeks. However, Jigami²³ concluded that both programs, land-based and water-based, even when the exercises were performed only once a week, improved overall physical activity and MS in the lower limbs of osteoarthritis patients.

Surface electromyography did not show any significant difference after the interventions of both groups. However, there seemed to be an improvement in the electromyography signal of all muscles evaluated in the AP group. The better result obtained in this group may be related to the greater amount of muscle fibers recruited in aquatic therapy added to the physical properties of water such as buoyancy and multidirectional strength. These findings are in agreement with the results reported by Kaneda et al.³⁰ who observed greater electromyography activity of all muscle movements performed in the water with floating devices.

The results of this study should be analyzed with caution as the sample size may have been a limiting factor and therefore, further studies are needed to confirm these results.

Conclusion

The results obtained here suggest that physical therapy is a resource capable of treating musculoskeletal dysfunction in SCD patients regardless of the technique used. However, exercises designed to stretch tone and strengthen the core and limb muscles carried out in the water require greater stabilization of the muscles and may justify the trend toward the better results obtained.

Funding

This work was supported by the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), and Fundação de Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES). The Hemocentro, UNICAMP, is a part of Instituto

Nacional de Ciência e Tecnologia do Sangue, Brazil (INCT do Sangue – CNPq/MCT/FAPESP).

Conflicts of interest

The authors declare no conflicts of interest.

Acknowledgments

The authors thank Prof. Margareth Castro Ozelo for kindly making the facilities at the Hemophilia Unit available; Dr. Marina Pereira Colella for helping recruit the patients, Márcia Matta, Janaína Bosso Ricciardi and Glenda Feldberg Andrade Pinto for their technical assistance; and Raquel Foglio for the editing and English revision.

REFERENCES

- Baldanzi G, Traina F, Marques Neto JF, Santos AO, Ramos CD, Olalla Saad ST. Low bone mass density is associated with hemolysis in Brazilian patients with sickle cell disease. *Clinics*. 2011;66(5):801–5.
- Almeida A, Roberts I. Bone involvement in sickle cell disease. *Br J Haematol*. 2005;129(4):482–90.
- Serarslan Y, Kalaci A, Ozkan C, Dogramaci Y, Cokluk C, Yanat AN. Morphometry of the thoracolumbar vertebrae in sickle cell disease. *Clin Neurosci*. 2010;17(2):182–6.
- Akinyoola AL, Adediran IA, Asaley CM, Bolarinwa AR. Risk factors for osteonecrosis of the femoral head in patients with sickle cell disease. *Int Orthop*. 2009;33(4):923–6.
- Huo MH, Friedlaender GE, Marsh JS. Orthopaedic manifestations of sickle-cell disease. *Yale J Biol Med*. 1990;63(3):195–207.
- Sheth S, Licursi M, Bhatia M. Sickle cell disease: time for a closer look at treatment options? *Br J Haematol*. 2013;162(4):455–64.
- Osunkwo I. An update on the recent literature on sickle cell bone disease. *Curr Opin Endocrinol Diabetes Obes*. 2013;20(6):539–46.
- Ohara DG, Ruas G, Castro SS, Martins PRJ, Walsh IA. Dor osteomuscular, perfil e qualidade de vida de indivíduos com doença falciforme. *Rev Bras Fisioter*. 2012;16(5):431–8.
- Neumayr LD, Aguilar C, Earles AN, Jergesen HE, Haberkern CM, Kammen BF, et al. Physical therapy alone compared with core decompression and physical therapy for femoral head osteonecrosis in sickle cell disease. Results of a multicenter study at a mean of three years after treatment. *J Bone Joint Surg Am*. 2006;88(12):2573–82.
- Barker AL, Talevski J, Morello RT, Brand CA, Rahmann AE, Urquhart DM. Effectiveness of aquatic exercise for musculoskeletal conditions: a meta-analysis. *Arch Phys Med Rehabil*. 2014;95(9):1776–86.
- Becker BE. Aquatic therapy: scientific foundations and clinical rehabilitation applications. *PM R*. 2009;1(9):859–72.
- Poyhonen T, Sipilä S, Keskinen KL, Hautala A, Savolainen J, Malkia E. Effects of aquatic resistance training on neuromuscular performance in healthy women. *Med Sci Sports Exerc*. 2002;34(12):2103–9.
- Doig GS, Simpson F. Randomization and allocation concealment: a practical guide for researchers. *J Crit Care*. 2005;20(2):187–91.
- Scales DC, Adhikari NK. Maintaining allocation concealment: following your SNOSE. *J Crit Care*. 2005;20(2):191–3.

15. Sadeghi H, Allard P, Prince F, Labelle H. Symmetric and limb in able-bodied gait: a review. *Gait Posture*. 2000;12(1):34-45.
16. Martin WL, Porac C. Patterns of handedness and footedness in switched and nonswitched Brazilian left-handers: cultural effects on the development of lateral preferences. *Dev Neurophysiol*. 2007;31(2):159-79.
17. Coelho RA, Siqueira FB, Ferreira PH, Ferreira ML. Responsiveness of the Brazilian-Portuguese version of the Oswestry Disability Index in subjects with low back pain. *Eur Spine J*. 2008;17(8):1101-6.
18. Marques AP. Manual de Goniometria: Segunda edição revisada e ampliada. Barueri: Manole; 2003.
19. Hermens HJ, Freriks B, Disselhorst-Klug C, Rau G. Development of recommendations for SEMG sensors and sensor placement procedures. *J Electromyogr Kinesiol*. 2000;10(5):361-74.
20. Wang TJ, Belza B, Thompson F, Whitney JD, Bennett K. Effects of aquatic exercise on flexibility, strength and aerobic fitness in adults with osteoarthritis of hip or knee. *J Adv Nurs*. 2007;57(2):141-52.
21. Hinman RS, Heywood SE, Day A. Aquatic physical therapy for hip and knee osteoarthritis: results of a single-blind randomized controlled trial. *Phys Ther*. 2007;87(1):32-43.
22. Batterham SI, Heywood S, Keating JL. Systematic review and meta-analysis comparing land and aquatic exercise for people with hip or knee arthritis on function, mobility and other health outcomes. *BMC Musculoskelet Disord*. 2011;12:123.
23. Jigami H, Sato D, Tsubaki A, Tokunaga Y, Ishikawa T, Dohmae Y, et al. Effects of weekly and fortnightly therapeutic exercise on physical function and health-related quality of life in individuals with hip osteoarthritis. *J Orthop Sci*. 2012;17(6):737-44.
24. Baena-Beato PA, Arroyo-Morales M, Delgado-Fernández M, Gatto-Cardia MC, Artero EG. Effects of different frequencies (2-3 days/week) of aquatic therapy program in adults with chronic low back pain. A non-randomized comparison trial. *Pain Med*. 2013;14(1):145-58.
25. Ariyoshi M, Sonoda K, Nagata K, Mashima T, Zenmyo M, Paku C, et al. Efficacy of aquatic exercises for patients with low-back pain. *Kurume Med J*. 1999;46(2):91-6.
26. Freitas CD, Greve JM. Estudo comparativo entre exercícios com dinamômetro isocinético e bola terapêutica na lombalgia crônica de origem mecânica. *Fisioter Pesqui*. 2008;15(4):380-6.
27. Carpenter DM, Nelson BW. Low back strengthening for the prevention and treatment of low back pain. *Med Sci Sports Exerc*. 1999;31(1):18-24.
28. Cochrane T, Davey RC, Matthes Edwards SM. Randomized controlled trial of the cost effectiveness of water-based therapy for lower limb osteoarthritis. *Health Technol Assess*. 2005;9(31), iii-iv, ix-xi, 1-114.
29. Rahmann AE, Brauer SG, Nitz JC. A specific inpatient aquatic physiotherapy program improves strength after total hip or knee replacement surgery: a randomized controlled trial. *Arch Phys Med Rehabil*. 2009;90(5):745-55.
30. Kaneda K, Sato D, Wakabayashi H, Nomura T. EMG activity of hip and trunk muscles during deep-water running. *J Electromyogr Kinesiol*. 2009;19(6):1064-70.