



Veterinary Microbiology

High frequency of hepatitis E virus infection in swine from South Brazil and close similarity to human HEV isolates



Ana Maria Passos-Castilho*, Celso Francisco Hernandes Granato

Universidade Federal de São Paulo, Departamento de Medicina, Divisão de Doenças Infecciosas, São Paulo, SP, Brazil

ARTICLE INFO

Article history:

Received 29 April 2016

Accepted 19 October 2016

Available online 3 January 2017

Associate Editor: João Araujo Jr.

Keywords:

Brazil

Genotype 3

Hepatitis E virus

Pigs

Zoonosis

ABSTRACT

Hepatitis E virus is responsible for acute and chronic liver infections worldwide. Swine hepatitis E virus has been isolated in Brazil, and a probable zoonotic transmission has been described, although data are still scarce. The aim of this study was to investigate the frequency of hepatitis E virus infection in pigs from a small-scale farm in the rural area of Paraná State, South Brazil. Fecal samples were collected from 170 pigs and screened for hepatitis E virus RNA using a duplex real-time RT-PCR targeting a highly conserved 70 nt long sequence within overlapping parts of ORF2 and ORF3 as well as a 113 nt sequence of ORF2. Positive samples with high viral loads were subjected to direct sequencing and phylogenetic analysis. hepatitis E virus RNA was detected in 34 (20.0%) of the 170 pigs following positive results in at least one set of screening real-time RT-PCR primers and probes. The swine hepatitis E virus strains clustered with the genotype hepatitis E virus-3b reference sequences in the phylogenetic analysis and showed close similarity to human hepatitis E virus isolates previously reported in Brazil.

© 2017 Sociedade Brasileira de Microbiologia. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

In endemic areas, hepatitis E virus (HEV) causes large epidemics and sporadic cases in humans, including genotype HEV-1 in Asia and Africa, genotype HEV-2 in Mexico and Africa, and genotype HEV-4 in Asia. In non-endemic areas, isolated cases of genotype HEV-3 occur in Europe, Japan and the Americas. Genotypes HEV-3 and HEV-4 are described as zoonotic, as they infect numerous mammalian species, including domestic pigs, and can be transmitted through the ingestion of raw or undercooked meat from infected animals.¹

HEV shows notable heterogeneity with several groups and genotypes. A recent consensus has classified this virus in one family of Hepeviridae, divided in 2 genera namely Orthohepevirus and Piscihepevirus. Orthohepevirus is further divided into 4 species from A to D. Orthohepevirus A is the species infecting humans and swine and other animals, such as boar, deer, mongoose, rabbit and camel. These include two genotypes isolated from humans alone (HEV-1 and HEV-2), two genotypes reported in both humans and different animal species and associated with the zoonotic cases (HEV-3 and HEV-4), two isolates from wild boar in Japan (genotype HEV-5 and HEV-6) and a single isolate from dromedary camel in Dubai

* Corresponding author.

E-mail: anampassos@gmail.com (A.M. Passos-Castilho).

<http://dx.doi.org/10.1016/j.bjm.2016.10.022>

1517-8382/© 2017 Sociedade Brasileira de Microbiologia. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

(genotype HEV-7). Rabbit HEV and closely related human isolate have been placed as a distant member in HEV-3. The moose virus appears to cluster closely to genotype HEV-3, while HEV isolates from mink to ferret virus (HEV C2) and HEV isolates from fox to rat virus (HEV C1). These viruses have not been placed in any specific genotype and need complete genomic sequences for a definite taxonomic classification. Orthohepevirus B includes all 3 avian HEV strains, Orthohepevirus C one species from rat and another one from ferret and Orthohepevirus D includes bat HEV. Piscihepevirus includes 2 trout HEV strains within a single species.^{2,3}

In Brazil, swine HEV was first isolated from pig fecal samples from the São Paulo State, Southeastern Brazil.⁴ Moreover, genotype HEV-3 was found in pigs and effluents from a pig slaughterhouse in Rio de Janeiro State, Southeastern Brazil,^{5,6} and in pigs from the Eastern Brazilian Amazon⁷ and from Southern Brazil.^{8,9} Nonetheless, hepatitis E has only been studied as a potential zoonotic disease in Latin America in the last ten years, and data on this subject are still scarce.

Data on human HEV in Brazil are limited as well. This country has been classified as moderately endemic for HEV, with seroprevalence from 1% to 4% in blood donors or the general population, 13% in individuals from an agricultural settlement in the Amazon Basin, and 15% in renal transplant recipients.^{10–13} A more recent study observed a seroprevalence of anti-HEV IgG antibodies of 10% among blood donors in the metropolitan area of Itajai Valley, Southern Brazil, a region of predominant German ancestry where cultural habits result in a high pork consumption.¹⁴ Genotype HEV-3 infection has been described in the country, both among immunocompetent and immunocompromised individuals.^{15–17}

Brazil is the fourth biggest pork exporter in the world, with an important increase in recent years. The State of Paraná has one of the largest swine productions in the country, and in 2014, this region was solely responsible for 20% of the national pig slaughter. In addition to export, pork consumption has also increased in Brazil.¹⁸ Specifically in Paraná State, the predominant European ancestry^{19,20} leads to higher pork consumption than in other parts of the country.^{21,22} The impact of such habits and the potential transmission of swine HEV to humans in this region remain unknown.

In the present study, we investigated the frequency of HEV infection in pigs from a small-scale farm in the rural area of Paraná State, South Brazil.

Materials and methods

Fecal samples

The study protocol was approved by the Institutional Committee for Ethics in the Use of Research Animals (CEUA-UNIFESP 2014/1004300914).

In September 2014, fecal samples were collected from 170 pigs from a small-scale farm in the municipality of Itapejara d'Oeste, in a rural area of Paraná State, Southern Brazil. This farm is one of the many small farms in the region, with a current population of approximately 70 sows, 5 boars and 580 pigs that are sold to slaughterhouses when raised to a median age

of 22 weeks. The sample size of 170 animals was calculated to allow determining a HEV RNA prevalence of approximately 15%⁹ with a 95% confidence interval (CI). Additionally, sampling was performed at the ages of 4, 7, 10, 13 and 16 weeks, according to the age distribution of all the pigs raised for slaughter.

RNA extraction, nested RT-PCR and quantitative RT-PCR

HEV RNA was extracted from the fecal samples using the QIAamp viral RNA mini kit (QIAGEN, Hilden, Germany). Briefly, fecal samples were first suspended in a 50% solution in ultrapure nuclease-free distilled water, then centrifuged at $20,000 \times g$ for 15 min, and the supernatant was used to extract viral RNA according to the manufacturer's instructions. Quantitative RT-PCR was performed according to a previously described modified one-step duplex real-time protocol,²³ with a primer and probe set targeting a highly conserved 70 nt long sequence within overlapping parts of ORF2 and ORF3²⁴ as well as a set specific for a 113 nt sequence of ORF2.²⁵ A previously characterized plasmid clone from a Brazilian human HEV strain (KF152884)¹⁷ was constructed with the TOPO[®] TA Cloning[®] Kit (Invitrogen, Carlsbad, CA, USA) and the described primers. Plasmid DNA was purified using the QIAprep Spin Miniprep Kit (Qiagen, Hilden, Germany) and then linearized and quantified with the Nanodrop ND-1000 instrument (Wilmington, DE, USA), followed by transcription to RNA with T7 RNA polymerase (Promega, Madison, WI, USA). Standard curves were generated using 10^{-1} – 10^{10} copies of plasmid RNA. HEV viral loads were determined based on standard curves and were reported as the log₁₀ number of copies of HEV RNA per mL of fecal suspension. The detection limit of the real-time RT-PCR was three copies of viral RNA per reaction ($2.40 \log_{10}$ copies per mL of fecal suspension), while the quantification limit was set at 10 copies of viral RNA per reaction ($3.00 \log_{10}$ copies per mL of fecal suspension). All screening reactions were run in duplicate with proper controls, whereas positive results were confirmed in separate confirmatory reactions.

The qualitative nested RT-PCR one-step reaction was conducted with primers to amplify partial regions of ORF1 and ORF2 of 287 nt and 348 nt, respectively, after second-round PCR.^{26,27} All precautions and procedures suggested to avoid the possibility of cross-contamination were employed. Amplified products were visualized in a 1.5% agarose gel stained with SYBR[®] Safe (Life Technologies, Austin, TX, USA).

Sequencing and phylogenetic analysis

Final fragments obtained from the nested RT-PCR analysis (ORF1 287 nt and ORF2 348 nt) were purified using the ExoSAP-IT PCR Clean-up Kit (GE Healthcare, Chalfont St. Giles, UK), and sequenced using the BigDye[®] Terminator v3.1 Cycle Sequencing Kit and the automated ABI 3100 DNA Sequencer (Applied Biosystems, Foster City, CA, USA), according to the manufacturer's instructions. Sequences from human and swine HEV were collected from public databases, and phylogenetic trees were constructed using the neighbor-joining method with

Table 1 – Molecular detection of HEV-RNA in pigs from the rural area of Paraná State, South Brazil, using real-time and conventional RT-PCR.

Pig no.	Age (weeks)	Real-time RT-PCR		Conventional RT-PCR	
		ORF2/3 (70 nt)	ORF2 (113 nt)	ORF1 (287 nt)	ORF2 (348 nt)
31	7	+	–	–	–
42	7	+	–	–	–
49	7	–	3.20 log ₁₀	–	+
51	7	–	+	–	–
59	7	–	+	–	–
63	7	+	–	–	–
69	10	–	+	–	–
71	10	+	–	–	–
74	10	+	–	–	–
77	10	+	–	–	–
78	10	+	–	–	–
79	10	–	6.48 log ₁₀	+	+
80	10	–	+	–	–
82	10	5.98 log ₁₀	5.54 log ₁₀	+	–
84	10	+	–	–	–
85	10	–	+	–	–
86	10	+	–	–	–
91	10	+	–	–	–
96	10	+	–	–	–
100	13	–	+	–	–
108	13	–	+	–	–
115	13	–	+	–	–
118	13	+	+	–	–
119	13	+	+	–	–
123	13	+	+	–	–
126	13	+	–	–	–
130	13	–	+	–	–
132	13	+	+	–	–
133	13	+	–	–	–
134	13	–	+	–	–
136	13	–	+	–	–
138	13	+	–	–	–
146	13	+	+	–	–
147	13	+	3.37 log ₁₀	–	+
n/N (%)		21/170 (12.4%)	20/170 (11.8%)	2/170 (1.2%)	3/170 (1.8%)

For positive samples, viral load is expressed as the log₁₀ number of copies of HEV-RNA per mL of fecal suspension if higher than the quantification limit of the real-time RT-PCR (>3.00 log₁₀ copies). Positive samples with viral load between 2.40 and 3.00 log₁₀ copies are expressed as the symbol +.

the Kimura 2-parameter model of nucleotide substitution in MEGA v. 5.0 (The Biodesign Institute, USA). Statistics was performed by bootstrap analysis with 1000 pseudoreplicates. The sequences reported in this study are available in the GenBank database under the accession numbers KP966825–KP966829.

Statistical analysis

All data were analyzed using SPSS version 11.0 (SPSS Inc., Chicago, IL, USA). Descriptive statistics consisted of the characterization of the studied population and the

Table 2 – HEV-RNA detection frequency in pigs from the rural area of Paraná State, South Brazil, by age group.

Age group (weeks)	HEV-RNA positive n (%)	HEV-RNA negative n (%)	p
4	0 (0.0)	30 (100.0)	0.001 ^a
7	6 (9.1)	30 (90.9)	
10	13 (40.6)	19 (59.4)	
13	15 (20.8)	34 (79.2)	
16	0 (0.0)	23 (100.0)	
Total	34 (20.0)	136 (80.0)	170 (100)

^a Significant at $p < 0.05$ with Pearson's Chi-square test.




Fig. 1 – Phylogenetic tree reconstructed by the neighbor-joining method with common 242-nt ORF1 sequences from 29 isolates, including 8 porcine isolates from Brazil, 1 human isolate from Brazil, and the 2 swine isolates described in this study, Brazil79.1sw and Brazil82sw (highlighted in red). The GenBank accession number in parentheses, the name of the country of origin, the species from which it was isolated, and the genotype/subtype of the isolate identify each viral strain. Bootstrap values of >50 are indicated for the major nodes as a percentage of the data obtained from 1000 replicates (bar, 0.02 substitutions per site). Major branches indicate genotypes. Avian HEV is the outgroup.

frequency of HEV-RNA detection with the respective percentages and 95% CI. The bivariate analysis to compare categorical values consisted of Pearson's Chi-square test. Non-conditional logistic regression was used to identify associations between dependent and independent variables by

the means of odds ratio (OR). For this analysis, the ages of 4 and 7 weeks were grouped to avoid empty cells, as well as the ages of 13 and 16 weeks. The statistical significance level was $p < 0.05$. All reported values are two-tailed.




Fig. 2 – Phylogenetic tree reconstructed by the neighbor-joining method with common 304-nt ORF2 sequences from 49 isolates, including 13 porcine isolates from Brazil, 4 human isolates from Brazil, and the 3 swine isolates described in this study, Brazil79sw, Brazil49sw and Brazil147sw (highlighted in red). The GenBank accession number in parentheses, the name of the country of origin, the species from which it was isolated, and the genotype/subtype of the isolate identify each viral strain. Bootstrap values of >50 are indicated for the major nodes as a percentage of the data obtained from 1000 replicates (bar, 0.02 substitutions per site). Major branches indicate genotypes. Avian HEV is the outgroup.

Results

HEV-RNA detection and sequencing

HEV-RNA was detected in 34 (20.0%) of 170 pigs following a positive result in at least one set of screening real-time RT-PCR primers and probes. Seven (20.6%) of these samples were positive with both sets of primers and probes. The 70 nt ORF2/3 set amplified 21 (12.4%) of 170 samples, while the 113 nt ORF2 set amplified 20 samples (11.8%). Among the 34 positive samples, only 4 (11.8%) presented viral loads higher than $3.00 \log_{10}$ copies per mL of fecal suspension (Table 1).

Table 2 shows the detection of HEV-RNA by age group. In the age group from 4 to 7 weeks, only 9.1% of pigs had detectable HEV-RNA, whereas 40.6% had detectable HEV-RNA at 10 weeks of age. These results show an approximate 7-fold higher risk of undergoing a HEV infection at the age of 10 weeks than at the ages of 4 or 7 weeks (OR = 6.84, 95% CI 2.29–20.48).

Due to the low viral load of the majority of the positive samples and the difference between the detection limit of the screening real-time RT-PCR and the conventional nested RT-PCR, only 4 (11.8%; 4/34) samples could be amplified with conventional nested RT-PCR and subjected to direct sequencing. Among these samples, two were amplified with the set of primers resulting in a final product of 287 nt, and three were amplified with the 384 nt set. Sample 079 was amplified using both sets of primers on the conventional nested RT-PCR.

Phylogenetic analysis

The two samples from the 287 nt ORF1 partial region shared 99% homology with each other and clustered with the genotype HEV-3b reference sequences in the phylogenetic analysis (Fig. 1). The three samples from the 384 nt ORF2 partial region also grouped with the genotype HEV-3b reference sequences in the phylogenetic analysis. Nucleotide identity among these three samples ranged from 98% to 99% (Fig. 2).

Discussion

The present data show a higher frequency of HEV infection (20.0%) in pigs than previously reported in Brazil. A study performed during 2009 in the same region with 170 fecal samples from 14 pig farms found HEV-RNA in 15.3% of samples.⁹ Another study that investigated serum, bile and fecal samples from 151 pigs from the eastern Brazilian Amazon, North Brazil, detected HEV-RNA in 9.9% of the animals.⁷ The higher infection rate observed in our study could be a result of the sanitary conditions of the small-scale pig farms in the rural area of Paraná State, where the contact of pigs of different ages may occur. However, it could also be due to the difference in methodology, since previous studies employed conventional RT-PCR techniques as opposed to the more sensitive real-time RT-PCR used in the present study. Additionally, the duplex RT-PCR technique employed in this study demonstrated the importance of using more than one set of primers and probes for higher detection during HEV screening, as only 20.6% of

the positive samples had detectable HEV-RNA with both sets of primers and probes.

Previous studies reported that HEV infection in pigs was more frequent from 12 to 16 weeks of age and that at slaughter age (20–24 weeks), the animals had already developed anti-HEV antibodies.²⁸ In the present study, HEV-RNA was more frequent in pigs aged 10 weeks (40.6%), although the frequency in pigs aged 13 was still high (20.8%). Nonetheless, none of the samples from the pigs aged 4 or 16 weeks tested positive. These observations confer with a study performed among swine herds in Rio de Janeiro, Southeast Brazil, showing that newborn pigs became susceptible to HEV between weeks 7 and 9, an age in which the serum levels of the maternal antibodies declined.⁵ These results are also in agreement with a study performed in Central Brazil in swine aged 20–30 weeks, with 81% of anti-HEV IgG positivity.²⁹

The ingestion of raw or undercooked pork has been associated with HEV infection,^{30,31} and a probable zoonotic HEV transmission has been reported in Brazil.¹⁵ Additionally, there is a known risk of HEV transmission to people who come in contact with feces from infected pigs, which have been reported as an important source of infection for slaughterhouse workers and butchers and have been associated with infection in non-endemic regions.^{32,33} HEV has also been described in sewage samples from a slaughterhouse in Southeastern Brazil.⁶

The ORF1 HEV isolates found in this study shared 88% homology with a human HEV sequence from Brazil¹⁵ and 86% to 96% homology with swine HEV sequences from Brazil.^{5,8,9} The ORF2 HEV isolates shared 86–93% homology with human HEV sequences previously characterized by our research group in renal transplant recipients in Brazil¹⁶ and 83–97% homology with swine HEV sequences from Brazil.^{5,9} Among all compared HEV sequences, the highest homology (98–99%) was with human sequences recently isolated in Southeastern Brazil from a pediatric liver transplant recipient with chronic HEV infection.¹⁷

Although more studies are needed to elucidate the real impact of swine HEV infection and zoonotic transmission in Brazil, taken together, these results reinforce the hypothesis that domestic pigs may be an important source for human hepatitis E virus infection in this setting.

Conclusions

In conclusion, this study confirms the circulation of the hepatitis E virus and shows a high frequency of HEV infection in pigs of different ages raised for slaughter in the rural area of Paraná State, South Brazil. The close similarity between the human HEV strains and those found herein indicates that adequate safety actions should be taken to prevent HEV infection when handling pigs and/or consuming pork.

Conflicts of interest

The authors declare no conflicts of interest.

Acknowledgements

This study was supported by the Fundação de Amparo à Pesquisa do Estado de São Paulo (Grant Nos. 2012/22925-3 and 2013.03701-0). We wish to thank Mr. Voitena, his family, and Veterinary Doctor Jessica Voitena for all their collaboration and technical assistance.

REFERENCES

- Kamar N, Dalton HR, Abravanel F, Izopet J. Hepatitis E virus infection. *Clin Microbiol Rev.* 2014;27:116–138.
- Smith DB, Simmonds P, Members of the International Committee on the Taxonomy of Viruses Hepeviridae Study G., et al. Consensus proposals for classification of the family Hepeviridae. *J Gen Virol.* 2014;95:2223–2232.
- Khuroo MS, Khuroo MS. Hepatitis E: an emerging global disease – from discovery towards control and cure. *J Viral Hepat.* 2016;23:68–79.
- Paiva HH, Tzaneva V, Haddad R, Yokosawa J. Molecular characterization of swine hepatitis E virus from Southeastern Brazil. *Braz J Microbiol.* 2007;38:693–698.
- dos Santos DR, Vitral CL, de Paula VS, et al. Serological and molecular evidence of hepatitis E virus in swine in Brazil. *Vet J.* 2009;182:474–480.
- dos Santos DR, de Paula VS, de Oliveira JM, Marchevsky RS, Pinto MA. Hepatitis E virus in swine and effluent samples from slaughterhouses in Brazil. *Vet Microbiol.* 2011;149:236–241.
- de Souza AJ, Gomes-Gouvêa MS, Soares MC, et al. HEV infection in swine from Eastern Brazilian Amazon: evidence of co-infection by different subtypes. *Comp Immunol Microbiol Infect Dis.* 2012;35:477–485.
- Gardinali NR, Barry AF, Otonel RA, Alfieri AF, Alfieri AA. Hepatitis E virus in liver and bile samples from slaughtered pigs of Brazil. *Mem Inst Oswaldo Cruz.* 2012;107:935–939.
- Gardinali NR, Barry AF, da Silva PF, de Souza C, Alfieri AF, Alfieri AA. Molecular detection and characterization of hepatitis E virus in naturally infected pigs from Brazilian herds. *Res Vet Sci.* 2012;93:1515–1519.
- Echevarría JM, González JE, Lewis-Ximenez LL, et al. Hepatitis E virus infection in Latin America: a review. *J Med Virol.* 2013;85:1037–1045.
- Hering T, Passos AM, Perez RM, et al. Past and current hepatitis E virus infection in renal transplant patients. *J Med Virol.* 2014;86:948–953.
- Vitral CL, da Silva-Nunes M, Pinto MA, et al. Hepatitis A and E seroprevalence and associated risk factors: a community-based cross-sectional survey in rural Amazonia. *BMC Infect Dis.* 2014;14:458.
- Passos-Castilho AM, de Sena A, Reinaldo MR, Granato CF. Hepatitis E virus infection in Brazil: results of laboratory-based surveillance from 1998 to 2013. *Rev Soc Bras Med Trop.* 2015;48:468–470.
- Passos-Castilho AM, de Sena A, Geraldo A, Spada C, Granato CF. High prevalence of hepatitis E virus antibodies among blood donors in Southern Brazil. *J Med Virol.* 2016;88:361–364.
- Lopes Dos Santos DR, Lewis-Ximenez LL, da Silva MF, de Sousa PS, Gaspar AM, Pinto MA. First report of a human autochthonous hepatitis E virus infection in Brazil. *J Clin Virol.* 2010;47:276–279.
- Passos AM, Heringer TP, Medina-Pestana JO, Ferraz ML, Granato CF. First report and molecular characterization of hepatitis E virus infection in renal transplant recipients in Brazil. *J Med Virol.* 2013;85:615–619.
- Passos-Castilho AM, Porta G, Miura IK, et al. Chronic hepatitis E virus infection in a pediatric female liver transplant recipient. *J Clin Microbiol.* 2014;52:4425–4427.
- Brazilian Institute of Geography and Statistics. *Livestock Production Estimates*; 2015. Available at: <http://www.ibge.gov.br/home/estatistica/indicadores/agropecuaria/producaoagropecuaria/default.shtm> Accessed 14.05.15.
- Brazilian Institute of Geography and Statistics (IBGE). *Brasil: 500 anos de povoamento [Brazil: 500 Years of Settlement]*. 2nd ed. Rio de Janeiro: IBGE; 2007 [in Portuguese].
- Lazier H. *Análise histórica da posse de terra no sudoeste paranaense [Historical Analysis of Land Possession in the Southeastern Paraná State]*. Fancisco Beltrão: Grafit Gráfica e Editora Ltd.; 1997 [in Portuguese].
- Schindwein MM, Kassouf AL. Análise da influência de alguns fatores socioeconômicos e demográficos no consumo domiciliar de carnes no Brasil [Analysis of the Influence of Some Socio-Economic and Demographic Factors in Household Consumption of Meat in Brazil]. *Rev Econ Sociol Rural.* 2006;44:549–572 [in Portuguese].
- Verbeke W, Perez-Cueto FJ, Barcellos MD, Krystallis A, Grunert KG. European citizen and consumer attitudes and preferences regarding beef and pork. *Meat Sci.* 2010;84:284–292.
- Vasickova P, Kralik P, Slana I, Pavlik I. Optimisation of a triplex real time RT-PCR for detection of hepatitis E virus RNA and validation on biological samples. *J Virol Methods.* 2012;180:38–42.
- Jothikumar N, Cromeans TL, Robertson BH, Meng XJ, Hill VR. A broadly reactive one-step real-time RT-PCR assay for rapid and sensitive detection of hepatitis E virus. *J Virol Methods.* 2006;131:65–71.
- Gyarmati P, Mohammed N, Norder H, Blomberg J, Belák S, Widén F. Universal detection of hepatitis E virus by two real-time PCR assays: TaqMan and Primer-Probe Energy Transfer. *J Virol Methods.* 2007;146:226–235.
- Wang Y, Ling R, Erker JC, et al. A divergent genotype of hepatitis E virus in Chinese patients with acute hepatitis. *J Gen Virol.* 1999;80:169–177.
- Huang FF, Haqshenas G, Guenette DK, et al. Detection by reverse transcription-PCR and genetic characterization of field isolates of swine hepatitis E virus from pigs in different geographic regions of the United States. *J Clin Microbiol.* 2002;40:1326–1332.
- Nakai I, Kato K, Miyazaki A, et al. Different fecal shedding patterns of two common strains of hepatitis E virus at three Japanese swine farms. *Am J Trop Med Hyg.* 2006;75:1171–1177.
- Guimaraes FR. Hepatitis E virus antibodies in swine herds of Mato Grosso state, Central Brazil. *Braz J Microbiol.* 2005;36:223–226.
- Colson P, Borentain P, Queyriaux B, et al. Pig liver sausage as a source of hepatitis E virus transmission to humans. *J Infect Dis.* 2010;202:825–834.
- Legrand-Abravanel F, Kamar N, Sandres-Saune K, et al. Characteristics of autochthonous hepatitis E virus infection in solid-organ transplant recipients in France. *J Infect Dis.* 2010;202:835–844.
- Hepatitis E virus in swine and meat carcasses. *Br J Gen Pract.* 2005;55:557–558.
- Perez-Gracia MT, Mateos ML, Galiana C, et al. Autochthonous hepatitis E infection in a slaughterhouse worker. *Am J Trop Med Hyg.* 2007;77:893–896.