

## ERRATUM

*Neotropical Entomology* 34(4):895-902 (2005)

The Presence of the Sexual Partner and Nutritional Condition Alter the *Anastrepha obliqua* MacQuart (Diptera: Tephritidae) Protein Discrimination Threshold *Carla Cresoni-Pereira and Fernando S. Zucoloto*

Please replace the following paragraph, at the page 897 by the one bellow:

The yeast amounts in the diet found as thresholds for the different groups were statistically compared (Figs. 3 and 6). In the non-deprived females group, statistical differences were found between the 1.6 g quantity of yeast (FF, F and FM groups threshold) and 1.8 g of yeast (FMS group threshold) (Fig. 3). In deprived females group, statistical differences were found among all yeast quantities found as threshold (0.7 g to F group; 1.2 g to **FF** group and 1.5 g to FMS group) (Fig. 3).

The yeast amounts in the diet found as thresholds for the different groups were statistically compared (Figs. 3 and 6). In the non-deprived females group, statistical differences were found between the 1.6 g quantity of yeast (FF, F and FM groups threshold) and 1.8 g of yeast (FMS group threshold) (Fig. 3). In deprived females group, statistical differences were found among all yeast quantities found as threshold (0.7 g to F group; 1.2 g to **FM** group and 1.5 g to FMS group) (Fig. 3).