# **Neotropical Entomology**

ISSN: 1519-566X

journal homepage: www.scielo.br/ne



# SCIENTIFIC NOTE

# First Record of *Anastrepha serpentina* (Wiedemann) (Diptera: Tephritidae) in Citrus in Brazil

WP LEMOS<sup>1</sup>, RA DA SILVA<sup>2</sup>, SCA ARAÚJO<sup>1</sup>, ELA OLIVEIRA<sup>1</sup>, WR DA SILVA<sup>3</sup>

<sup>1</sup>Embrapa Amazônia Oriental, Belém, PA, Brasil

#### Keywords

Fruit fly, host plant, Amazonian, Pará, Belém, Capitão Poço, Tomé-Açu

#### Correspondence

RICARDO A DA SILVA, Embrapa Amapá, Rodovia JK, km 5, n. 2600, 68903-419, Macapá, AP, Brasil; adaime@cpafap.embrapa.br

Edited by Roberto A Zucchi – ESALQ/USP

Received 21 December 2010 and accepted 18 April 2011

#### **Abstract**

Anastrepha serpentina (Wiedemann) is recorded for the first time in citrus (Rutaceae) in Brazil. Specimens were obtained from sweet orange (*Citrus sinensis*) sampled in the municipalities of Belém and Capitão Poço, and from mandarin orange (*Citrus reticulata*) from Tomé-Açu, state of Pará, Brazil.

Anastrepha serpentina (Wiedemann) occurs in 20 Brazilian states (Zucchi 2008) and in the Federal District (Zahler 1991) associated with 16 host species belonging to Sapotaceae (10 hosts), Rubiaceae (2), Anacardiaceae (1), Clusiaceae (1) Hippocrateaceae (1) and Moraceae (1) (Zucchi 2008). The preferred hosts of *A. serpentina* in Brazil belong to the Sapotaceae, particularly species of *Pouteria* and *Manilkara*, with four and three host species, respectively.

This paper reports the first record of *A. serpentina* in fruits of Rutaceae in Brazil. Data were obtained from an intensive survey of potential host fruits of *Anastrepha* species carried out at major growing areas in the state of Pará (eastern Amazon region), from January 2008 to August 2010.

Two males and one female of *A. serpentina* were reared from sweet oranges, *Citrus sinensis*, purchased at a street market in Belém, and 23 females and 14 males from Capitão Poço (Table 1). The highest infestation was 8.6 puparia/kg. However, no specimens of *A. serpentina* were reared from four samples, but from one of them one specimen of *Doryctobracon areolatus* (Szépligeti)

emerged. Also for one sample, only two males were obtained, which were identified as *A. serpentina* based on external features. Although puparia have been obtained, no fly or parasitoid emergence was observed for three samples. One unique female of *A. serpentina* was reared from mandarin (*Citrus reticulata*) from Tomé-Açu (Table 1). All specimens sampled were identified by Wilson Rodrigues da Silva and confirmed by Dr. Miguel Francisco de Souza Filho (Instituto Biológico, Campinas, SP). Voucher specimens are deposited at the Embrapa Amapá collection.

Twenty-five other samples, totaling 49.26 kg and 386 fruits of five *Citrus* species – the sweet orange *C. sinensis*; the mandarin orange *C. reticulata*; the lime *C. aurantifolia*; the lemon *C. limon* and the Murcott mandarin *C. reticulata* x *C. sinensis* were not infested by fruit flies.

This report represents a new fruit fly species association related to the host range evolution. The record is not a casual fact, since *A. serpentina* females were reared from one mandarin orange sample in Tomé-Açu (2008), and 11 orange samples – one in Belém (2008) and 10 in Capitão Poço (2009). These three municipalities are

<sup>&</sup>lt;sup>2</sup>Embrapa Amapá, Macapá, AP, Brasil

<sup>&</sup>lt;sup>3</sup>Secretaria de Estado da Educação do Amapá, Macapá, AP, Brasil

Vernacular names Scientific names	Samples	Municipalities	Dates	Fruits (n)	Mass (g)	Puparia (n)	Infestations (puparia/kg)	Females (n)	Males (n)
Sweet orange Citrus sinensis (L.) Osbeck	1	Belém	05/16/2008	8	1,300	5	3.9	1	2
	2	Capitão Poço	03/25/2009	29	5,760	12	2.1	0	0
	3	Capitão Poço	03/25/2009	40	6,950	12	1.7	3	2
	4	Capitão Poço	03/26/2009	32	3,650	14	3.8	3	4
	5	Capitão Poço	03/26/2009	27	3,390	4	1.2	1	2
	6	Capitão Poço	03/26/2009	11	1,250	5	4.0	3	0
	7	Capitão Poço	03/26/2009	21	3,470	13	3.8	5	0
	8	Capitão Poço	03/26/2009	13	1,770	5	2.8	2	1
	9	Capitão Poço	03/26/2009	10	1,200	2	1.7	2	0
	10	Capitão Poço	03/26/2009	12	1,100	3	2.7	1	2
	11	Capitão Poço	08/19/2009	19	3,150	22	7.0	3	1
	12	Capitão Poço	08/19/2009	19	1,700	5	2.9	0	2
	13	Capitão Poço	08/19/2009	15	1,950	1	0.5	0	0
	14	Capitão Poço	08/19/2009	21	1,750	15	8.6	0	0
	15	Capitão Poço	08/19/2009	12	1,800	5	2.8	0	0
Mandarin orange	4.6	T	04/00/2000	10	4.254	4	2.0	4	

04/08/2008

Table 1 *Citrus* infestation by *Anastrepha serpentina* in the state of Pará.

around 200 km far from one another, forming a triangle. Capitão Poço has the largest planted area with *Citrus* in the state of Pará, which may have favored the new association in here reported due to the high availability of *Citrus* plants during the year.

16

Tomé-Açu

Anastrepha fraterculus (Wiedemann), A. obliqua (Macquart), A. striata Schiner, A. turpiniae Stone and Ceratitis capitata (Wiedemann) are reported in citrus in Brazil (Raga 2005, Zucchi 2008). However, for the state of São Paulo, the largest citrus producer in Brazil, only A. fraterculus and C. capitata are economically important (Raga 2005).

Considering that Brazil is the largest citrus producer in the world (about 800,000 ha of citrus groves), and that the state of Pará is the seventh Brazilian citrus producer and the first in the North region (12,277 hectares) (Agrianual 2010), this record strengthens the need to conduct applied research to study the spatial and temporal distribution and potential damage of *A. serpentina* in *Citrus* in this state.

### **Acknowledgments**

Citrus reticulata Blanco

To MSc. Júlia Daniela Braga Pereira for her contribution to this manuscript, Dr Miguel F. de Souza Filho for

confirming the identification of *Anastrepha serpentina*, and Dr Moacyr B. Dias Filho for reviewing the English version of an early draft of this manuscript. To "Conselho Nacional de Desenvolvimento Científico e Tecnológico" (CNPq) for financial support.

3.0

## References

10

1,354

Agrianual: anuário da agricultura brasileira (2010) São Paulo, FNP Consultoria & Comércio, 520p.

Raga A (2005) Incidência, monitoramento e controle de moscasdas-frutas na citricultura paulista. Laranja 26: 307-322.

Zahler PM (1991) Moscas-das-frutas (Diptera, Tephritidae) em dois pomares de manga (*Mangifera indica*) do Distrito Federal: levantamento de espécies e flutuação populacional. Rev Ceres 38: 206-216.

Zucchi RA (2008) Fruit flies in Brazil - *Anastrepha* species and their host plants. Available at: <www.lea.esalq.usp.br/anastrepha/> Accessed in 05 July 2010.

0

1