











# Electric eels galore: microsatellite markers for population studies

 Lenice Souza-Shibatta<sup>1</sup>,  Dhiego G. Ferreira<sup>2</sup>,  Kátia F. Santos<sup>2</sup>,  
 Bruno A. Galindo<sup>2</sup>,  Oscar A. Shibatta<sup>3</sup>,  Silvia H. Sofia<sup>4</sup>,  
 Renata M. Giacomini<sup>5</sup>,  Douglas A. Bastos<sup>6</sup>,  
 Raimundo N. G. Mendes-Júnior<sup>7</sup> and  
 Carlos David de Santana<sup>8</sup>

Correspondence:  
Lenice Souza-Shibatta  
[lenicesouza@hotmail.com](mailto:lenicesouza@hotmail.com)

Submitted August 19, 2020

Accepted October 1, 2020

by Claudio Oliveira

Epub November 16, 2020

Fourteen novel microsatellite loci are described and characterized in two species of electric eels, *Electrophorus varii* and *E. voltai* from floodplains and rivers of the Amazon rainforest. These loci are polymorphic, highly informative, and have the capacity to detect reliable levels of genetic diversity. Likewise, the high combined probability of paternity exclusion value and low combined probability of genetic identity value obtained demonstrate that the new set of loci displays suitability for paternity studies on electric eels. In addition, the cross-amplification of electric eel species implies that it may also be useful in the study of the closely related *E. electricus*, and to other Neotropical electric fishes (Gymnotiformes) species as tested herein.

**Keywords:** Amazon Rainforest, *Electrophorus*, Genetic diversity, Gymnotiformes, SSR.

Online version ISSN 1982-0224

Print version ISSN 1679-6225

Neotrop. Ichthyol.  
vol. 18, no. 4, Maringá 2020

1 Laboratório de Sistemática Molecular, Programa de Pós-Graduação em Ciências Biológicas, Universidade Estadual de Londrina, Rod. Celso Garcia Cid, km 380, 86051-970 Londrina, PR, Brazil. (LSS) [lenicesouza@hotmail.com](mailto:lenicesouza@hotmail.com) (corresponding author).

2 Laboratório de Genética e Conservação (GECON), Universidade Estadual do Norte Paraná, Rua Portugal, 340 86300-000 Cornélio Procopio, PR, Brazil. (DGF) [dhiegouenp@gmail.com](mailto:dhiegouenp@gmail.com); (KFS) [kfabianadossantos@gmail.com](mailto:kfabianadossantos@gmail.com); (BAG) [bruno@uenp.edu.br](mailto:bruno@uenp.edu.br).

3 Universidade Estadual de Londrina, Museu de Zoologia, Departamento de Biologia Animal e Vegetal, Centro de Ciências Biológicas, 86051-990 Londrina, PR, Brazil. (OAS) [shibatta@uel.br](mailto:shibatta@uel.br).

4 Laboratório de Genética e Ecologia Animal (LAGEA), Departamento de Biologia Geral, Universidade Estadual de Londrina, Rod. Celso Garcia Cid, km 380, 86051-970 Londrina, PR, Brazil. (SHS) [shsofiabelh@gmail.com](mailto:shsofiabelh@gmail.com).

5 Departamento de Biologia Geral, Universidade Estadual de Londrina, 86051-990 Londrina, PR, Brazil. (RMG) [giacomini.rm@gmail.com](mailto:giacomini.rm@gmail.com).

6 Programa de Pós-Graduação em Ciências Biológicas (BADPI), Instituto Nacional de Pesquisas da Amazônia, 69060-001 Manaus, AM, Brazil. (DAB) [avizdoug@gmail.com](mailto:avizdoug@gmail.com).

7 RESEX do Rio Cajari, Instituto Chico Mendes de Conservação da Biodiversidade, Rua Hamilton Silva, 1570, 68906-440 Macapá, AP, Brazil. (RNMJ) [raimundo.mendes-junior@icmbio.gov.br](mailto:raimundo.mendes-junior@icmbio.gov.br).

8 Division of Fishes, Department of Vertebrate Zoology, MRC-159, National Museum of Natural History, P.O. Box 37012, Smithsonian Institution, 20013-7012 Washington, DC, WA, USA. (CDS) [desantanac@si.edu](mailto:desantanac@si.edu).

Catorze novos loci microsatélites são descritos e caracterizados em duas espécies de poraquês, *Electrophorus varii* e *E. voltai* de planícies alagadas e rios da floresta amazônica. Esses loci são polimórficos, altamente informativos e têm a capacidade de detectar níveis confiáveis de diversidade genética. Da mesma forma, o alto valor de exclusão de paternidade combinado com a baixa probabilidade de identidade genética demonstra que o novo conjunto de loci exibe adequação para estudos de paternidade em poraquês. Além disso, a amplificação cruzada de espécies de peixes elétricos implica que também pode ser útil no estudo da espécie intimamente relacionada *E. electricus*, e de outras espécies de peixes elétricos neotropicais (Gymnotiformes).

**Palavras-chave:** Diversidade genética, *Electrophorus*, Floresta amazônica, Gymnotiformes, SSR.

## INTRODUCTION

Electric eels (*Electrophorus* Gill, 1864) share with other species of Neotropical electric fishes (Gymnotiformes) a specialized electrogenic-electrosensory system used to navigate, and communication (Crampton, 2019). In addition to low-voltage electric organ discharges (EODs), electric eels generate high-voltage EODs for stunning prey and defense, as reported in the field by Humboldt in the 18th Century, and elegantly demonstrated in the laboratory by Catania (2019). For centuries, electric eels captivate minds, inspire scientific innovation, like the electric battery, which has been used as a model for understanding bioelectrogenesis (Finger, Piccolino, 2011; Gallant *et al.*, 2014). Despite the broad public and scientific community interest, only recently species diversity on *Electrophorus* began to be explored in extent (de Santana *et al.*, 2019). As a result, three electric eel species occurring in very distinct ecological environments were recognized: *E. electricus* (Linnaeus, 1766) and *E. voltai* de Santana, Wosiacki, Crampton, Sabaj, Dillman, Castro e Castro, Bastos & Vari, 2019 from Brazilian and Guyana shields in Highlands Amazon and *E. varii* de Santana, Wosiacki, Crampton, Sabaj, Dillman, Mendes-Júnior & Castro e Castro, 2019 from the Lowlands Amazon (de Santana *et al.*, 2019).

The new finds offer an opportunity to study the genetics of populations of those distinct ecological and unique animals by characterizing their genetic variation, within and between populations, and the forces that affect their frequencies, such as migration, mutation, selection, and genetic drift. An excellent way to study the genetic composition of natural fish populations is by using molecular markers, which are powerful tools for quantifying genetic variation in individuals and populations, contributing to the management and conservation of species (Allendorf *et al.*, 2010). According to Zane *et al.* (2002), the microsatellites (SSR – Simple Sequence Repeats), for instance, are considered useful for population studies because they are highly polymorphic markers. The population genetic analysis of species in the wild is of paramount importance for elucidating the factors and conditions that allow populations and species to be maintained and in the development of a strategy for its effective management (Moysés *et al.*, 2005).

Published population genetic studies in Neotropical electric fishes are inexistent, and only a few attempts to develop microsatellite primers for Gymnotiformes were made (e.g. Moysés *et al.*, 2005).

This study aims to develop candidate microsatellite loci to accurately assess genetic diversity and help in future studies of population genetics of electric eels. Thus, this paper reports the development and characterization of novel microsatellite loci for *E. varii* and evaluates it in *E. voltai* to cross-amplification. Additionally, the primers were tested for cross-amplification in four species across Gymnotiformes.

## MATERIAL AND METHODS

A partial enriched genomic library was constructed, and microsatellites were isolated and characterized following the protocol of Billotte *et al.* (1999). Tissue samples from *E. varii* and *E. voltai* were donated by the Instituto Nacional de Pesquisas da Amazônia (INPA), with invoice number: 009/96. Total genomic DNA was extracted from muscle tissue from a sample of *E. varii* (INPA 41112), according to Almeida (Almeida *et al.*, 2010). Genomic DNA (5 µg) was digested, and the blunt-ended fragments were ligated to the adaptors (Edwards *et al.*, 1996). Fragments were selected, amplified, and cloned into pGem-T Easy (Promega; www.promega.com) vectors using 5 µL of the amplification product, 50 ng of vector, and 1 U of T4 DNA ligase in reaction buffer at 4°C (overnight). Cloning products were used to transform *Escherichia coli* (DH5 – α lineage) cells. The recombinant clones were selected and sequenced on an ABI 3500 XL automated sequencer. Sequences were analyzed, and primers were designed according to Hall (1999) and Rozen, Skaletsky (2000), respectively. The selected forward primers were marked with the M13 at the 5' end (Schuelke, 2000). To test the potential presence of hairpin structures and problems with the primer-dimer, we follow the protocol of Vallone, Butler (2004). PCR amplifications were carried out on a panel consisting of 13 individuals of *E. varii* (INPA 41112 – 41122 and INPA 41124 – 41125) from three localities along the Curiaú River; and 14 individuals of *E. voltai* (LIA 4802 – 4806; INPA 41123; INPA 050453) – five from two localities of the Xingu River, one specimen collected in the Curiaú River and eight collected in the Iriri River. All specimens of electric eels were collected in the Amazon basin, Brazil. Cross-amplification tests were performed using four other Gymnotiformes species whose voucher specimens are deposited in the Museu de Zoologia da Universidade Estadual de Londrina (MZUEL) as follows: *Apteronotus cf. caudimaculosus* de Santana, 2003 (n=4; MZUEL 09538; Apteronotidae); *Eigenmannia trilineata* López & Castello, 1966 (n=5; MZUEL 09552; Sternopygidae); *Gymnotus sylvius* Albert & Fernandes-Matioli, 1999 (n=5; MZUEL 09546; Gymnotidae); and *Sternopygus macrurus* (Bloch & Schneider, 1801) (n=5; MZUEL 09454; Sternopygidae), all collected in the Laranjinha River, Paraná river basin. Reactions were performed according to Apolinário-Silva *et al.* (2018). Amplifications were made with an initial denaturation step at 94°C for 4 min, followed by 35 cycles at 94°C for 40 s, 48°C, 54°C, or 60°C (Tabs. 1–2) for 1 min, 72°C for 1 min, and a final extension at 72°C for 30 min. The PCR products were submitted to electrophoresis on an automated sequencer. GeneScan 600 Liz (Applied Biosystems) was used as the molecular weight standard.

Individuals were genotyped with GeneMarker 1.85 (SoftGenetics, State College, PA), followed by manually editing. Tests for Hardy-Weinberg Equilibrium (HWE) and the presence of linkage disequilibrium among the pairs of loci were calculated using GENEPOP 4.0.10;  $P$  values were subsequently adjusted applying the sequential Bonferroni correction (Rice, 1989). GenAEx v.6.41 was used to estimate the observed ( $H_o$ ) and expected ( $H_e$ ) heterozygosities and the average number of alleles per locus. The paternity exclusion probability ( $Q$ ) (Weir, 1996) and genetic identity probabilities ( $I$ ) (Paetkau *et al.*, 1995) were estimated using Identity 1.0. Estimates of the polymorphic information content (PIC) and potential null alleles were obtained through Cervus v.3.0 and Micro-Checker v.2.2.3, respectively. Default settings were used for all tests.

## RESULTS

A set of 13 polymorphic and highly informative microsatellite loci for genetic studies of populations of *Electrophorus* were developed: a total of 45 out of 96 clones sequenced contained microsatellite regions, with 25 being suitable for primer design and PCR reactions. After testing different amplification conditions, 14 loci (almost all dinucleotide repeats) were successfully amplified. From those, one was monomorphic, and 13 were polymorphic for two electric eel species.

In *E. varii*, a total of 85 different alleles were detected, varied from 2 (*Elec24*) to 15 (*Elec39*), with an average of 6.4 alleles per locus. The observed and expected heterozygosity ranged from 0.000 (*Elec24*) to 1.000 (*Elec14*) and from 0.334 (*Elec49*) to 0.902 (*Elec39*), respectively. After sequential Bonferroni correction for multiple comparisons ( $\alpha = 0.05$ ,  $k = 91$ ), no evidence of linkage disequilibrium between any pair of loci examined was observed. In the HWE tests, two loci, *Elec24* and *Elec241*, presented significant deviation after correction for multiple tests (sequential Bonferroni correction  $\alpha = 0.05$  and  $k = 14$ ). These loci were also the only ones showing possible null alleles, inferred from excess homozygous genotypes, explaining the observed deviation from HWE. It was observed that the same loci that had a significant deviation in the HWE, plus loci *Elec22* and *Elec31*, also had significant values of the endogamic coefficient ( $F_{IS}$ ; Tab. 1). The mean PIC for the 13 polymorphic loci was 0.572 following a scale proposed by Botstein *et al.* (1980), 10 loci (*Elec12*, *Elec14*, *Elec 21*, *Elec31*, *Elec39*, *Elec43*, *Elec53*, *Elec241*, *Elec246* and *Elec247*) were highly informative and three loci (*Elec22*, *Elec24* and *Elec49*) were moderately informative. The probabilities of identity and paternity exclusion were equal to  $2.665^{-12}$  and 0.999, in that order (Tab. 1).

All 14 microsatellite primers developed for *E. varii* were successfully cross-amplified in *E. voltai*. Thirteen are polymorphic loci and produced a total of 74 different alleles, with allele number ranging from 2 (*Elec31* and *Elec451*) to 12 (*Elec49*), with an average of 5.2 alleles per locus (Tab. 2). The observed and expected heterozygosity varied from 0.071 (*Elec12*, *Ele24* and *Elec451*) to 1.000 (*Elec14*) and from 0.069 (*Elec451*) to 0.908 (*Elec49*), correspondingly. After Bonferroni sequential correction for multiple comparisons ( $\alpha = 0.05$ ,  $k = 91$ ), no evidence of linkage disequilibrium between any pair of loci examined was detected.

**TABLE 1** | Description and characterization of 14 microsatellite loci isolated from *Electrophorus varii*. Flanking primers,  $T_a$  = optimal annealing temperatures,  $k$  = number of alleles,  $H_o$  = observed heterozygosity,  $H_e$  = expected heterozygosity estimated from 13 individuals,  $Q$  = paternity exclusion probability,  $I$  = probability of genetic identity,  $F_{IS}$  = endogamy coefficient, PIC = polymorphic information content, GenBank accession numbers. \* Significant value for the endogamy coefficient ( $F_{IS}$ ).

Locus	Sequence repeat	Primer sequences (5' - 3')	$T_a$ (°C)	$k$	Allele size range (bp)	$H_o$	$H_e$	PIC	( $Q$ )	( $I$ )	$F_{IS}$	Genebank Accession numbers
<i>Elec12</i>	(CA) <sub>14</sub>	F: CAGTTCAGTAGCAGGAGTATACAGG	52°	7	203 – 241	0.769	0.692	0.661	0.483	0.125	-0.071	MN967054
		R: TTAGTGTGAGGTGGATTAACAATG										
<i>Elec14</i>	(TG) <sub>28</sub>	F: GCTCTGTTGTGTTACGGC	52°	9	191 – 260	1.000	0.795	0.774	0.663	0.049	-0.216	MN967055
		R: TGA CT CGCAGGCTAACAGG										
<i>Elec22</i>	(TG) <sub>15</sub>	F: GGAGCAGCAACCGACTC	48°	4	171 – 177	0.231	0.388	0.363	0.214	0.399	0.437*	MN967056
		R: GGCACACTACAGTCTCTCCAA										
<i>Elec24</i>	(GT)13(GAAA) <sub>4</sub>	F: GATACTTCGAGCTCAGTCTTAG	56°	2	214 – 216	0.000	0.355	0.292	0.146	0.479	1.000*	MN967057
		R: TCCTCATGTATCCATTACCAAG										
<i>Elec31</i>	(AG) <sub>18</sub>	F: TTGATCAITTAGCGTGGACTTAAC	45°	5	144 – 166	0.538	0.751	0.711	0.524	0.102	0.319*	MN967058
		R: AGGCCACACTACTAATCAGAACG										
<i>Elec39</i>	(GT) <sub>37</sub>	F: TCCAGGGACAGGACGTTG	56°	15	166 – 228	0.846	0.902	0.895	0.805	0.017	0.102	MN967059
		R: TCCAGCACACTCAGGTAGAGG										
<i>Elec43</i>	(TG) <sub>16</sub>	F: CCTGTTAGGCTGGTTAGATAATATG	60°	5	263 – 279	0.769	0.701	0.649	0.451	0.141	0.057	MN967060
		R: CAAGAAGCTAGACGCCATGC										
<i>Elec49</i>	(GT) <sub>17</sub>	F: ACTATCAGGTCTCAAAGGATTTTC	56°	4	178 – 202	0.231	0.334	0.317	0.184	0.460	0.345	MN96705461
		R: GAGCACAGATCTGGTCATCTAGG										
<i>Elec53</i>	(GA)10(TG)8(AG) <sub>19</sub>	F: GCAATATGATTCTGTTGACTTCG	52°	6	177 – 225	0.692	0.710	0.662	0.472	0.131	0.064	MN967062
		R: GCACTGCCTGACAGATGG										
<i>Elec241</i>	(GT) <sub>14</sub>	F: CTGGTGGAGTTGATTACAGAGAG	56°	8	147 – 215	0.455	0.740	0.706	0.604	0.070	0.245*	MN967063
		R: AACTAACATATCCATCCACAAG										
<i>Elec244</i>	(TG) <sub>14</sub>	F: GAGGTGGATTAACAATGTAACTGG	56°	8	202 – 243	0.714	0.769	0.732	0.567	0.084	0.024	MN967064
		R: CAGTTCAGTAGCAGGAGTATACAGG										
<i>Elec246</i>	(TG) <sub>24</sub>	F: CTCGGTCTCCAGTCTTGC	52°	4	280 – 338	0.692	0.678	0.613	0.400	0.168	0.018	MN967065
		R: GTGACTCGCAGGCTAACAGG										
<i>Elec247</i>	(TG) <sub>13</sub>	F: TTAGTGTGAGGTGGATTAACAATG	56°	7	156 – 196	0.538	0.689	0.639	0.450	0.146	0.256	MN967066
		R: CATA CAT ATGCACGTTCTCTTGC										
<i>Elec451</i>	(GT) <sub>14</sub>	F: GTAAGGAGAGCCGACAGCAC	52°	1	169	–	–	–	–	–	–	MN967067
		R: AAGGCAGTGTGGAGTACC										
All loci				85		0.538	0.607	0.572	0.999	2.665 <sup>-12</sup>	0.153*	

Hardy-Weinberg Equilibrium deviations were significant for four loci (*Elec22*, *Elec24*, *Elec241* and *Elec247*) after correction for multiple tests (sequential Bonferroni correction,  $\alpha = 0.05$  and  $k = 14$ ). At the same time, these loci were the only ones showing null alleles (inferred from excess homozygous genotypes), which could explain the observed deviation from HWE. In addition, these same loci, plus *Elec12* and *Elec53*, also showed significant values of the inbreeding coefficient ( $F_{IS}$ ; Tab. 2). The mean Polymorphic Information Content (PIC) for the 13 loci was 0.500, indicating that the loci set is highly informative (Tab. 2). Seven loci (*Elec14*, *Elec39*, *Elec43*, *Elec49*, *Elec53*, *Elec241* and *Elec246*) were highly informative (PIC > 0.5); four loci (*Elec22*, *Elec24*, *Elec31* and *Elec247*) were moderately informative (PIC > 0.25 and < 0.5); and two loci

**TABLE 2** | Cross-amplification of 14 microsatellite loci and genetic diversity per locus in *Electrophorus voltai*. Flanking primers,  $k$  = number of alleles,  $H_o$  = observed heterozygosity,  $H_e$  = expected heterozygosity estimated from 14 individuals,  $Q$  = paternity exclusion probability,  $I$  = probability of genetic identity,  $F_{IS}$  = endogamy coefficient, PIC = polymorphic information content. \* Significant value for the endogamy coefficient ( $F_{IS}$ ).

Locus name	$k$	Allele size range (bp)	$H_o$	$H_e$	PIC	( $Q$ )	( $I$ )	$F_{IS}$
<i>Elec12</i>	3	203 – 211	0.071	0.135	0.131	0.068	0.752	0.500*
<i>Elec14</i>	6	209 – 260	1.000	0.719	0.679	0.492	0.119	-0.358
<i>Elec22</i>	4	173 – 179	0.143	0.403	0.364	0.209	0.395	0.666*
<i>Elec24</i>	3	212 – 216	0.071	0.564	0.466	0.266	0.287	0.881*
<i>Elec31</i>	2	148 – 166	0.500	0.375	0.305	0.152	0.460	-0.300
<i>Elec39</i>	6	150 – 172	0.857	0.760	0.724	0.547	0.093	-0.090
<i>Elec43</i>	7	251 – 315	0.760	0.791	0.724	0.594	0.074	0.228
<i>Elec 49</i>	12	220 – 242	0.786	0.908	0.901	0.812	0.015	0.171
<i>Elec53</i>	8	199 – 225	0.643	0.832	0.811	0.668	0.048	0.261*
<i>Elec241</i>	10	161 – 191	0.538	0.861	0.846	0.749	0.027	0.408*
<i>Elec244</i>	1	212	0.000	0.000	–	–	–	–
<i>Elec246</i>	5	280 – 362	0.500	0.548	0.516	0.338	0.236	0.125
<i>Elec247</i>	5	154 – 162	0.214	0.508	0.478	0.306	0.272	0.602*
<i>Elec451</i>	2	169 – 177	0.071	0.069	0.067	0.033	0.869	0.001
All loci	74	–	0.429	0.532	0.500	0.999	2.981 <sup>-11</sup>	0.228*

(*Elec12* and *Elec451*) had low informative potential ( $PIC < 0.2$ ). The loci set showed a low value of genetic identity combined probability ( $2.9 \times 10^{-11}$ ) and high-shared probabilities of paternity exclusion (0.999), which suggest a high discriminatory power for population genetic studies (Tab. 2).

Cross-amplification testing of all 14 *Electrophorus* loci in four other Gymnotiformes was conducted. Six microsatellite loci successfully amplified in *Apteronotus albifrons* (Linnaeus, 1766) (*Elec12*, *Elec39*, *Elec49*, *Elec53*, *Elec247* and *Elec451*) and *E. trilineata* (*Elec12*, *Elec14*, *Elec39*, *Elec49*, *Elec247*, *Elec451*). Three effectively worked in *G. sylvius* (*Elec12*, *Elec53*, *Elec247*), and two in *S. macrurus* (*Elec12*, *Elec49*). The locus *Elec12* was polymorphic for all species tested, ranging from three (*G. sylvius* and *S. macrurus*) to five (*A. albifrons* and *E. trilineata*) alleles per locus. *Elec47* presented four alleles in *G. sylvius*, three in *A. albifrons*, and two in *E. trilineata*. On the other hand, the microsatellite loci *Elec14*, *Elec39*, and *Elec451* were monomorphic for tested species, and loci *Elec22*, *Elec24*, *Elec31*, *Elec43*, *Elec241*, and *Elec244*, did not amplify for any of the four tested species.

## DISCUSSION

Deviations of the HWE and significant  $F_{IS}$  values for some loci, mainly in *E. voltai*, are likely to be caused by the mixture of individuals originating from different populations. Freeland (2005) suggested that the inclusion of elements of multiple genetic units in a single panel could cause the Wahlund effect, *i.e.*, excess homozygosity and significant estimations of  $F_{IS}$ . Similar results were observed by Apolinário-Silva *et al.* (2018), which

used a panel consisting of 34 individuals derived from genetically distinct units for microsatellite validation.

Microsatellite primers are generally highly species-specific (Zane *et al.*, 2002). However, we have verified that all 14 primers pairs, developed for *Electrophorus varii*, satisfactorily amplify for *E. voltai*. The cross-species amplification implies that it may also be useful in *E. electricus* (which is more closely related to *E. voltai* – see de Santana *et al.*, 2019) as well as in other Gymnotiformes species not tested herein. Heterologous primers can be successfully used in different species of fishes, and the quality of amplification depends on the degree of genetic conservation of positions bordering microsatellite regions (Abdul-Muneer, 2014). Consequently, the low amplification rate primers in the four species of Gymnotiformes can be explained by the lack of conservation of microsatellite sites. Equally, the successful amplification described in *E. voltai* can be attributed to the elevated conservation of the microsatellite flanking regions, which according to Barbará *et al.* (2007), is expected among closely related species. Accordingly, the lowest cross-amplification found for *Gymnotus*, currently hypothesized as the putative sister taxon to *Electrophorus* (Alda *et al.*, 2018), was unexpected (see discussion on *Electrophorus* interrelationships in de Santana *et al.*, 2019), indicating that *Electrophorus* current hypothesis of interrelationships deserves further attention.

## ACKNOWLEDGMENTS

We are grateful to JAR Capiberibe for his financial support to this research (Parliamentary Amendment n° 20470007), IBAMA/SISBIO IBAMA (Instituto Brasileiro do Meio Ambiente e dos Recursos Renováveis), ICMBio (Instituto Chico Mendes-MMA), to INPA for donating tissue samples, and to C. Ruas (UEL), for helping build the library. OAS was granted by the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq 303685/2018–2). This paper was supported by the Project Diversity and Evolution of Gymnotiformes from the São Paulo Science Foundation (FAPESP)/ Smithsonian Institution (# 2016/19075–9) and Global Genome Initiative (GGI–Peer–2017–149) to CDS.

## REFERENCES

- **Abdul-Muneer PM.** Application of microsatellite markers in conservation genetics and fisheries management: Recent advances in population structure analysis and conservation strategies. *Genet Res Int.* 2014;691759. <http://dx.doi.org/10.1155/2014/691759>
- **Alda F, Tagliacollo VA, Bernt MJ, Waltz BT, Ludt WB, Faircloth BC, Alfaro ME, Albert JS, Chakrabarty P.** Resolving deep nodes in an ancient radiation of Neotropical fishes in the presence of conflicting signals from incomplete lineage sorting. *Syst Biol.* 2018; 68(4):573–93. <https://doi.org/10.1093/sysbio/syy085>
- **Allendorf FW, Hohenlohe PA, Luikart G.** Genomics and the future of conservation genetics. *Nat Rev Genet.* 2010; 11:698–709. <https://doi.org/10.1038/nrg2844>
- **Almeida FS, Fungaro MHP, Sodr  LMK.** RAPD and isoenzyme analysis of genetic variability in three allied species of catfish (Siluriformes: Pimelodidae) from the Tibagi River, Brazil. *J Zool.* 2001; 253(1):113–20. <https://doi.org/10.1017/S0952836901000103>

- **Apolinário-Silva C, Ferreira DG, Cavenagh AF, Aprígio NGO, Galindo BA, Carlsson J, Sofia SH.** Development and characterization of fifteen polymorphic microsatellite loci in *Bryconamericus aff. iheringii* (Teleostei: Characidae) and cross-amplification in related Characidae species. *Neotrop Ichthyol.* 2018; 16(1):e170135. <https://doi.org/10.1590/1982-0224-20170135>
- **Barbará T, Palma-Silva C, Paggi GM, Bered F, Fay MF, Lexer C.** Cross-species transfer of nuclear microsatellite markers: potential and limitations. *Mol Ecol.* 2007; 16(18):3759–67. <http://doi.org/10.1111/j.1365-294X.2007.03439.x>
- **Billotte N, Lagoda PJ, Risterucci AM, Baurens FC.** Microsatellite enriched libraries: applied methodology for the development of SSR markers in tropical crops. *Fruits.* 1999; 54(4):277–88.
- **Botstein D, White LR, Skolnick M, Davis RW.** Construction of a genetic linkage map in man using restriction fragment length polymorphisms. *Am J Hum Genet.* 1980; 32(3):314–31.
- **Catania KC.** The astonishing behavior of electric eels. *Front Integr Neurosci.* 2019; 13:23. <http://dx.doi.org/10.3389/fnint.2019.00023>
- **Crampton WGR.** Electrosensory, electrogenesis and signal evolution in freshwater fish. *J Fish Biol.* 2019; 95(1):92–134. <https://doi.org/10.1111/jfb.13922>
- **Edwards KJ, Barker JHA, Daly A, Jones C, Karp A.** Microsatellite libraries enriched for several microsatellite sequences in plants. *Biotechniques.* 1996; 20(5):758–60. <https://doi.org/10.2144/96205bm04>
- **Finger S, Piccolino M.** The shocking history of electric fishes: from ancient epochs to the birth of modern neurophysiology. Oxford: Oxford University Press; 2011.
- **Freeland JR.** *Molecular Ecology.* Chichester: J Wiley; 2005.
- **Gallant JR, Traeger LL, Volkening JD, Moffett H, Po-Hao C, Novina CD et al.** Genomic basis for the convergent evolution of electric organs. *Science.* 2014; 344(6191):1522–25. <https://doi.org/10.1126/science.1254432>
- **Hall TA.** BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symp Ser.* 1999; 41:95–98.
- **Moysés CB, Mockford S, Almeida-Toledo LF, Wright JM.** Nine polymorphic microsatellite loci in the Neotropical electric eel *Eigenmannia* (Teleostei: Gymnotiformes). *Mol Ecol Notes.* 2005; 5(1):7–9. <https://doi.org/10.1111/j.1471-8286.2004.00803.x>
- **Paetkau D, Calvert W, Stirling I, Strobeck C.** Microsatellite analysis of population structure in Canadian polar bears. *Mol Ecol.* 1995; 4(3):347–54. <https://doi.org/10.1111/j.1365-294x.1995.tb00227.x>
- **Rice WR.** Analyzing tables of statistical tests. *Evolution.* 1989; 43(1):223–25. <https://doi.org/10.2307/2409177>
- **Rozen S, Skaletsky HJ.** Primer 3 on the www for general users and for biologist programmers. In: Misener S, Krawetz SS, editors. *Bioinformatics methods and protocols: Methods in molecular biology.* Totowa: Humana Press; 2000. p.365–86.
- **de Santana CD, Crampton WGR, Dillman CB, Frederico RG, Sabaj MH, Covain R et al.** Unexpected species diversity in electric eels with a description of the strongest living bioelectricity generator. *Nat Commun.* 2019; 10:4000. <https://doi.org/10.1038/s41467-019-11690-z>
- **Schuelke M.** An economic method for the fluorescent labeling of PCR fragments. *Nat Biotechnol.* 2000; 18:223–34. <https://doi.org/10.1038/72708>
- **Vallone MV, Butler JM.** AutoDimer: a screening tool for primer-dimer and hairpin structures. *BioTechniques.* 2004; 37(2):226–31. <https://doi.org/10.2144/04372ST03>
- **Weir BS.** *Genetic Data Analysis II: Methods for discrete population genetic data.* Sunderland: Sinauer Associates Inc.; 1996.
- **Zane L, Bargelloni L, Patarnello T.** Strategies for microsatellite isolation: A review. *Mol Ecol.* 2002; 11(1):1–16. <https://doi.org/10.1046/j.0962-1083.2001.01418.x>



## AUTHOR CONTRIBUTION

**Lenice Souza-Shibatta:** Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Project administration, Supervision, Writing–original draft, Writing–review and editing.

**Dhiego G. Ferreira:** Formal analysis, Investigation, Methodology, Writing–original draft, Writing–review and editing.

**Kátia F. Santos:** Formal analysis, Investigation, Methodology.

**Bruno A. Galindo:** Formal analysis, Investigation, Writing–original draft.

**Oscar A. Shibatta:** Investigation, Writing–original draft, Writing–review and editing.

**Sílvia H. Sofia:** Investigation, Methodology, Writing–review and editing.

**Renata M. Giacomín:** Formal analysis, Investigation, Methodology.

**Douglas A. Bastos:** Data curation, Investigation, Resources, Writing–original draft, Writing–review and editing.

**Raimundo N. G. Mendes-Júnior:** Data curation, Investigation, Methodology, Resources, Writing–review and editing.

**Carlos David de Santana:** Conceptualization, Data curation, Funding acquisition, Project administration, Resources, Supervision, Writing–original draft, Writing–review and editing.

## ETHICAL STATEMENT

This study was carried out in strict accordance with the recommendations provided in the Guide for the Care and Use of Laboratory Animals. The collection was authorized by the System of Authorization and Information on Biodiversity – SISBIO (n°. 40522–6). The sampling protocol was approved by the Ethics Committee on the Use of Animals – CEUA of the Instituto Nacional de Pesquisas da Amazônia (n°. 044/2016).

## COMPETING INTERESTS

The authors declare no competing interests.

## HOW TO CITE THIS ARTICLE

- **Souza-Shibatta L, Ferreira DG, Santos KF, Galindo BA, Shibatta OA, Sofia SH, Giacomín RM, Bastos DA, Mendes-Júnior RNG, de Santana CD.** Electric eels galore: microsatellite markers for population studies. *Neotrop Ichthyol.* 2020; 18(4):e200081. <https://doi.org/10.1590/1982-0224-2020-0081>

Neotropical Ichthyology

OPEN ACCESS



This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

Distributed under Creative Commons CC-BY 4.0

© 2020 The Authors. Diversity and Distributions Published by SBI



Official Journal of the Sociedade Brasileira de Ictiologia