

ARTISANAL FISHERIES PRODUCTION IN THE COASTAL ZONE OF ITAIPU, NITERÓI, RJ, BRAZIL*

Rafael de Almeida Tubino^{1**}; Cassiano Monteiro-Neto¹; Luiz Eduardo de S. Moraes² & Eduardo Tavares Paes²

¹Universidade Federal Fluminense, Departamento de Biologia Marinha
Pós-Graduação em Biologia Marinha
(Caixa Postal 100644, 24001-970 Niterói, RJ, Brasil)
**rafaeltubino@bol.com.br

²Instituto Nacional de Pesquisas Espaciais – INPE
Grupo de Sensoriamento Remoto Aplicado à Pesca – Divisão de Sensoriamento Remoto
(Av. dos Astronautas, 1.758, Jd. Granja, 12227-010 São José dos Campos, SP, Brasil)

ABSTRACT

Artisanal fisheries in the coastal zone of Itaipu-RJ play an important role in the local market, but little is known about production and productivity of the fisheries. From April 2001 to March 2003, we monitored a total of 618 landings from the beach-seine (99), the gill nets (“corvineira” - 331, “linguadeira” - 40 and “rede alta” - 25), the hook and line (98), the squid jigging (17), and the spear fishing and trap fisheries (five and three, respectively). The total monitored production was 21866 kg and the catch per unit effort (CPUE) 35.4 kg.trip⁻¹. Considering the number of effective fishing days and the average daily production per fishing gear, the annual production was estimated at 136687 kg (0.2% of the state’s production in 2004). Predominant resources were: *Micropogonias furnieri*, *Trichiurus lepturus*, squids (*Loligo* spp.), clupeiform fishes (*Sardinella brasiliensis*, *Cetengraulis edentulus*, *Pellona harrowerii* and *Harengula clupeiola*), *Eucinostomus* spp., *Cynoscion* spp., *Menticirrhus* spp., *Caranx crysos* and *Selene setapinnis*. The combined analysis using hierarchical and non-hierarchical classification methods, separated four fishing gears and five species groups, based on their similarity patterns, identifying gears that most effectively capture the different fish groups. The beach-seine was less selective and most productive fishery in the summer. The “corvineira” was the second most productive fishery with higher yields in the summer and fall. Catches of “linguadeira” included rays and *Paralichthys* spp., whereas the “rede alta” caught primarily *M. furnieri*, *Mugil* spp. and sharks.

RESUMO

As pescarias artesanais na zona costeira de Itaipu-RJ desempenham um papel importante no mercado local, mas pouco se conhece sobre a produção e a produtividade da pesca. De abril-2001 a março-2003, nós monitoramos um total de 618 desembarques das pescarias de arrastos-de-praia (99), de redes de emalhe (corvineira - 331, linguadeira - 40 e rede alta - 25), de linha-e-anzol (98), de zangareio (17), e de arpão e puçá (cinco e três respectivamente). A produção monitorada total foi de 21866 kg e a captura por unidade de esforço (CPUE) de 35,4 kg.viagem⁻¹. Considerando o número de dias efetivos de pesca e a produção média diária por arte de pesca, a produção anual foi estimada em 136687 kg (0,2% da produção do estado em 2004). Os recursos predominantes foram: *Micropogonias furnieri*, *Trichiurus lepturus*, lulas (*Loligo* spp.), peixes clupeiformes (*Sardinella brasiliensis*, *Cetengraulis edentulus*, *Pellona harrowerii* e *Harengula clupeiola*), *Eucinostomus* spp., *Cynoscion* spp., *Menticirrhus* spp., *Caranx crysos* e *Selene setapinnis*. A análise combinada utilizando métodos de classificação hierárquica e não-hierárquica separou quatro grupos de artes de pesca e cinco de espécies, baseada nos seus padrões de similaridade, identificando as artes de pesca que melhor capturam os diferentes grupos de recursos pesqueiros. O arrasto-de-praia foi a arte de pesca menos seletiva e mais produtiva no verão. A corvineira foi a segunda pescaria mais produtiva com rendimentos maiores no verão e outono. As capturas com linguadeira incluíram raias e *Paralichthys* spp., enquanto que a rede alta capturou principalmente *M. furnieri*, *Mugil* spp. e tubarões.

Descriptors: Artisanal fisheries, Production, CPUE, Coastal zone, Itaipu-RJ, Brazil.

Descritores: Pesca artesanal, Produção, CPUE, Zona costeira, Itaipu-RJ, Brasil.

(*) Paper presented at the 1st Brazilian Congress of Marine Biology, on 15-19 May 2006. Rio de Janeiro, Brazil.

INTRODUCTION

Human demand for marine resources is increasing worldwide, especially in large urban centers of developing countries. Under this scenario, small scale artisanal fisheries play an important role for supporting local and regional markets. Reported small scale marine fisheries account for 40% of the marine fish taken for human food (FAO, 1998) and comprise about 50 million fishers (Berkes *et al.*, 2001). Along the Brazilian coast, the artisanal fishery sector plays an important economic, social and cultural role, responsible for a great part of the overall catches (Diegues, 1999). However, data on this sector are frequently scattered and isolated, thus limiting the possibilities for establishing catch and productivity patterns for supporting management strategies and initiatives.

Due to their dynamics and multispecies catches, artisanal fisheries use several fishing gears and affect different fish stocks. Jablonski & Silva (1996) considered that, for the most part, the fishing effort of artisanal fishers is within sustainable limits and are usually adequate for the exploitation of local resources. On the other hand, several authors (Begossi, 1992, 2006; Nehrer & Begossi, 2000; Kant de Lima & Pereira, 1997) demonstrated that most artisanal fishers have complained about the drastic decline of fish

stocks, mostly due to competition with the industrial fleet, but also as a consequence of coastal environment degradation. Furthermore, competition for space and exclusion from their traditional fishing areas, also limited access to the fish stocks and reduced the artisanal production capability.

The coastal region of Itaipu ($22^{\circ}53'14''\text{S}$, $43^{\circ}22'48''\text{W}$) shelters intensive artisanal fisheries, established in the area since the 18th century, and currently involving about 200 fishers. The area is located to the west of the mouth of Guanabara Bay, and forms a semi-sheltered cove protected by three coastal islands (Fig. 1). The coastal water mass is a mix of Guanabara Bay waters, contributions from the Itaipu-Piratininga lagoons, and coastal oceanic waters, seasonally influenced by weak upwelling of the South Atlantic Central Water, following the general pattern for the South Atlantic Bight (Castro-Filho *et al.*, 1987).

This study aims to identify the fisheries catch composition and provide estimates of the total fishing production and productivity. We further used a combined hierarchical and non-hierarchical classification analysis to evaluate similarities between the species composition and productivity from the different fishing gears used, to define seasonal species groups that constituted the artisanal fishery harvests in the Itaipu coastal zone.




Fig. 1. Location of the fishing areas in the coastal zone of Itaipu (RJ): beach-seine (★); gill nets (□); hook and line and squid jigging (□); trap (●) and spear fishing (▲).

MATERIAL AND METHODS

We monitored the artisanal fisheries landings, one to four times a month, at Itaipu, Niterói-RJ, from April 2001 to March 2003 (618 trips in 77 days), recording data on the fishing fleet, fishing gear, catch composition and abundance for estimating fisheries effort, production, CPUE and seasonality. Sampling effort was determined mostly by the availability of the research crew.

Data regarding the catch composition, the fishing gear used, the area where fishing occurred, and time spent fishing was gathered through interviews with the fishers. Data on fisheries production were obtained from estimates of the total number of boxes landed of a given species, taking that one box weights 22.0 kg of fish. This measure was also checked at times, by weighing randomly selected boxes with a 100 kg scale with 0.5 kg precision. We also weighed individuals of selected species for calibration and length-weight relationships using several precision scale dynamometers.

The fishing areas outlined in Figure 1, were established based in the literature (Kant de Lima & Pereira, 1997; Begossi, 2006), personal observations and fishers interviews in which they were asked to mark, in a nautical chart, fishing grounds for different gears used

Fishing operations and gears used in the fishing area included the beach seining, the gill netting, the hook and line, squid jigging, spear fishing and a trap fisheries. In order to standardize fishing effort we considered boat trips as our effort unit. Beach seine fishing operations were also considered as boat trips. All trips were monitored at the landing site during the sampling day, by teams of two to three observers. Once a month one beach-seine operation was followed throughout in order to record the full fishing operation, fish sorting, and marketing.

To estimate annual fisheries production, separated by fishing gear and by season, we first calculated the average catch (kg) per trip (CPUE) for each gear within a season. We then estimated the average number of trips (T) based on a standard monitoring day in which we were certain that, for each particular gear, all trips were monitored within that day. To estimate the effective fishing days (DF) we subtracted from the total number of days in each season: (a) the number of cold fronts passing through the region, obtained from climate bulletins. We assumed, based on interviews and personal observations, that for each cold front fishers don't go out fishing for three days. (b) Sundays and holidays in which fishing does not occur. We then used the following empirical relationship to calculate

production within a season: $Production = CPUE * T * DF$.

The seasonal groups considered followed approximately the conventional southern hemisphere four seasons definition: Summer – January to March; Fall – April to June; Winter – July to September; and Spring – October to December.

Species were identified according to the specialized literature (Carvalho-Filho, 1999; Figueiredo, 1977; Figueiredo & Menezes, 1978, 1980; Menezes & Figueiredo, 1980; 1985; 2000; Nelson, 1994). Species data were also checked with Fishbase website (www.fishbase.org).

To evaluate the similarity in the catch specific composition among the fishing gears, species catch data from the 610 monitored fishing operations were rearranged on a 22 column (samples – fishing gear by season) by 74 row (species) matrix. Spear fishing and trap fisheries were not considered in this analysis due to their small frequency of occurrence. The squid jigging fishery did not occur in the winter and spring. Only species that presented catches greater than 0.5 kg were included in the matrix. Values within the matrix represent the average catch of a given species, captured by one fishing gear at a certain time of the year (Table 2). Data were log transformed [$\ln(\text{catch} + 1)$] and later standardized using the "Ranging for variables with arbitrary zero" procedure proposed by Milligan & Cooper (1988), defined as:

$$X_p = \frac{X - \text{Min}(X)}{\text{Max}(X) - \text{Min}(X)}, \text{ where:}$$

X_p = standardized value X = original value, $\text{Min}(X)$ = smallest column value, $\text{Max}(X)$ = largest column value. After standardization values within each column vary between 0 and 1. The transformed and standardized matrix was submitted to q-mode (grouping samples - columns) and r-mode (grouping species - rows) classification analyses (Legendre & Legendre, 1998). Q-mode classification was conducted using SHAM (Sequential Hierarchical Agglomerative Method), using the Ward method for generating hierarchical dendrograms. This method uses analysis of variance to define clusters by minimizing the sum of squares and maintaining within group variance as homogeneous as possible (Milligan & Cooper, 1987; Rossi-Wongtschowski & Paes, 1993). The analysis was accomplished using the free access software: FITOPAC1 (Shepherd, 1995).

R-mode classification was accomplished using a non hierarchical K-means algorithm that gathers similar variables in a pre-established number of groups (Legendre & Legendre, 1998). Such groups

are geometrically as compact as possible around their respective centroids. The method is particularly efficient when classifying widely distributed variables with many null values without the need for establishing a relative scale of dissimilarity (Legendre *et al.*, 2002). The best number of groups present in a data set was defined by maximum value of the *C-H* index (Calinski & Harabasz, 1974) for different partitions (number of groups), where:

$$C - H = \left[\frac{R^2 / (K - 1)}{(1 - R) / (n - K)} \right]$$

where, $R^2 = (SST - SSE) / SST$.

SST is the total sum of squared distances to the overall centroid and SSE is the sum of squared distances of the objects to their group's own centroids. The classification was accomplished with the aid of the free software K-MEANS2 (Legendre, 2001), available in: (<http://www.bio.umontreal.ca/casgrain/en/labo/kmeans.html>). Finally, both classifications were jointly analyzed to verify species groups that were strongly associated with particular fishing gear groups. This approach of combining q-mode and r-mode classifications in two-way tables is usually called nodal analysis (Rossi-Wongtschowski & Paes, 1993).

RESULTS

Monitored Landings

During the sampling period we recorded eight types of fisheries operating different fishing gears. A description of the fisheries methods is given by Kant de Lima & Pereira (1997). The beach-seine fishery operates from shore, at several stations distributed over Itaipu and Camboinhas beaches. Three types of gill net fisheries ("corvineira", "linguadeira" and "rede alta") operate within the cove and sometimes behind the islands. The hook and line and the squid jigging are conducted in more sheltered waters, off Itaipu Beach and between the Menina and Mãe Islands. Spear fishing is performed near the rocky coast and around all three islands. Traps ("Puçás") are mostly used in shallow waters at Itaipu Beach (Fig. 1).

Landings from the "corvineira" gill net fishery were the most common (53.6%) among all monitored trips, followed by the beach-seine (16.0%), hook and line (15.9%), the "linguadeira" and "rede alta" (6.5% and 4.0% respectively), squid jigging (2.5%) and spear fishing (0.8%) (Table 1).

A total of 98 species were captured in the monitored trips. Nevertheless most species occurred sporadically at low abundances. The most important species or species categories caught (>1.0% of the total) were: croaker (*Micropogonias furnieri*), largehead hairtail (*Trichiurus lepturus*), squids (*Loligo* spp.), clupeiform fishes (*Sardinella brasiliensis*, *Cetengraulis edentulus*, *Pellona harrowerii* and *Harengula clupeola*), mojarras (*Eucinostomus* spp.), weakfish (*Cynoscion jamaicensis*, *C. leiarchus*, and *C. guatucupa*), kingfish (*Menticirrhus littoralis*, *M. americanus*), banded croaker (*Paralonchurus brasiliensis*), blue runner (*Caranx crysos*), atlantic moonfish (*Selene setapinnis*) and smooth puffer (*Lagocephalus laevigatus*) (Table 1).

The total monitored production was 21.8 tons with an average of 35.4 kg.trip⁻¹. The beach-seine was responsible for more than half of the volume caught (10793 kg) and showed the highest CPUE (109 kg.trip⁻¹). Gear selectivity was low capturing several species and individuals of all sizes. The largehead hairtail was the predominant fish in the beach-seine catch, contributing 18.8% of the total monitored production, followed by squids (12.0%), croaker (11.2%), clupeiform fishes (10.5%) and mojarras (9.8%) (Table 1).

Catches from the "corvineira" fishery yielded a total of 8378 kg, representing approximately 38.0% of the total monitored catch, with a CPUE of 25.3 kg.trip⁻¹. The fishery focuses on demersal fishes, especially croaker, which represented about 82.3% of the total "corvineira" catch (Table 1).

Hook and line was the third most important fisheries monitored in the period. The total catch was equal to 2063 kg and CPUE to 21.0 kg.trip⁻¹. The fishery is highly selective capturing mostly the largehead hairtail (Table 1).

Production from both "linguadeira" and "rede alta" gill nets were low (282 kg and 248 kg respectively). The yield from the "linguadeira" fishery was 7 kg.trip⁻¹ and the catch included flounders (*Paralichthys* spp.) and croaker, whereas for the "rede alta" CPUE was 10 kg.trip⁻¹, and caught mostly demersal-pelagic resources (*M. furnieri*, *Mugil* spp., *Rhizoprionodon* spp. and *Sphyrna lewini*) (Table 1).

The squid jigging fishery is highly seasonal and focused in a single resource. Total monitored production was equal to 48 kg and CPUE to 2.8 kg.trip⁻¹ (Table 1). Nevertheless, squids reach a high market value making them an important fishery resource.

The trap fisheries and the spear fishing, each targeting crabs and octopus respectively, were the least frequent fisheries. Trap monitored production was 34 kg and CPUE 11 kg.trip⁻¹. Spear fishing showed a total catch of 20 kg and a CPUE of 4 kg.trip⁻¹ (Table 1).

Fisheries Cluster Analysis

The Q-mode classification analysis of fisheries resulted in four sample groups reflecting the gear type as the primary factor of dissimilarity among groups. Group 1 was characterized by beach-seine catches and the whole group was well differentiated from the rest, attaining the maximum dissimilarity value. Causes of such differences may be attributed to the low specificity of the fisheries, not targeting any specific resource, and the highest diversity of the catch. Also, the beach-seine showed the highest average catch (Table 2, Fig. 2). Group 2 aggregated the “corvineira” and “linguadeira” gill net fisheries, reflecting similarities between bottom oriented fishing gears, despite their differences in mesh size, and consequent selectivity. Group 3 was formed by the “rede alta” fisheries, which concentrated catches on highly mobile pelagic fishes including sharks, mullets (*Mugil* spp.), and bluefish (*Pomatomus saltatrix*) in the summer. Group 4 clustered together the hook and line and the squid jigging fisheries. The squid jigging is highly selective and oriented exclusively for squids, with no by-catch in the fisheries. Also, the hook and line is quite selective and captures less species than the other fishing gears monitored. Therefore, the association of the squid jigging and hook and line in group 4 probably reflected the large number of absent species or null occurrences shared by both fisheries. The clustering of group 4 could be considered as a classification artifact, yet, it allowed the aggregation of the most selective fisheries, both focused on the capture of pelagic species (Table 2, Fig. 2).

The highest partition of the C-H index in the R-mode classification yielded two groups: one included croaker and largehead hairtail, and the other the remaining species. This classification is trivial and adds very little to the understanding of fisheries patterns. Therefore we considered the second highest partition of the C-H value, which resulted in 5 groups of species or species categories, which still kept the croaker and largehead hairtail as a separate group (Table 2, Fig. 2).

The five groups (A, B, C, D and E) showed distinct species composition and catches associated with the fishing gears used by each one of the fishing methods. For instance, group A included seven species or species categories often captured by the “rede alta” gill net, including sharks *Rhizoprionodon* spp. and *Sphyrna* spp., and bluefish as the most representative. Group B comprised 39 elements, most of them caught exclusively in beach-seines, at catches between 5 - 10 kg.trip⁻¹ (Table 2). Representative species were the clupeiform fishes and rough scad (*Trachurus lathami*). Group C, with 11 species, was less important in beach-seines but predominated in the “linguadeira” catches,

including larger individuals such as sting rays (*Dasyatis* spp.) and angel sharks (*Squatina* spp.). Group D, had 15 species and predominated in beach-seines. Species occurred almost exclusively in this fishing gear and included squids, mojarras, Brazilian sardine (*Sardinella brasiliensis*), blue runner, false herring (*Harengula clupeiola*) and striped weakfish (*Cynoscion guatucupa*). Group E represented the two most abundant and important species, croaker and largehead hairtail, in Itaipu. Both resources were not only abundant and frequent, but were also shared by most of the fisheries conducted in the area such as beach-seine, the hook and line, and nearly all of the gill net fisheries (Table 2, Fig. 2).

Production Estimates

The annual fisheries production, considering 200 effective fishing days was equal to 136687 kg (Table 3). The beach seine fishery accounted for approximately 53.1% of this total, followed by the “corvineira” (27.8%) and the hook and line (13.4%) fisheries. The production estimate of the other fisheries together added up to a total of 7651 kg which represented approximately 5.6% of the total estimated annual production.

Seasonal production estimates revealed a consistent pattern in which summer and fall were the most productive, and winter and spring the least productive periods. Summer alone accounted for 41.9% of the total annual production (Table 3, Fig. 3).

Beach-seine and hook and line showed production peaks in the summer, whereas peak production from the “corvineira” occurred in the fall slightly different from the summer production (Table 3). The production from “rede alta” peaked in the fall and was smallest in the spring, whereas the “linguadeira” peaked in the spring and was the smallest in the winter.

The squid jigging occurs only in the summer and fall following the target species seasonality. The sporadic occurrence and low frequency of the trap fishery and spear fishing, limit the possibilities to infer any seasonal pattern from these fishing modalities (Table 3).

DISCUSSION

According with the classification suggested by Reis (1993), fisheries in the coastal region of Itaipu may be characterized as artisanal due to its small fishing capacity and low technology. Also, fishers play an important role in the marketing process, promoting the auction of the catch at their arrival on the beach.




Fig. 2. Comparison among the grouping of the fishing gears for seasons (mode-Q) using analysis of grouping type SHAM and grouping of species (mode-R) using analysis of grouping type K-means. Fishing gears: BSN – beach-seine, COR – corvineira, LIG – linguadeira, RDA – rede alta, HKL – hook and line and SQJ – squid jigging. Seasons: FA – fall, SU – summer, SP – spring and WI – winter.

Table 3. Estimates of annual and seasonal fisheries production (kg), by fishing gear at Itaipu-RJ, Brazil. T = average number of trips per day; CPUE = kg.trips⁻¹.

Fishing Gear	Seasons		Fall				Winter				Spring				Summer				Annual Production
	T	CPUE	Production		Production		Production		Production		Production		Production						
			Daily	Total	Daily	Total	Daily	Total	Daily	Total	Daily	Total	Daily	Total					
Beach-seine	4	96	384	20349	2	95	190	9143	2	121	243	11176	5	121	603	31966	72633		
Corvineira	8	30	240	12729	8	20	159	7631	7	19	131	6038	6	37	220	11663	38060		
Linguadeira	1	8	8	437	1	6	6	272	3	8	23	1060	2	7	15	772	2541		
Rede alta	1	28	28	1487	1	8	8	404	1	5	5	226	1	12	12	632	2749		
Hook and line	2	18	36	1909	1	16	16	750	4	26	102	4707	8	26	207	10977	18343		
Trap					1	5	5	255					1	14	14	749	1005		
Squid jigging	2	4	8	441									3	2	6	315	756		
Speare fishing					1	4	4	173	1	5	5	207	1	4	4	220	600		
Total			37	705	37352	26	388	18628	36	509	23414	47	1081	57293	136687				
Total Days	91				92				92				90						
Cold Fronts	25				31				33				25						
Sundays, Holidays	13				14				13				12						
Effective Fishing Days	53				48				46				53						




Fig. 3. Seasonal variation of estimated production (kg) of the fisheries in the coastal zone at Itaipu (RJ), from April/01 to March/03.

The classification analysis provided an important tool for identifying species groups associated with the different fishing modalities used in the area on a seasonal basis. Also, the classification suggests a strong association between the different fisheries and their catch. This partition is quite clear as we look at the three different gill net fisheries. The large diversity in the types of fishing gears and specific equipments used towards the exploitation of certain fish resources (e.g. “corvineira” - croaker; “linguadeira” - flounders and angel shark; and “rede alta” – pelagic sharks, mullets and bluefish) result in a multi-specific fisheries showing high overlap regarding the use of the most important fish resources such as the croaker, the largehead hairtail and squids.

The use of several fishing gears and the exploitation of different fishing areas amplify the access to resources, increasing total production. However, Kant de Lima & Pereira (1997) observed that in the early 60’s the introduction of the gill net fisheries raised conflicts with the traditional beach-seiners, which suggested that many of the migrating stocks would not come into the beach-seining area due to gill nets set offshore in neighboring waters. For instance, the mullet beach-seine fishery disappeared from the area in the mid 80’s, and fishers blamed it on the gill nets. Whether this hypothesis is true, still remains to be tested.

The estimated total annual production for the area represents approximately 0.7% of the Guanabara Bay fisheries production, following Jablonski *et al.* (2006), and 0.2% of the Rio de Janeiro total production in 2004 (0.8% from artisanal production) - (IBAMA, 2005). In spite of its small contribution, the fisheries production at Itaipu is locally significant, supplying restaurants and kiosks, but also attracting buyers from the Niterói fish market, when large catches are made or special resources are captured.

The largest yields found during the summer may be related to two major factors: a) the period of more favorable oceanographic conditions for conducting fishing operations, and b) the arrival of many different species near the coast, such as croaker (Vazzoler, 1991; Vazzoler *et al.*, 1999), several species of clupeiform fishes (Saccardo & Rossi-Wongtschowski, 1991), mojarras and squids (Costa & Haimovici, 1990; Perez & Pezzuto, 1998; Perez, 2002). The winter months are significantly less productive. Two possible hypotheses may explain this difference. Unfavorable oceanographic conditions and cold fronts affecting all types of fishing may prevent fishers to go out at sea, two to three days at a time. Such conditions may happen mostly in the winter and spring. Also, artisanal fish marketing in the winter may be less favorable due to cold weather and less buyers on the beach.

In spite of the low yields presented from “linguadeira”, “rede alta”, squid jigging, traps and spear fishing, their catches included flounders, guittarfish, mullets, squids, crabs, octopus and groupers, usually reaching high price in the local market, representing important alternative resources compared, for instance, with the beach-seine catches.

Magro (2005), indicated that, in spite of the lack of information about the artisanal and sport fishing, catches of largehead hairtail with hook and line are considerable in the State of Rio de Janeiro, and peak abundance in the spring and summer seems to be linked to recruitment periods (Bellini, 1980). Still, the harvest of this species appears to be resilient and conservative over time, as Kant de Lima & Pereira (1997) observed in the late 70's, local fishers harvesting similar large catches in January.

The high squid yields in the fall and summer were associated with persistent and abundant spawning groups near the islands. A similar fact had been previously observed near the Arvoredo Island in the State of Santa Catarina (Perez *et al.*, 1997). According to local fishers, the squid harvest occurs from December to February, associated with the occurrence of cold and clear waters near the coast (Lima & Pereira, 1997).

The clupeiform fishes constitute an exclusive fishing resource of the beach-seine. In spite of their low economic value, they represent a resource of interest, for instance, when the Brazilian sardine occurs in great volumes. The production peak in the fall, matches the highest frequency of sardine schools in coastal waters of Rio de Janeiro State, after reproduction (Paiva & Motta, 2000).

ACKNOWLEDGMENTS

We thank the fishers of Itaipu for their collaboration in obtaining the fishing data. Special thanks to Mr. Aurelino Mattos de Souza (Mestre Cambuci) for sharing his ideas and insights about his lifelong experience as a local fisher. Thanks also to everyone at NUFEP, Núcleo Fluminense de Estudos e Pesquisas and Department of Anthropology at UFF. Financial support came from the Ministry of Science and Technology (PADCT-MCT), Fundação Carlos Chagas de Amparo à Pesquisa – FAPERJ. CAPES and CNPq provided fellowships at different levels for the authors and students involved in the project. Finally we thank the students at the Necton Biology and Fisheries Ecology Laboratory, specially our technician Davilma Antonio Borges for her unconditional help on many of the less glamorous tasks of a fisheries lab; Laís de Carvalho Chaves, Tarsila Ferreira Seara and Kathlyn Erin Levasseur who helped in the translation of the manuscript.

REFERENCES

- BEGOSSI, A. The use of optimal foraging theory to understand fishing strategies: a case from Sepetiba Bay (State of Rio de Janeiro, Brazil). **Hum. Ecol.**, v. 20, p. 463-475, 1992.
- BEGOSSI, A. Temporal stability in fishing spots: conservation and co-management in Brazilian artisanal coastal fisheries. **Ecology and Society**, v. 11, n. 1, p. 5., 2006 Disponível em: <http://www.ecologyandsociety.org>. Acesso em: 05 Jun. 2006.
- BERKES, F. *et al.* **Managing small-scale fisheries.** Alternative directions and methods. Ottawa: IDRC, 2001. 320 p.
- CALINSKI, T.; HARABASZ, J. A dendrite method for cluster analysis. **Commun. Stat.**, v. 3, p. 1-27, 1974.
- CARVALHO-FILHO, A. **Peixes:** costa brasileira. São Paulo: Melro, 1999. 320 p.
- CASTRO-FILHO, B. M.; MIRANDA, L. B.; MIYAO, S. Y. Condições hidrobiológicas na plataforma continental ao largo de Ubatuba: variações sazonais e de média escala. **Bolm Inst. Oceanogr.**, S.Paulo, v. 35, n. 2, p. 135-151, 1987.
- COSTA, P. A. S.; HAIMOVICI, M. A pesca de lulas e polvos no litoral do Rio de Janeiro. **Ciência Cult.**, S Paulo, v. 42, n. 12, p. 1124-1130, 1990.
- DIEGUES, A. C. Human populations and coastal wetlands: conservation and management in Brazil. **Ocean coast. Mgmt.**, v.42, p. 187-210, 1999.
- FAO. **Integrated Coastal Area Management and Agriculture, Forestry, and Fisheries.** Rome: Food and Agriculture Organization of the United Nations, 1998. 183 p.
- FIGUEIREDO, J. L. **Manual de peixes marinhos do sudeste do Brasil.** I. Introdução: cações, raias e quimeras. São Paulo: Museu de Zoologia da Universidade de São Paulo, 1977. 104 p.
- FIGUEIREDO, J. L.; MENEZES, N. A. **Manual de peixes marinhos do sudeste do Brasil.** II. Teleostei (1). São Paulo: Museu de Zoologia da Universidade de São Paulo, 1978. 110 p.
- FIGUEIREDO, J. L.; MENEZES, N. A. **Manual de peixes marinhos do sudeste do Brasil.** III. Teleostei (2). São Paulo: Museu de Zoologia da Universidade de São Paulo, 1980. 90 p.
- IBAMA. **Estatística da pesca - Brasil - Grandes regiões e unidades da federação.** Brasília: Ministério do Meio Ambiente - Instituto Brasileiro do Meio Ambiente e dos Recursos Naturais Renováveis, 2005. 138 p. 2004.
- JABLONSKI, S.; SILVA, C. E. M. **Inventário da Pesca no estado do Rio de Janeiro.** Rio de Janeiro: Departamento de Oceanografia – UERJ, 1996. 200 p.
- JABLONSKI, S.; AZEVEDO, A. F.; MOREIRA, L. H. A. Fisheries and conflicts in Guanabara Bay, Rio de Janeiro, Brazil. **Braz Arch. Biol. Tech.**, v. 49, n. 1, p. :79-91, 2006.
- LIMA, R. K. DE; PEREIRA, L. F. **Pescadores de Itaipu:** meio ambiente, conflito e ritual no litoral do Estado do Rio de Janeiro. Niterói, RJ: [s.n.], 1997. 331 p.
- LEGENDRE, P. **Program K-means User's Guide.** Pierre Legendre. Département de sciences biologiques. Université de Montréal, 2001. 12 p. 2001.

- LEGENDRE, P. *et al.* Acoustic seabed classification: improved statistical method. **Can. J. Fish. aquact. Sci.**, v. 59, p. 1085-1089, 2002.
- LEGENDRE, P.; LEGENDRE, L. **Numerical Ecology**. 2. ed. Amsterdam: Elsevier, 1998. 853 p.
- MAGRO, M.; CERGOLE, M. C.; ROSSI-WONGTSHOWSKI, C. L. D. B. **Síntese de conhecimento dos principais recursos pesqueiros costeiros potencialmente exploráveis na costa sudeste-sul do Brasil**: Peixes. Avaliação do potencial sustentável de recursos vivos na zona econômica exclusiva – REVIZEE. Rio de Janeiro: Graflin, 2000. 143 p.
- MENEZES, N. A.; FIGUEIREDO, J. L. **Manual de peixes marinhos do sudeste do Brasil**. IV. Teleostei (3). São Paulo: Museu de Zoologia da Universidade de São Paulo, 1985. 96 p.
- MENEZES, N. A.; FIGUEIREDO, J. L. **Manual de peixes marinhos do sudeste do Brasil**. V. Teleostei (4). São Paulo: Museu de Zoologia da Universidade de São Paulo, 1985. 105 p.
- MENEZES, N. A.; FIGUEIREDO, J. L. **Manual de peixes marinhos do sudeste do Brasil**. V Teleostei (5). São Paulo: Museu de Zoologia da Universidade de São Paulo, 2000. 90 p.
- MILLIGAN, G. W.; COOPER, M. C. A study of standardization of variables in cluster analysis. **J. Classific.**, v. 5, p. 181-204, 1988.
- MILLIGAN, G. W.; COOPER, M. C. Methodological review: clustering methods. **Appl. Psychol. Meas.**, v. 11, p. 329-354, 1987.
- NELSON, J. S. **Fishes of the World**. 3. ed. New Jersey: John Wiley, 1994. 599 p.
- NEHRER, R.; BEGOSSI, A. Fishing at Copacabana (Rio de Janeiro): local strategies in a global city. **Ciencia Cult.**, S Paulo, v. 52, n. 1, p. 26-30, 2000.
- PAIVA, M. P.; MOTTA P. C. S. Cardumes da sardinha-verdadeira, *Sardinella brasiliensis* (Steindachner), em águas costeiras do estado do Rio de Janeiro, Brasil. **Revta bras. Zool.**, v. 17, n 2, p. 339-346, 2000.
- PEREZ, J. A. A. Biomass dynamics of the squid *Loligo plei* and the development of a small-scale seasonal fishery off southern Brazil. **Bull. mar. Sci.**, v. 71, n. 2, p. 633-651, 2002.
- PEREZ, J. A. A. *et al.* A pesca de lulas na Ilha do Arvoredo: Características e relações ecológicas. Verão de 1996. **Notas Técnicas da Facimar**, Itajaí, SC, v. 1, p. 9-21, 1997.
- PEREZ, J. A. A.; PEZZUTO, P. R. Valuable shellfish species in the by-catch of shrimp fishery in southern Brazil: spatial and temporal patterns. **J. Shellfish Res.**, v. 17, n. 1, p. 303-309, 1998.
- REIS, E. G. Classificação das atividades pesqueiras na costa do Rio Grande do Sul e qualidade das estatísticas de desembarque. **Atlântica**, Rio Grande, v. 15, p. 107-114, 1993.
- ROSSI-WONGTSHOWSKI, C. L. D. B.; PAES, E. T. Padrões espaciais e temporais da comunidade de peixes demersais do Litoral norte do estado de São Paulo, Ubatuba, Brasil. **Publ. Esp. Inst. Oceanogr. USP**, n. 10, p. 169-186, 1993.
- SACCARDO, S. A.; ROSSI-WONGTSHOWSKI, C. L. D. B. Biologia e avaliação do estoque da sardinha *Sardinella brasiliensis*: uma compilação. **Atlântica**, Rio Grande. v. 13, n. 1, p. 29-43, 1991.
- SHEPHERD, G. J. **Fitopac 1**. Manual do Usuário. Campinas, SP: Depto. de Botânica, IB, UNICAMP, 1995. 17 p.
- Smith, I. R. A research framework for traditional fisheries. **ICLARM Stud. Rev.**, n. 2, 45 p, 1979.
- VAZZOLER, A. E. A. de M.; SOARES, L. S. H.; CUNNINGHAM, P. T. M. Ictiologia da costa brasileira. In: LOWE-MCCONNELL, R. H. (Ed.). **Estudos Ecológicos de comunidades de peixes tropicais**. São Paulo: Editora Universidade de São Paulo, 1999. p. 424-468.
- VAZZOLER, A. E. A. de M. Síntese do conhecimento sobre a biologia da corvina *Micropogonias furnieri* (Desmarest, 1823), da costa do Brasil. **Atlântica**, Rio Grande, v. 13, n. 1, p. 55-74, 1991.

Sources of Unpublished Material

- BELLINI, A. T. **Biologia e bionomia de *Trichiurus lepturus* (Linneus, 1758) (Trichiuridae; Perciformes; Teleostei), da costa brasileira, entre Cabo Frio (23°00 S) e Torres (29°21 S)**. 1980. 97p. Dissertação (Mestrado) - Instituto Oceanográfico, USP, São Paulo.

(Manuscript received 09 June 2006; revised 23 October 2006; accepted 29 March 2007)