

## CLINICAL SCIENCES

---

### AUTOPSY-PROVEN CAUSES OF DEATH IN LUNGS OF PATIENTS IMMUNOCOMPROMISED BY SECONDARY INTERSTITIAL PNEUMONIA

Alberto Antonio Terrabuio Junior, Edwin Roger Parra, Cecília Farhat, Vera Luiza Capelozzi

---

Terrabuio Junior AA, Parra ER, Farhat C, Capelozzi VL. Autopsy-proven causes of death in lungs of patients immunocompromised by secondary interstitial pneumonia. Clinics. 2007;62(1):69-76.

**PURPOSE:** To present the more frequent associations found in autopsies of immunocompromised patients who developed secondary interstitial pneumonia as well as the risk of death (odds ratio) in having specific secondary interstitial pneumonia according to the cause of immunocompromise.

**METHOD:** From January 1994 to March 2004, 17,000 autopsies were performed at Hospital das Clínicas, São Paulo University Medical School. After examining the pathology report review, we selected 558 of these autopsies (3.28%) from patients aged 15 years or more with primary underlying diseases who developed radiologically diffuse infiltrates of the lung during their hospital course and died after secondary interstitial pneumonia (bronchopneumonia, lobar pneumonia, interstitial pneumonia, diffuse alveolar damage, pulmonary recurrence of underlying disease, drug-induced lung disease, cardiogenic pulmonary edema, or pulmonary embolism). Histology slides were reviewed by experienced pathologists to confirm or not the presence of secondary interstitial pneumonia. Statistical analysis included the Fisher exact test to verify any association between histopathology and the cause of immunocompromise; a logistic regression was used to predict the risk of death for specific histological findings for each of the independent variables in the model.

**RESULTS:** Secondary interstitial pneumonia was histologically represented by diffuse interstitial pneumonitis ranging from mild nonspecific findings (n = 213) to a pattern of diffuse alveolar damage (n = 273). The principal causes of immunocompromise in patients with diffuse alveolar damage were sepsis (136 cases), neoplasia (113 cases), diabetes mellitus (37 cases), and transplantation (48 cases). A high risk of death by pulmonary edema was found for patients with carcinoma of colon. Similarly, in patients with lung cancer or cachexia, a high risk of death by bronchopneumonia (OR = 3.6; OR = 2.6, respectively) was found. Pulmonary thromboembolism was associated with an appreciable risk of death (OR = 2.4) in patients with arterial hypertension. The risk of death was also high in patients presenting hepatic cancer (OR = 2.5) or steroid therapy (OR = 2.4) who developed pulmonary hemorrhage as the histological pattern of secondary interstitial pneumonia. The risk of death by lung metastasis was also elevated (OR = 1.6) for patients that were immunosuppressed after radiotherapy.

**CONCLUSION:** Patients with secondary immunosuppression who developed secondary interstitial pneumonia during treatment in hospital should be evaluated to avoid death by diffuse alveolar damage, pulmonary edema, bronchopneumonia, lung hemorrhage, pulmonary thromboembolism, or lung metastasis. The high-risk patients are those immunosuppressed by hematologic disease; those under steroid treatment; or those with colon or hepatic carcinoma, cachexia, or arterial hypertension.

**KEYWORDS:** Autopsies. Secondary interstitial pneumonia. Immunocompromised patients. Diffuse alveolar damage.

---

Department of Pathology, São Paulo University Medical School - São Paulo/SP, Brazil.

Email: vcapelozzi@lim05.fm.usp.br

Received for publication on September 11, 2006.

Accepted for publication on October 10, 2006.

---

## INTRODUCTION

Acute pulmonary complications in an immunocompromised host are important causes of death in such pa-

tients. Mortality has been reported to be higher than 40% to 50%, especially in patients who develop diffuse infiltrates.<sup>1,2</sup> Pneumonia of bacterial, viral, or fungal origin is the most frequent complication as a consequence of altered immunological status of the host.<sup>3,4</sup> Nevertheless, an immunocompromised host can develop diffuse radiographic lung infiltrates without evidence of active lower respiratory tract infection,<sup>5-7</sup> a condition that has been named secondary interstitial pneumonia. The differential diagnosis of pulmonary disease in these patients includes infectious pneumonia, recurrence of underlying disease, drug-induced lung disease, diffuse alveolar damage, as well as unrelated processes such as cardiogenic pulmonary edema or pulmonary embolism, or any combination of these.<sup>8-10</sup> Prompt investigation and diagnosis are essential for improving the chances of patient survival.<sup>11,12</sup> Clinical and radiographic findings of acute pulmonary complications in the immunocompromised host are nonspecific,<sup>5-7,13</sup> and computed tomography has been shown to be useful in the assessment of these patients,<sup>5</sup> mainly to assist in guiding biopsy. In this context, the complexity of the clinical presentations turns diagnosis by the clinician into a constant challenge. In spite of recent advances, most types of diagnostic support are still expensive. Not infrequently, the clinician initiates treatment, trying to avoid rapid progression of the disease or more invasive procedures. It is thus important to know the main causes of death in this population to establish correct prophylactic actions, which are the cheaper and more intelligent ways to prevent secondary interstitial pneumonia or the eventual indication for lung biopsy.<sup>14</sup>

The aim of this paper is to present the most frequent associations found in autopsies of immunocompromised patients who developed secondary interstitial pneumonia as well as the risk of death (odds ratio) in having specific secondary interstitial pneumonia according to the cause of immunocompromise.

## MATERIALS AND METHODS

**Patients and Autopsies:** From January 1994 to March 2004, 17,000 autopsies were performed at Hospital das Clínicas, São Paulo University School of Medicine. After examining the pathology report review, we selected 558 of these autopsies (3.28%) from patients aged 15 years or more with primary underlying diseases who developed radiologically diffuse infiltrates of the lung during their hospital course and died after secondary interstitial pneumonia (bronchopneumonia, lobar pneumonia, interstitial pneumonia, diffuse alveolar damage, pulmonary recurrence of underlying disease, drug-induced lung disease, cardiogenic pulmonary edema, or pulmonary embolism). The rapid and fatal clinical course was marked by progres-

sive respiratory failure and did not allow a pathological evaluation of lung biopsies or bronchoalveolar lavage specimens. Complete autopsies were performed in all cases. Tissue samples were taken from all organs, including from each lobe of both lungs. They were immediately fixed in 10% phosphate-buffered formaldehyde solution and embedded in paraffin. Serial sections were cut (3 mm), deparaffinized, and tissues were routinely processed. In all cases, staining with hematoxylin and eosin was done for histological examination, silver stains or mucicarmine for *Pneumocystis* and fungi, Ziehl-Neelsen stain for mycobacterium, and Gram stains for bacteria.

**Histopathology:** Histology slides from the autopsies were reviewed by experienced pathologists to confirm the presence or not of secondary interstitial pneumonia, according to the following patterns:

- Acute or chronic interstitial pneumonia, defined as a thickened interstitium infiltrated by neutrophils and macrophages (acute) or lymphocytes and plasmacytes (chronic) in uniform temporal appearance;
- Diffuse alveolar damage, defined as interstitial acute inflammation, with edema and neutrophils in uniform temporal appearance and diffuse distribution; alveolar collapse and hyaline membranes are also present.
- Pulmonary recurrence of underlying disease,
- Pulmonary edema, defined as proteinaceous intra-alveolar transudate
- Pulmonary hemorrhage, defined as alveolar spaces containing red cells
- Pulmonary thromboembolism, defined as arterial or venous occlusion by thrombi

**Statistical Analysis:** First analysis included the verification of association between histopathology and immunocompromise causes and was done using the Fisher exact test. For this purpose, neoplasia cases were stratified according to localization in the pancreas, liver, stomach, esophagus, lung, breast, blood, colon, or nervous system. Transplantation was equally stratified according to the affected organ—in bone marrow, kidney, or liver. We then performed a logistic regression analysis to predict the risk of death for specific histological findings and to estimate the odds ratios for each of the independent variables in the model. We considered results with *P* values of less than 0.05 to be statistically significant.

## RESULTS


Patient gender was 54.3% male and 45.7% female. Patient ages ranged from 15 to 91 years (median 51). The

primary underlying diseases of the 558 patients with diffuse pulmonary infiltrates included cancer (266 cases, 48%), sepsis (266 cases, 48%), transplantation (80 cases, 14%), diabetes mellitus (73 cases, 13%), chemotherapy (46 cases, 8.2%), steroid treatment (40 cases, 7.2%), malnutrition (26 cases, 5%), and radiotherapy (12 cases, 2.2%).

On histological examination, the lungs showed the most evident lesions. Thus, secondary interstitial pneumonia that developed during treatment in the hospital was histologically represented by diffuse alveolar damage, primary or metastatic lung cancer, cardiogenic pulmonary edema, and pulmonary thromboembolism. Two hundred and seventy three cases (49%) presented diffuse alveolar damage, 92 (16%) cardiogenic pulmonary edema, 72 (13%) bronchopneumonia, 62 (11%) pulmonary thromboembolism, 40 (7%) pulmonary hemorrhage, and 17 (4%) cases presented primary or metastatic lung cancer (Figure 1).

Among the patients who developed diffuse alveolar damage, we found viral inclusions in 17 cases, all of which were associated with cytomegalovirus; fungi infection in 11 cases, associated with *Aspergillus* sp (n = 9), *Cryptococcus* sp (n = 1), and *Pneumocystis carinii* (n = 1) (Figure 2). Tuberculosis was found in 9 cases of diffuse alveolar damage. The principal causes of immunocompromise in patients with diffuse alveolar damage were sepsis (136 cases), neoplasia (113 cases), diabetes mellitus (37 cases), and transplantation (48 cases). Chemotherapy and radiotherapy were done in 25 and 8 cases, respectively. Twenty-eight patients received steroid treatment, and 24 patients had cachexia. It is important to emphasize that more than 1 type of immunocompromise might be present in the same patient. After comparing all of these associations, we found

that the highest risk for dying from diffuse alveolar damage was found in patients with hematological neoplasia or under steroid treatment (Table 1). The same was found for pulmonary edema bronchopneumonia, pulmonary thromboembolism, pulmonary hemorrhage, and primary or metastatic lung cancer. A high risk of death by pulmonary edema was found for patients with carcinoma of colon (Table 1).




**Figure 1** - Secondary interstitial pneumonia / diffuse alveolar damage in immunocompromised patients: (A) diffuse alveolar damage, showing a homogeneous distribution of the inflammatory process and hyaline membranes; (B) cardiogenic pulmonary edema, showing the proteinaceous intra-alveolar transudate; (C) diffuse alveolar damage in patients who developed concomitant bronchopneumonia, showing the numerous neutrophils inside of the alveolar space; (D) pulmonary thromboembolism, showing the significant vascular occlusion; (E) pulmonary hemorrhage, showing alveolar spaces containing red cells; and (F) a case with primary lung cancer. Hematoxylin & eosin X 200 (panels A, D), X 100 (panels B, E), and X 50 (panels C, F).

**Table 1** - Results of the logistic regression used to predict the risk of death (odds ratio) in patients with SIP, according to the underlying disease

Variable	B	SE	Wald	Sig	Exp (B)
<b>DAD</b>					
Steroid treatment	1.0229	0.3739	7.4834	0.006	2.7811
Hematological disease	0.6201	0.3014	4.2321	0.03	1.8592
<b>EDEMA</b>					
Colon carcinoma	0.9647	0.5331	3.2755	0.05	2.6241
<b>BCP</b>					
Lung cancer	1.2830	0.5909	4.7139	0.02	3.6075
Caquexia	0.9619	0.4826	3.9724	0.04	2.6167
<b>THROMBOEMBOLISM</b>					
Systemic hypertension	8901	0.4084	4.7516	0.02	2.4355
<b>LUNG HEMORRHAGE</b>					
Hepatic cancer	0.9202	0.5101	3.2540	0.05	2.5098
Steroid therapy	0.8791	0.4025	4.7690	0.02	2.4086
<b>LUNG METASTASIS</b>					
Radiotherapy	0.4934	0.6764	0.5322	0.04	1.6379

SIP = secondary interstitial pneumonia; DAD = diffuse alveolar disease



**Figure 2** - Histological examination of lungs with diffuse alveolar damage show viral inclusions and fungi infection: (A) diffuse alveolar damage, showing notable amounts of viral inclusions; (B) detail of the cytomegalovirus inclusion (arrows); (C) Fungi infection represented by *Aspergillus* sp. (arrows); (D) detail of numerous *Aspergillus* sp. inside of the alveolar space (stained with silver). (E) diffuse alveolar damage, showing the presences of *Cryptococcus* sp. (arrows); (F) detail of this fungi stained with mucicarmine; (G) Diffuse alveolar damage, showing infection by *Pneumocystis carinii* inside the alveolar space; (H) detail of *Pneumocystis carinii* (arrows) (stained with silver). Hematoxylin & eosin X 200 (panels A, E), X 100 (panels C, G), and X 50 (panel B); Silver stain X 100 (panel D), and X 50 (panel H); Mucicarmine stain X 50 (panel F).

Similarly, in patients with lung cancer or cachexia, the risk of death by bronchopneumonia as determined histologically was high (OR = 3.6; OR=2.6, respectively) . Pulmonary thromboembolism, as determined histologically, was associated with an appreciable risk of death (OR = 2.4) in patients with arterial hypertension. Of interest was the high risk of death for patients with hepatic cancer (OR = 2.5) or steroid therapy (OR = 2.4) who developed pulmonary hemorrhage as the histological pattern of secondary interstitial pneumonia . Equally interesting was the elevated risk of death by lung metastasis (OR = 1.6) for patients that were immunosuppressed after radiotherapy.

## DISCUSSION

Conditions that contribute to the requirement for specific histopathological and etiological criteria for definitive diagnosis in patients with immunosuppression are as follows: 1) the nonspecificity of clinical and radiological findings, which are common features for immunosuppressed patients who develop pulmonary involvement during the course of their disease; and 2) co-infections caused by other endemic patho-

gens in Brazil, such as fungi and tuberculosis.<sup>2,6,16-20</sup> The high cost for special techniques, such as molecular methods, still poses a problem in developing countries.

The above-mentioned considerations prompted us to carry out this study using histopathology of autopsy cases. In doing so, we tried to verify the causes of death in the population of immunocompromised patients who developed secondary interstitial pneumonia .

In this series of adult patients (median age, 51 years), men and women had similar prevalences of secondary interstitial pneumonia . We found that the primary underlying diseases of the 558 patients with diffuse pulmonary infiltrates included cancer (48%), sepsis (48%), transplantation (14%), diabetes mellitus (13%), chemotherapy (8.2%), steroid treatment (7.2%), malnutrition (5%) and radiotherapy (2.2%). A similar distribution was found by Juric et al in 3117 autopsies.<sup>21</sup> They found that the most common causes of death were cardiovascular diseases (40.9%), followed by malignancies (25.2%) and infections (12.9%). Among all cardiovascular diseases, myocardial infarction was the most frequently diagnosed (17.9%) but it should be noted that it was misdiagnosed by clinicians in 16.5% of the cases. Incorrectly diagnosed malignancies were found in only 5.7% of the cases; hematological and lymphoid malignancies (48.8%) were the most common neoplasms and were usually confirmed before death. Infections were found in 46.9% of all autopsies. Bacterial pneumonias and peritonitis were overlooked in 67.5% and 23.5% of the cases, respectively, in which they existed together with another serious condition.

On autopsy, we found that in patients with secondary immunosuppression who developed secondary interstitial pneumonia during treatment in hospital, the lungs were the site with the most evident lesions that were directly responsible by death. These patients died from diffuse alveolar damage, primary or metastatic lung cancer, cardiogenic pulmonary edema, or pulmonary thromboembolism.

The most prevalent autopsy finding as cause of secondary interstitial pneumonia in patients with immunosuppression was diffuse alveolar damage, identified in 273 cases. Thus, it is important to emphasize 2 observations made in this study regarding diffuse alveolar damage. The first is the association of this type of damage with patients under mechanical ventilation at the time of death, thus raising the question of whether diffuse alveolar damage could be secondary to mechanical ventilation injury and not related to immunodeficiency. This very important question probably will not be answered by this study because our review was performed on pathologic reports after necropsy, where the information about mechanical ventilation is not present among the data. With respect to the question of whether dif-



fuse alveolar damage could be secondary to mechanical ventilation injury and not related to immunodeficiency, our answer is yes, because most of the patients undergoing mechanical ventilation already have diffuse alveolar damage. In addition, the morphological pattern of diffuse alveolar damage is different when secondary to barotraumas; in the case of barotrauma, this damage is almost always represented by thin hyaline membranes, while in case of immunodeficiency, all of the diffuse alveolar damage cases presented thicker hyaline membranes and had notable changes in alveolar septae as well as extensive alveolar collapse. Brun-Buisson et al<sup>22</sup> studied the epidemiology and outcome of acute lung injury in European intensive care units (ICUs). They found that acute lung injury occurred in 463 (7.1%) of 6522 admissions and 16.1% of all mechanically ventilated patients; 65.4% cases occurred on ICU admission.

The second important observation about diffuse alveolar damage is the association with infectious etiology represented by fungi, citomegalo, and tuberculosis. Among the fungi, *Aspergillus* sp was the most prevalent. Association between *Aspergillus* and diffuse alveolar damage was similarly found in immunosuppressed patients by other authors.<sup>23-29</sup> Another important association was found between diffuse alveolar damage and citomegalo. In our study, 17 immunosuppressed patients presented diffuse alveolar damage and citomegalo viral inclusions, thus confirming the prevalence of citomegalo in immunosuppressed patients, who are usually very sick patients and are kept in the ICU for long periods; it is possible that in these patients, the diffuse alveolar damage is not secondary to the cytomegalic infection.<sup>30-36</sup>

Diffuse alveolar damage was a prevalent morphological finding in patients immunocompromised by sepsis, neoplasia, diabetes mellitus, or transplantation. As expected, all of these conditions are very prevalent in a series of immunosuppressed patients, probably contributing to their deaths. Diffuse alveolar damage was also a final morphological event in 2 immunosuppressed patients who underwent chemotherapy and radiotherapy but were not undergoing that therapeutic regimen at the time of the death. This finding is very important because if they were under thoracic radiotherapy or under the effects of drugs, the interpretation of the pulmonary findings was completely different, as previously reported by Camus and Rossi.<sup>37,38</sup> Also expected was the prevalence of diffuse alveolar damage as the cause of death in patients with cachexia, which was frequently observed in series of patients with immunodeficiency. However, the highest risk of death by diffuse alveolar damage was found in patients with hematological diseases or under steroid treatment, in agreement with previously reported data.<sup>39-46</sup>

In our study, pulmonary edema was the second most important cause of death, with highest risk for patients with carcinoma of colon. This finding is very similar to that found by Hoelz et al in their study concerning morphometric differences in pulmonary lesions in primary and secondary ARDS.<sup>47</sup>

We found that pulmonary thromboembolism as the morphological pattern for secondary interstitial pneumonia occurred in 11% of patients with arterial hypertension, an appreciable risk of death. Madani et al<sup>48</sup> carried out a retrospective review of all postmortem reports during the period 1991 to 2000 at King's College Hospital, London. They found that during that 10-year period, 16,104 deaths occurred, and 6,833 (42.4%) necropsies were performed. The outcome occurrence, fatal pulmonary embolism, was recorded as cause of death in 265 cases (3.9% of all necropsies; 5.2% of adult cases). No deaths from pulmonary embolism occurred in patients under 18 years of age; 80.0% occurred in patients older than 60 years. Of the fatal emboli, 214 of 265 (80.8%) occurred in patients who had not undergone recent surgery. Of these patients, 110 (51.4%) had suffered an acute medical illness in the 6 weeks before death, most often an acute infectious episode (26 cases).

Of interest in our study was the high prevalence of patients presenting hepatic cancer or steroid therapy who developed fatal pulmonary hemorrhage as the histological pattern of secondary interstitial pneumonia. Parambil et al.<sup>49</sup> studied causes and presenting features of pulmonary infarctions in 43 cases identified by surgical lung biopsy. The underlying cause was identifiable in 31 cases (72%) based on a review of clinical, laboratory, radiologic, and histopathologic data. The 2 most common causes were pulmonary thromboembolism (18 cases, 42%) and pulmonary infections (5 cases, 12%). Thromboembolic pulmonary infarctions typically presented as solitary or multiple nodules located in the subpleural regions. Other causes included diffuse alveolar damage in 2 cases (5%), pulmonary torsion in 2 cases (5%), and 1 case each of lung cancer, amyloidosis, embolotherapy, and catheter embolism. In 12 cases (28%), the underlying cause was not directly identifiable but was probably due to previous pulmonary thromboembolism.

We conclude that patients with secondary immunosuppression who develop secondary interstitial pneumonia during treatment in the hospital should be evaluated to avoid death cause by diffuse alveolar damage, pulmonary edema, bronchopneumonia, lung hemorrhage, pulmonary thromboembolism, or lung metastasis. The high-risk patients are those immunosuppressed by hematological disease; who are under steroid treatment; or who have colon or hepatic carcinoma, cachexia, or arterial hypertension.

## ACKNOWLEDGMENTS

This study was supported by the following Brazilian agencies: the Fundação de Amparo a Pesquisa do Estado

de São Paulo [FAPESP 2000/14336-0, 2001/14566-9 and 2003/13007-1]; and the Laboratories for Medical Research [LIM 05], Hospital das Clínicas, University of São Paulo School of Medicine.

## RESUMO

---

Terrabuio Junior AA, Parra ER, Farhat C, Capelozzi VL. Causas de óbito por pneumonia intersticial secundária em autópsias pulmonares de pacientes imunocomprometidos. Clinics. 2007;62(1):69-76.

**OBJETIVO:** Apresentar as associações mais frequentes encontradas em autópsias de pacientes imunossuprimidos que desenvolveram pneumonia intersticial secundária bem como o risco de óbito (Odds Ratio) de desenvolver PIS associada à causa da imunossupressão.

**MÉTODO:** De janeiro de 1994 a março de 2004, 17000 autópsias foram realizadas no Hospital das Clínicas da Faculdade de Medicina da Universidade de São Paulo. A partir da revisão dos laudos patológicos foram selecionados 558 destas autópsias (3,28%) de pacientes com 15 anos de idade ou mais, com alguma doença de base que desenvolveu um infiltrado pulmonar radiologicamente difuso durante o curso da hospitalização e que depois foi para óbito com pneumonia intersticial secundária (broncopneumonia, pneumonia lobar, pneumonia intersticial, dano alveolar difuso, doença pulmonar recorrente, doença pulmonar induzida por drogas, edema pulmonar cardiogênico e embolismo pulmonar). As lâminas histológicas foram revisadas por patologistas experientes para confirmar ou não a presença de pneumonia intersticial secundária. A análise estatística incluiu o "Teste exato de Fisher" para verificar associação entre a histopatologia e causa de imunocomprometimento; e regressão logística para prever o risco de óbito por achados histológicos específicos para cada variável independente do modelo.

**RESULTADOS:** A pneumonia intersticial secundária foi representada histologicamente por pneumonite intersticial

difusa variando de características não específicas leves (n=213) ao padrão histológico de dano alveolar difuso (n=273). A principal causa de imunossupressão nos pacientes com dano alveolar difuso foi sepse (136 casos), neoplasia (113 casos), diabetes melito (37 casos) e transplantados (37 casos). O maior risco de morte por edema pulmonar foi encontrado nos pacientes com carcinoma de cólon. Da mesma forma, nos pacientes com câncer pulmonar ou cachexia ocorreu um alto risco de morte (OR=3.6; OR=2.6, respectivamente) por broncopneumonia. O tromboembolismo pulmonar ofereceu um importante risco de morte (OR=2.4) nos pacientes com hipertensão arterial. Observou-se também risco de morte por câncer hepático (OR=2.5) ou terapia esteroidea (OR=2.4) nos pacientes que desenvolveram hemorragia pulmonar com padrão histológico de pneumonia intersticial secundária. Da mesma forma houve alto risco de morte por metástase pulmonar (OR= 1.6) nos pacientes imunossuprimidos após radioterapia.

**CONCLUSÃO:** Pacientes com imunossupressão secundária que desenvolveram pneumonia intersticial secundária durante o tratamento dentro do hospital podem ser avaliados para evitar como evento final o dano alveolar difuso, o edema pulmonar, a broncopneumonia, a hemorragia pulmonar, o tromboembolismo pulmonar e a metástase pulmonar. Os pacientes com aumento de risco são aqueles imunossuprimidos por doença hematológica, sob tratamento com esteroides, carcinoma hepático, cachexia e hipertensão.

**UNITERMOS:** Autópsias. Pneumonia intersticial secundária. Pacientes imunocomprometidos. Dano alveolar difuso.

## REFERENCES

1. Perez R. Interstitial lung disease: causes, treatment, and prevention. *Ethn Dis.* 2005;15 (2 suppl 2):S45-8.
2. Maimon N, Almog Y. Pulmonary infiltrates in the immunocompromised host. *Isr Med Assoc J.* 2003;5:112-5.
3. Cunha B. Pneumonias in the compromised host. *Infect Dis Clin North Am.* 2001;15:591-612.
4. Chan K, Allen S. Infectious pulmonary complications in lung transplant recipients. *Semin Respir Infect.* 2002;17:291-302.
5. Brown M, Miller R, Muller N. Acute lung disease in the immunocompromised host: CT and pathologic examination findings. *Radiology.* 1994;190:247-54.
6. Hiorns M, Sreaton N, Muller N. Acute lung disease in the immunocompromised host. *Radiol Clin North Am.* 2001;39:1137-51.
7. Gosselin M. Diffuse lung disease in the immunocompromised non-HIV patient. *Semin Roentgenol.* 2002;37:37-53.
8. Crawford S. Noninfectious lung disease in the immunocompromised host. *Respiration.* 1999;66:385-95.
9. Alam S, Chan K. Noninfectious pulmonary complications after organ transplantation. *Curr Opin Pulm Med.* 1996;2:412-18.
10. Conces D. Noninfectious lung disease in immunocompromised patients. *J Thorac Imaging.* 1999;14:9-24.
11. Suh G, Kang E, Chung M, Lee K, Han J, Kitaichi M, et al. Early intervention can improve clinical outcome of acute interstitial pneumonia. *Chest.* 2006;129:753-61.
12. Qefatieh A, Stone C, DiGiovine B, Toews G, Hyzy R. Low hospital mortality in patients with acute interstitial pneumonia. *Chest.* 2003;124:554-9.
13. Roychowdhury M, Pambuccian S, Aslan D, Jessurun J, Rose A, Manival J, et al. Pulmonary complications after bone marrow transplantation: an autopsy study from a large transplantation center. *Arch Pathol Lab Med.* 2005;129:366-71.
14. Lee Y, Wu C, Hsu H, Huang P, Chang Y. Surgical lung biopsy for diffuse pulmonary disease: experience of 196 patients. *J Thorac Cardiovasc Surg.* 2005;129:984-90.
15. Canzian M, Soeiro A de M, Taga MF de L, Barbas CSV, Capelozzi VL. Correlation between surgical lung biopsy and autopsy findings and clinical data in patients with diffuse pulmonary infiltrates and acute respiratory failure. *Clinics.* 2006;61:425-432.
16. Carvalho MGF, Rodrigues MAM, Marques ME, Franco M, Montenegro MR. Lesões do trato gastrointestinal na síndrome da imunodeficiência adquirida: estudo de 45 necrópsias consecutivas. *Rev Soc Bras Med Trop.* 1994;27:135-41.
17. Gutierrez EB, Zanetta DMT, Saldiva PHN, Capelozzi VL. Autopsy-proven determinants of death in HIV-infected patients treated for pulmonary tuberculosis in São Paulo, Brazil. *Pathol Res Pract.* 2002;198:339-46.
18. Lowndes CM, Bastos FI, Giffin KM, Vaz dos Reis AC, d'Orsi E, Alary M. Differential trends in mortality from AIDS in men and women in Brazil (1984-1995). *AIDS.* 2000;14:1269-73.
19. Nash G. Pathologic features of the lung in the immunocompromised host. *Hum Pathol.* 1982;13:841-58.
20. de Medeiros CR, Moreira VA, Pasquini R. Cytomegalovirus as a cause of very late interstitial pneumonia after bone marrow transplantation. *Bone Marrow Transplant.* 2000;26:443-4.
21. Juric G, Tentor D, Jakic-Razumovic J. Autopsy findings and clinical diagnoses: retrospective study of 3,117 autopsies. *Croat Med J.* 1999;40:71-6.
22. Brun-Buisson C, Minelli C, Bertolini G, Brazzi L, Pimentel J, Lewandowski K, et al. ALIVE Study Group. Epidemiology and outcome of acute lung injury in European intensive care units. Results from the ALIVE study. *Intensive Care Med.* 2004;30:51-61. Epub 2003 Oct 16.
23. Hagerty J, Ortiz J, Reich D, Manzarbeitia C. Fungal infections in solid organ transplant patients. *Surg Infect (Larchmt).* 2003;4:263-71. Review.
24. Fortun J, Martin-Davila P, Moreno S, De Vicente E, Nuno J, Candelas A, et al. Risk factors for invasive aspergillosis in liver transplant recipients. *Liver Transpl.* 2002;8:1065-70.
25. Marik PE. Fungal infections in solid organ transplantation. *Expert Opin Pharmacother.* 2006;7:297-305.
26. Kubak BM. Fungal infection in lung transplantation. *Transpl Infect Dis.* 2002; 4 Suppl 3:24-31. Review.
27. Chang GC, Chang KM, Wu CL, Chiang CD. Clinical patterns among invasive pulmonary aspergillosis patients with and without recent intensive immunosuppressive therapy. *J Formos Med Assoc.* 2001;100:762-6.
28. Iida H, Ri M, Niimi K, Harada T, Sao H. Histopathological findings of pulmonary complications in 16 patients who received allogeneic hematopoietic stem cell transplants. *Rinsho Ketsueki.* 2005;46:1273-8.
29. Chamilos G, Luna M, Lewis R, Bodey G, Chemaly R, Tarrand J, et al. Invasive fungal infections in patients with hematologic malignancies in a tertiary care cancer center: an autopsy study over a 15-year period (1989-2003). *Haematologica.* 2006; 91:986-9. Epub 2006 Jun 1.
30. Humbert M, Lombard-Roux P, Cerrina J. Soluble TNF receptors (TNF-sR55 and TNF-sR75) in lung allograft recipients displaying cytomegalovirus pneumonitis. *Am J Respir Crit Care Med.* 1994;149:1681-5.
31. Barry SM, Johnson MA, Janossy G. Cytopathology or immunopathology? The puzzle of cytomegalovirus pneumonitis revisited. *Bone Marrow Transplant.* 2000;26:591-7.
32. Stals F, Steinhoff G, Wagenaar S, van Breda Vriesman J, Haverich A, Dormans P, et al. Cytomegalovirus induces interstitial lung disease in allogeneic bone marrow transplant recipient rats independent of acute graft-versus-host response. *Lab Invest.* 1996;74:343-52.
33. Oldakowska-Jedynak U, Niewczas M, Ziolkowski J, Mucha K, Foronczewicz B, Bartłomiejczyk I, et al. Cytomegalovirus infection as a common complication following liver transplantation. *Transplant Proc.* 2003;35:2295-7.
34. Magro C, Ali N, Williams JD, Allen J, Ross P Jr. Cytomegalovirus-associated pulmonary septal capillary injury sine inclusion body change: a distinctive cause of occult or macroscopic pulmonary hemorrhage in the immunocompetent host. *Appl Immunohistochem Mol Morphol.* 2005;13:268-72.

35. Dworniczak S, Ziara D, Basta L, Piwowarska G, Jastrzebski D, Oklek K, et al. Human cytomegalovirus serological status in patients with interstitial lung diseases. *Przegl Epidemiol.* 2003;57:431-7.
36. Delbos V, Abgueguen P, Chenebault JM, Fanello S, Pichard E. Acute cytomegalovirus infection and venous thrombosis: Role of antiphospholipid antibodies. *J Infect.* 2006 May 12; [Epub ahead of print].
37. Camus P, Costabel U. Drug-induced respiratory disease in patients with hematological diseases. *Semin Respir Crit Care Med.* 2005;26:458-81. Review.
38. Rossi SE, Erasmus JJ, McAdams HP, Sporn TA, Goodman PC. Pulmonary drug toxicity: radiologic and pathologic manifestations. *Radiographics.* 2000;20:1245-59. Review.
39. Jandrljic M, Kalenic S, Labar B, Nemet D, Jakic-Razumovic J, Mrcic M, et al. An autopsy study of systemic fungal infections in patients with hematologic malignancies. *Eur J Clin Microbiol Infect Dis.* 1995;14:768-74.
40. Gentile G, Micozzi A, Girmenia C, Iori AP, Donati PP, Capria S, et al. Pneumonia in allogeneic and autologous bone marrow recipients. *Chest.* 1993;104:371-5.
41. Stoblen F, Blumhard G, Kaisers U, Raakow R, Bechstein WO, Neuhaus P. Outcome of severe pneumonias after orthotopic liver transplantation. *Transplant Proc.* 1994;26:3681-2.
42. Wade JJ, Rolando N, Hayllar K, Philpott-Howard J, Casewell MW, Williams R. Bacterial and fungal infections after liver transplantation: An analysis of 284 patients. *Hepatology.* 1995;21:1328-36.
43. Sharma S, Nadrous HF, Peters SG, Tefferi A, Litzow MR, Aubry MC, et al. Pulmonary complications in adult blood and marrow transplant recipients: autopsy findings. *Chest.* 2005;128:1385-92.
44. Li GS, Ye QF, Xia SS, Chen ZS, Zeng FJ, Lin ZB, et al. Acute respiratory distress syndrome after liver transplantation: etiology, prevention and management. *Hepatobiliary Pancreat Dis Int.* 2002;1:330-4.
45. Nakase K, Tsuji K, Nagaya S, Tamaki S, Tanigawa M, Ikeda T, et al. Acute interstitial pneumonitis during chemotherapy for haematological malignancy. *Eur J Cancer Care (Engl).* 2005;14:336-41.
46. Walter EA, Bowden RA. Infection in the bone marrow transplant recipient. *Infect Dis Clin North Am.* 1995;9(4):823-47. Review.
47. Hoelz C, Negri EM, Lichtenfels AJ, Concecao GM, Barbas CS, Saldiva PH, et al. Morphometric differences in pulmonary lesions in primary and secondary ARDS. A preliminary study in autopsies. *Pathol Res Pract.* 2001;197:521-30.
48. Madani MM, Jamieson SW. Chronic thromboembolic pulmonary hypertension. *Curr Treat Options Cardiovasc Med.* 2000;2:141-8.
49. Parambil JG, Savci CD, Tazelaar HD, Ryu JH. Causes and presenting features of pulmonary infarctions in 43 cases identified by surgical lung biopsy. *Chest.* 2005;127:1178-83.