

CLINICAL SCIENCE

Metformin, but not glimepiride, improves carotid artery diameter and blood flow in patients with type 2 diabetes mellitus

Helena Atroch Machado,¹ Marcelo Vieira,¹ Maria Rosaria Cunha,¹ Marcia Regina Soares Correia,¹ Rosa Tsunehiro Fukui,² Rosa Ferreira dos Santos,¹ Dalva Marreiro Rocha,¹ Bernardo Leo Wajchenberg,² Silvia G. Lage,² Maria Elizabeth Rossi da Silva¹

¹Hospital das Clínicas da Faculdade de Medicina da Universidade de São Paulo, Laboratory of Medical Investigation (LIM-18), São Paulo/SP, Brazil.

²Hospital das Clínicas da Faculdade de Medicina da Universidade de São Paulo, Heart Institute (InCor) São Paulo/SP, Brazil.

OBJECTIVE: To compare the effects of glimepiride and metformin on vascular reactivity, hemostatic factors and glucose and lipid profiles in patients with type 2 diabetes.

METHODS: A prospective study was performed in 16 uncontrolled patients with diabetes previously treated with dietary intervention. The participants were randomized into metformin or glimepiride therapy groups. After four months, the patients were crossed over with no washout period to the alternative treatment for an additional four-month period on similar dosage schedules. The following variables were assessed before and after four months of each treatment: 1) fasting glycemia, insulin, catecholamines, lipid profiles and HbA_{1c} levels; 2) t-PA and PAI-1 (antigen and activity), platelet aggregation and fibrinogen and plasminogen levels; and 3) the flow indices of the carotid and brachial arteries. In addition, at the end of each period, a 12-hour metabolic profile was obtained after fasting and every 2 hours thereafter.

RESULTS: Both therapies resulted in similar decreases in fasting glucose, triglyceride and norepinephrine levels, and they increased the fibrinolytic factor plasminogen but decreased t-PA activity. Metformin caused lower insulin and pro-insulin levels and higher glucagon levels and increased systolic carotid diameter and blood flow. Neither metformin nor glimepiride affected endothelial-dependent or endothelial-independent vasodilation of the brachial artery.

CONCLUSIONS: Glimepiride and metformin were effective in improving glucose and lipid profiles and norepinephrine levels. Metformin afforded more protection against macrovascular diabetes complications, increased systolic carotid artery diameter and total and systolic blood flow, and decreased insulin levels. As both therapies increased plasminogen levels but reduced t-PA activity, a coagulation process was likely still ongoing.

KEYWORDS: Treatment; Metabolic profile; Vascular reactivity; Hemostatic factors; Type 2 diabetes.

Machado HA, Vieira M, Cunha MR, Correia MRS, Fukui RT, Santos RF et al. Metformin, but not glimepiride, improves carotid artery diameter and blood flow in patients with type 2 diabetes mellitus. Clinics. 2012;67(7):711-717.

Received for publication on November 28, 2011; First review completed on December 28, 2011; Accepted for publication on March 5, 2012

E-mail: maria_cunha@uol.com.br

Tel.: 55 11 3061 7259

INTRODUCTION

Cardiovascular disease is the leading cause of mortality in patients with type 2 diabetes mellitus (DM2). The effective control of glycemia can delay but not prevent vascular complications, which are likely related to many other poorly controlled atherogenic factors, such as hyperlipidemia,

hypertension, oxidative stress, accelerated aging, hyperinsulinemia, disturbances in coagulation and fibrinolysis (1).

Biguanides and sulfonylureas remain the principal oral therapeutic options for treating patients with DM2 (2). The biguanide metformin has been established as a first-line drug for the management of type 2 diabetes. Its indications are supported by its potency, lack of weight gain, low risk of hypoglycemia and mode of action in countering insulin resistance (3). The drug's anti-atherosclerotic and cardio-protective effects appear to reflect a combination of glucose-independent effects on the vascular endothelium, suppressant effects on glycation, oxidative stress and the formation of adhesion molecules, and anti-inflammatory properties, in addition to stimulating fibrinolysis and favorable effects on lipid profiles (4).

Copyright © 2012 CLINICS – This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

No potential conflict of interest was reported.

In contrast, sulfonylureas enhance insulin release primarily by closing the ATP-sensitive K⁺ channels of pancreatic β-cells. The glucose-lowering potency of sulfonylureas is similar to that of metformin. However, by stimulating insulin secretion, sulfonylureas are believed to favor the development of hypoglycemia and weight gain, accelerate beta-cell apoptosis and beta-cell exhaustion and impair endothelial function, thereby increasing the risk for ischemic complications. The blocking of potassium channels in the heart by sulfonylureas has raised concern regarding the drugs' potential adverse effects in cases of ischemic heart episodes (5).

Glimepiride is a long-acting and low potency insulin secretagogue sulfonylurea that has an insulin-sensitizer effect on the muscles and liver (6) and rarely causes hypoglycemia (5). Data from studies in humans are scarce but have suggested that the risks of developing coronary artery disease and mortality do not appear to be increased by glimepiride (7-8). In addition, no deleterious effects of glimepiride on brachial vasodilatation — an acute effect (9) — or ischemic preconditioning — acute (10) and chronic (11) effects — have been observed

Only two studies have investigated the chronic vascular effects of glimepiride measured by forearm arterial blood flow, and the results were similar to those of glibenclamide (12-13) and metformin (12). There have been no reports comparing carotid artery blood flow during glimepiride and metformin therapy, and only a few have measured hemostatic factors (14-19), specifically plasminogen activator inhibitor (PAI-1), fibrinogen and adhesion molecules. In our study, the effects of glimepiride in patients with type 2 diabetes on vascular reactivity, hemostatic factors, and fat and carbohydrate metabolism were compared with those of metformin, the first-line therapy. This comparison is valuable in that these different classes of oral anti-diabetic agents — metformin and glimepiride — target the two main pathophysiological defects of type 2 diabetes, and their safety is fundamental.

METHODS

Patients

A prospective study was performed in 16 uncontrolled patients with type 2 diabetes according to the ADA criteria (2); the cohort included ten women and six men with a mean age of 51.8 ± 6.5 years (mean ± SD) previously treated with dietary intervention. Subjects who had fasting blood glucose values >7.78 mmol/L and/or glycated hemoglobin exceeding the normal range (4-8.5%) by 1.0% or more after two or more months of a diet therapy program without medications (basal values) were included. The participants were randomly assigned to receive either metformin (M group) or glimepiride (G group). The drug dosage was titrated to achieve fasting glucose levels lower than 7.0 mmol/L using domiciliary capillary glucose measurements. After four months, the patients were crossed over with no washout period to the alternative treatment for an additional four-month period on a similar dosage schedule. The subjects were followed on an outpatient basis every 1-2 weeks throughout the study period for drug and weight-maintaining diet adjustments. The clinical characteristics of the patients are depicted in Table 1. Three of the sixteen subjects smoked, and six had systemic arterial hypertension, which was treated with converting-enzyme inhibitors.

Table 1 - Anthropometric data.

	Pre-treatment	Metformin Group	Glimepiride Group	
Age (yr)	51.8 ± 6.5			
Sex (F:M)	10:06			
Dose (mg/day)		1907 ± 558	4 ± 2	
Weight (kg)	71.1 ± 13.4	68.9 ± 12.5	70.5 ± 14.2	Ns
BMI (kg/m ²)	27.6 ± 3.6	26.7 ± 3.4	27.2 ± 3.7	Ns
W/H	0.9 ± 0.1	0.9 ± 0.1	0.9 ± 0.1	Ns
DBP (mm Hg)	83.3 ± 9.3	79.4 ± 7.5	81.1 ± 8.5	Ns
SBP (mm Hg)	141.5 ± 14.0	129.2 ± 16.4	134.0 ± 14.0	Ns

Values are expressed as means ± SDs. DBP = diastolic blood pressure; SBP = systolic blood pressure; BMI = body mass index; W/H = waist-to-hip ratio.

Seven of the ten women were postmenopausal, and none had received hormone replacement therapy, while the premenopausal women were tested up to the 8th day of the follicular phase of their cycles. At the time of enrollment, a complete medical history, physical examination, and laboratory evaluation, including urinalysis, renal, hepatic, and thyroid function tests and serum lipid and electrolyte levels, were obtained for all of the subjects. An ECG and echocardiogram were performed, and subjects with left ventricular systolic dysfunction, valve abnormalities, arrhythmias and ischemic heart disease were not enrolled. None of the patients exercised on a regular basis. Other exclusion criteria included any severe concomitant illness, nephropathy (serum creatinine >1.6 mg/dL or microalbuminuria), uncontrolled hypertension (BP >190x120 mm Hg), stroke, peripheral vascular disease, marked dyslipidemia (total cholesterol >6.5 mmol/L and triglyceride levels >2.8 mmol/L), coagulopathy, proliferative diabetic retinopathy and use of hypolipidemic and anticoagulant medications. None of the subjects demonstrated clinical evidence of autonomic neuropathy as assessed by blood pressure response to standing, beat-to-beat heart rate variation, the Valsalva maneuver and the handgrip test. The Medical Ethics Committee of Hospital das Clínicas and Heart Institute (INCOR) approved the study protocol, and all of the subjects provided written informed consent.

Study protocol

The patients were instructed to continue a similar food intake and abstain from the use of tobacco, alcohol, coffee, salty food, and any physical activity for 24 h before test days and discontinue converting-enzyme inhibitors 72 h before the evaluation.

The following procedures were performed before (basal values) and after each four-month treatment period (M and G groups): 1) hormonal and metabolic evaluations: fasting plasma glucose, insulin and catecholamine levels, lipid profiles and HbA_{1c}; 2) hemostatic factor determination: tissue plasminogen activator (t-PA) antigen and activity, plasminogen activator inhibitor (PAI-1) antigen and activity, platelet aggregability, fibrinogen and plasminogen levels; and 3) a cardiovascular evaluation: by high-resolution ultrasound of the carotid and brachial arteries.

At the end of each treatment period, a 12-h metabolic profile, including measurements of glucose, insulin, glucagon, proinsulin, and triglyceride levels at fasting and every 2 h (from 7 am to 7 pm), was obtained. The meals offered to patients (breakfast, lunch, and dinner) contained 50% of

their total calories as carbohydrates, 20% as protein, and 30% as fat, and the meals were provided by the Metabolic Unit of the Hospital das Clínicas.

Cardiovascular evaluation

The study was performed in a laboratory setting at a temperature of 23°C and in low luminosity, with the patients in the supine position. A catheter was inserted into the forearm vein 30 minutes before blood sample collection for catecholamine, insulin, sodium, and potassium determinations and maintained with a saline infusion.

During the study, the heart rate and systolic, diastolic, and mean blood pressure were registered using a non-invasive, automatic oscillometric device (Dinamap 1486, Critikon, Inc., Tampa, FL, USA). The images for flow velocity and diameters of the arteries were first established after a 10-minute equilibration period using a transducer (Apogee-800 Plus, ATL Inc., Bothell, WA, USA). Intima-media thickness (IMT), the compliance and distensibility of the carotid artery, and the total and systolic blood flow indices (TFI and SFI) were obtained. Flow-mediated vasodilation of the brachial artery was measured using a method reported elsewhere (20). A cuff placed on the left forearm was inflated to 200 mm Hg for 5 minutes. Imaging of the artery was performed before cuff inflation (baseline; B) and at 60 seconds after cuff deflation (reactive hyperemia; RH) to obtain measurements of flow velocity and the diameter of the artery. Fifteen minutes after acquisition of the post-occlusion image, the baseline image was reobtained (rebase; RE), after which nitroglycerin was administered sublingually; 3 and 5 minutes later, another image was acquired (N3 and N5, respectively) to establish endothelium-independent vasodilation. The total and systolic flow indices (TFI and SFI) of the brachial artery were also calculated based on these images.

Biochemical and hormonal analyses

Glucose was determined by the glucose oxidase method (Labtest, São Paulo, Brazil) (21), and HbA_{1c} (normal values: 4 to 8.5%) was determined by ionic chromatography (Labtest, São Paulo, Brazil) (22). Total cholesterol was measured by employing the cholesterol oxidase/peroxidase method; HDL cholesterol was separated using the phosphotungstic acid/Mg²⁺ method and measured using the oxidase/peroxidase method, while triglycerides were measured by the lipase/glycerol kinase method (Labtest, São Paulo, Brazil) (23). LDL was estimated using the Friedewald equation (LDL cholesterol = total cholesterol minus HDL cholesterol minus 0.2 × triglycerides). The intra-assay and inter-assay coefficients of variation (CVs) for the glucose and lipid determinations were <3% and 0%, respectively. Insulin, proinsulin and glucagon were quantified by a double-antibody radioimmunoassay (Linco Research, St. Louis, MO, USA) (24). Catecholamines were measured by high-performance liquid chromatography (25). The intra-assay and inter-assay CVs for the hormonal analyses were 6.8% and 9.6% for insulin, 4.4% and 6.5% for glucagon, 5.5% and 6.8% for catecholamines, and 5% and 5.3% for proinsulin, respectively.

Hemostatic factors were measured using the same assay. Plasminogen was measured by chromogenic assay (Plasminogen Accucolor™ Sigma Diagnostic, ST Louis, MO, USA), and the intra-assay coefficient of variation was <3.0%. Fibrinogen was determined by the CLAUSS method (26) with Fibriquick Assay (Sigma Diagnostics, ST Louis, USA). The

intra-assay CV was 8%. Platelet aggregation was performed using a method described by Born (27). The activities of tissue plasminogen activator (t-PA) and plasminogen activator inhibitor (PAI-1) were determined by quantitative assays (Chromolize™ t-PA and Chromolize™ PAI-1, respectively, Biopool, Umea, Sweden); the intra-assay CVs were 3.9% for t-PA and 3.7% for PAI-1. PAI-1 and t-PA antigens were determined by Imulyse and Tint-Elize (Biopool, Umea, Sweden), respectively. The intra-assay CVs were 5% for PAI-1 and 5.5% for t-PA. All of the analyses were performed in duplicate.

Statistical methods

Numerical data are reported as means and standard deviations, and nominal data are reported as proportions. Differences (95% CI) between the treatment groups were initially tested for treatment-time interaction (28) and then compared by an analysis of variance for repeated measurements and a Tukey's post-test or Student's two-tailed test, with $p < 0.05$ considered statistically significant.

RESULTS

A significant treatment period interaction effect was demonstrated for triglycerides, VLDL cholesterol, plasminogen and norepinephrine levels; hence, only the values from the first treatment period were analyzed.

Anthropometric, biochemical, hormonal and hemostatic factor measurements

Weight, waist-to-hip ratio, and systolic and diastolic blood pressure did not change after glimepiride or metformin treatment (Table 1). Fasting plasma HbA_{1c} ($p = 0.000009$) and glucose levels ($p = 0.00009$) decreased by equal amounts in both treatment groups (Table 2). Fasting insulin levels were greater in the G group ($p = 0.009$) compared with the M group. Similar decreases in VLDL cholesterol ($p = 0.007$), triglycerides ($p = 0.023$) and norepinephrine ($p = 0.042$) levels were obtained in both groups. There was an increase in plasminogen levels ($p = 0.025$) after the initial four months of M ($118.2 \pm 8.2 \times 142.4 \pm 32.0$) or G ($128.4 \pm 8.6 \times 130.2 \pm 8.1$) therapy, which although persistent did not remain significant when the groups switched to the alternative drugs. LDL and HDL cholesterol and epinephrine levels were unchanged. Both therapies decreased t-PA activity ($p = 0.024$). There was no significant effect of either of the therapies on the other hemostatic factors measured (PAI-1 antigen and activity, fibrinogen levels and platelet aggregation — data not shown).

12-h metabolic profile

Only the areas under the curve during glimepiride and metformin therapy were analyzed (Figure 1). The 12-h integrated areas for glucose (M: 87.7 ± 11.03 vs. G: 104.61 ± 36.63 mmol/L/h) and triglycerides (M: 20.47 ± 9.68 vs. G: 22.4 ± 11.03 mmol/L/h) were similar for both therapies.

Treatment with metformin was associated with higher glucagon (M: 1361.69 ± 473.25 vs. G: 1044.22 ± 326.90 ng/L/h; $p = 0.0046$) and lower insulin-integrated (M: 1076.61 ± 389.02 vs. G: 1718.69 ± 837.03 pmol/L/h — $p = 0.02$) and proinsulin-integrated (M: 565.38 ± 279.11 vs. G: 834.71 ± 299.96 pmol/L/h — $p = 0.0016$) areas compared with the G group (Figure 1). The proinsulin-to-insulin molar ratios during the M (0.58 ± 0.32) and G (0.60 ± 0.37) therapies did not differ.

Table 2 - Glucose, glycated hemoglobin, insulin, total and fractional cholesterol, triglycerides and catecholamine levels.

	Pre-treatment	Metformin Group	Glimepiride Group	p-value
Glucose (mmol/L)	15.1 ± 5.12	7.10 ± 1.34	8.3 ± 1.7	^a 0.00011 ^b 0.037
HbA _{1c} (%)	10.8 ± 2.3	8.2 ± 1.4	7.8 ± 1.3	^a 0.009 ^b 0.001
Insulin (pmol/L)	59.5 ± 22.9	55.9 ± 29.4	79.6 ± 25.8	^c 0.009
Total cholesterol (mmol/L)	5.41 ± 1.29	5.05 ± 0.93	5.23 ± 1.09	Ns
HDL cholesterol (mmol/L)	0.99 ± 0.21	1.01 ± 0.18	0.95 ± 0.22	Ns
LDL cholesterol (mmol/L)	3.45 ± 1.16	3.32 ± 0.80	3.64 ± 0.90	Ns
VLDL cholesterol (mmol/L)	0.92 ± 0.54	0.66 ± 0.31	0.70 ± 0.25	^d 0.007
Triglyceride (mmol/L)	2.01 ± 1.19	1.38 ± 0.67	1.46 ± 0.60	^d 0.023
Norepinephrine (nmol/dL)	1481.0 ± 986.4	913.3 ± 497.5	958.7 ± 414.7	^d 0.042
Epinephrine (nmol/dL)	104.7 ± 176.8	26.1 ± 72	16.3 ± 49.6	Ns
t-PA activity (IU/mL)	1.1 ± 0.5	0.7 ± 0.3	0.8 ± 0.3	^d 0.024

^a(pre-treatment vs. metformin); ^b(pre-treatment vs. glimepiride); ^c(metformin vs. glimepiride); ^d(pre. vs. post-treatment). Values are expressed as means ± SDs.




Figure 1 - 12-h metabolic profiles of glucose, triglyceride, insulin, proinsulin and glucagon levels. Mean levels are indicated.

Table 3 - Total (TFI) and systolic (SFI) flow indices and percentage changes in the systolic diameter of the carotid artery.

	Pre-treatment	Metformin Group	Glimepiride Group	p-value
TFI (L/min/m ²)	0.319 ± 0.079	0.382 ± 0.094	0.285 ± 0.091	^a 0.024; ^c 0.003
SFI (L/min/m ²)	0.599 ± 0.155	0.738 ± 0.167	0.563 ± 0.152	^a 0.006
% Systolic Diameter		2.75 ± 9.93	-3.50 ± 11.46	^c 0.003 <0.05

^a(pre-treatment vs. metformin); ^c(metformin vs. glimepiride). Values are expressed as means ± SDs.

Vascular reactivity

The elastic arterial properties were evaluated.

Carotid artery (Table 3): Both the total ($p=0.004$) and systolic ($p=0.002$) carotid flow indices increased with metformin therapy compared with baseline values and glimepiride therapy. Moreover, the carotid systolic diameter changed in opposite directions during the therapies; its percentage change (from basal levels) increased with metformin and decreased with glimepiride, and these different behaviors reached significance ($p=0.028$). The compliance, distensibility and intima-media thickness of the carotid artery did not change significantly with any therapy (data not shown).

Brachial artery: There was a similar increase in flow-mediated vasodilatation of the brachial artery after reactive hyperemia (endothelium-dependent) and sublingual nitroglycerin (endothelium-independent) stimuli across the groups. There were no significant effects of medications on systolic diameter or the total and systolic flow indices of the brachial artery (data not shown).

DISCUSSION

This study compared the actions of two different classes of drugs (the biguanide metformin, an insulin sensitizer, and the sulfonylurea glimepiride, an insulin secretagogue) on carbohydrate and lipid metabolism, hemostatic factors and vascular reactivity. We evaluated the same patients with type 2 diabetes before and after four months of treatment with metformin or glimepiride. The aim of this strategy was to minimize the influence of metabolic control on specific drug effects other than glucose control. The data in the literature remain poor and contradictory regarding the direct actions of these drugs on the cardiovascular risk-related factors analyzed in the present study (15-19,30-33).

Carbohydrate and lipid metabolism

Both treatment groups achieved similar and significant mean decreases from baseline in fasting plasma glucose and HbA1 levels. The lower insulin and proinsulin levels (at fasting and during the 12-h metabolic profile) observed during metformin therapy compared with glimepiride were in agreement with reports of metformin's sparing effects on beta cell function, thus lowering the basal and postprandial insulin requirements for the same metabolic control (3,17). These differences in insulin secretion could not be accounted for by changes in body weight, which were unaffected in both groups. Although some authors have found no increases in insulin levels (18) and others have reported that glimepiride's insulin trophic effect might diminish in the presence of normoglycemia (34), the insulin levels in this study were found to be higher during therapy with glimepiride than metformin.

Despite these findings, the proinsulin-to-insulin ratio areas under the curve during the 12-h metabolic profile were similar for both therapies, suggesting that glimepiride did not worsen the previous secretor dysfunction of beta cells, as reported with other sulfonylureas (35).

The overall effects on plasma lipids were small, with similar lowering of fasting VLDL cholesterol and triglyceride levels after the initial four months of both therapies. TG levels during the 12-h metabolic profile were also similar between the two drugs. LDL and HDL cholesterol levels were unaffected by treatment. Metformin and glimepiride-associated improvements in lipid metabolism were expected, but the reported changes have been small (3,16,30). The near normal triglyceride and cholesterol levels of our patients prior to both therapies were likely factors that influenced these modest results.

Despite causing lower insulin and higher glucagon secretion, metformin kept glucose and lipid profiles at similar levels compared with glimepiride. The blood glucose-lowering actions of metformin result primarily from an amelioration of insulin resistance, increase in peripheral glucose disposal, decrease in fatty acid oxidation and activation of the enzyme adenosine monophosphate (AMP) kinase to increase glucose transporter 4 (GLUT4) translocation to plasma membrane cells in the muscles and fat and reduce gluconeogenesis in the liver (3,36). The higher glucagon levels are unlikely to have interfered with metformin's action, as glucagon appears to have little effect on the presence of insulin, suggesting that its diabetogenic action occurs only under conditions of high insulin deficiency (37). In addition, glucagon's effects on hepatic glucose production could have been strongly counteracted by metformin.

Hemostatic factors

Glimepiride and metformin had similar effects on hemostatic factors; both increased plasminogen levels (significant after four months on each therapy) and decreased t-PA activity. These findings observed in the G group have not been reported previously.

The improvements in fibrinolysis after metformin and glimepiride therapy suggested by the increases in plasminogen levels occurred along with an unexpected decrease in another marker of fibrinolysis — t-PA activity. In contrast to our results, an increase in t-PA activity during metformin therapy has been described (38). However, as active t-PA declines as a function of increasing concentrations of PAI-1 and considering that PAI-1 antigen and activity did not change in our experiment, we can speculate that the formed complexes of PAI-1 and t-PA cleared in an accelerated fashion by the liver (39) contributed to the decrease in t-PA activity and that an impairment of fibrinolysis continued to occur. These data are likely implicated in the high frequency of cardiovascular disease in type 2 diabetes. No additional effects of any of the therapies were observed on the other

hemostatic factors analyzed (fibrinogen levels and platelet aggregation — data not shown).

Vascular reactivity

Increased carotid artery systolic diameter and blood flow were demonstrated only with metformin therapy, whereas an opposite trend in carotid diameter was observed with glimepiride use. Similar trends were observed even after patients treated for hypertension were excluded. No changes were observed for the other properties of the carotid (compliance, distensibility, and intima-media thickness) or brachial arteries (flow and diameter measurements after stimulus with reactive hyperemia and nitroglycerin).

Metformin's effects on carotid artery flow, independent of glucose decay or changes in systolic and diastolic blood pressure, likely afford more protection against cerebral diabetes complications. Its countering of insulin resistance action is likely implicated. Furthermore, we observed a negative correlation between fasting serum insulin levels and carotid compliance in the M group ($r = -0.5$; $p = 0.04$). Several observational analyses have suggested cardioprotective benefits with metformin use in patients with cardiovascular disease (3-4,40). Patients on metformin exhibited reduced nitroxidant metabolites and increased nitric oxide levels (30). As the carotid artery is more elastic and more proximal to the heart than the brachial artery, it might be more amenable to these improvements.

As both therapies improved glucose control in very similar manners, their beneficial effects on lipid profiles, hemostatic factors and norepinephrine levels can likely be ascribed to improvement in the metabolic milieu and not to a specific drug effect. Our study is the first to demonstrate decreases in norepinephrine levels after the first four months of glimepiride therapy and similar trends for epinephrine levels. The same results were obtained for metformin.

Although metformin has been confirmed as the first-line option for treating diabetes, troublesome gastrointestinal intolerance occasionally precludes its use (3). Thus, sulfonylureas remain important adjuvants for patients with intolerance to metformin or limited insulin secretion (5). When compared with metformin, glimepiride achieved similar efficacy in controlling weight and improving metabolic and hemostatic factors. Although it led to greater insulin secretion, glimepiride did not worsen beta cell function, as measured by the proinsulin-to-insulin ratio or vascular reactivity. Long-term studies are needed to ascertain whether glimepiride can reduce beta cell exhaustion or apoptosis.

Our study has some limitations. The four-month treatment duration could have not been sufficient to demonstrate all of the effects of these medications. Additionally, as a crossover study with no washout period, a treatment period interaction effect was demonstrated for some variables (triglyceride, VLDL cholesterol, plasminogen and norepinephrine levels). To minimize this problem, only the values from the first treatment period were analyzed.

In patients with type 2 diabetes inadequately controlled by dietary therapy, M and G resulted in similar overall improvements in glycemic control, lipid profiles, norepinephrine levels, and levels of the fibrinolytic factor plasminogen. All of these beneficial effects were likely due to improvement of the metabolic environment and were not drug-specific. However, as both therapies reduced t-PA

activity, the coagulation process continued, which can worsen cardiovascular disease. Only metformin countered insulin resistance and induced an increase in carotid artery diameter and blood flow indices. Neither drug affected small brachial artery vasodilation.

ACKNOWLEDGMENTS

We are grateful to Greci de Paula Souza for assistance with procedures. This study was supported by the Fundação de Amparo à Pesquisa do Estado de São Paulo-FAPESP.

AUTHOR CONTRIBUTIONS

Machado HA and Cunha MR participated in designing the studies, conducting the clinical and experimental procedure and writing the paper. Vieira M provided the cardiovascular evaluation. Correia MRS performed the biochemical analysis. Fukui RT performed the hormonal analysis. Santos RF, Wajchenberg BL and Rocha DM collaborated on the clinical study. Lage SG collaborated on the clinical study and the cardiovascular evaluation. Silva MER collaborated on the clinical study and wrote, edited and finalized the manuscript for publication.

REFERENCES

1. Stolar M. Glycemic control and complications in type 2 diabetes mellitus. *Am J Med.* 2010;123(Suppl. 3):S3-S11, <http://dx.doi.org/10.1016/j.amjmed.2009.12.004>.
2. American Diabetes Association - Standards of medical care in diabetes. *Diabetes Care.* 2009;32(Suppl. 1):S13-S61.
3. Scarpello JHB, Howlett HCS. Metformin therapy and clinical uses. *Diabetes and Vascular Disease Research* 2008;5:157-67.
4. Bailey CJ. Metformin: Effects on micro and macro vascular complications in type 2 diabetes. *Cardiovascular Drugs Ther.* 2008; 22(3):215-24, <http://dx.doi.org/10.1007/s10557-008-6092-0>.
5. Draeger KE, Wernicke-Panten K, Lomp HJ, Schuler E, Rosskamp R. Long-term treatment of type 2 diabetic patients with the new oral antidiabetic agent glimepiride (Amaryl): a double-blind comparison with glibenclamide. *Horm Metab Res.* 1996;28(9):419-25, <http://dx.doi.org/10.1055/s-2007-979830>.
6. Mori RC, Hirabara SM, Hirata AE, Okamoto MM, Machado UF. Glimepiride as insulin sensitizer: increased liver and muscle responses to insulin. *Diabetes Obes Metab.* 2008;10(7):596-600.
7. Sadikot SM, Mogensen CE. Risk of coronary artery disease associated with initial sulphonylurea treatment of patients with type 2 diabetes: a matched case-control study. *Diabetes Res Clin Pract.* 2008;82(3):391-5, <http://dx.doi.org/10.1016/j.diabres.2008.09.004>.
8. Monami M, Luzzi C, Lamanna C, Chiasserini V, Addante F, Desideri CM, et al. Three-year mortality in diabetic patients treated with different combinations of insulin secretagogues and metformin. *Diabetes Metab Res Rev.* 2006;22(6):477-82, <http://dx.doi.org/10.1002/dmrr.642>.
9. Bijlstra PJ, Lutterman JA, Russel FGM. Interaction of sulfonylurea derivative with vascular ATP-sensitive potassium channels in humans. *Diabetologia.* 1996;39(9):1083-90, <http://dx.doi.org/10.1007/BF00400658>.
10. Klepzig H, Kober G, Matter C, Luus H, Schneider H, Boedeker KH, et al. Sulfonylureas and ischemic preconditioning. A double-blind, placebo controlled evaluation of glimepiride and glibenclamide. *European Heart Journal.* 1999;20(6):439-46.
11. Lee TM, Chou TF. Impairment of myocardial protection in type 2 diabetic patients. *J Clin Endocrinol Metab.* 2003;88(2):531-7, <http://dx.doi.org/10.1210/jc.2002-020904>.
12. Abbink EJ, Pickkers P, Jansen van RA, Lutterman JA, Tack CJ, Russel FG, et al. Vascular effects of glibenclamide vs. glimepiride and metformin in Type 2 diabetic patients. *Diabet Med.* 2002;19(2):136-43.
13. Spallarossa P, Schiavo M, Rossettin P, Cordone S, Olivotti L, Cordera R, et al. Sulfonylurea treatment of type 2 diabetic patients does not reduce the vasodilator response to ischemia. *Diabetes Care.* 2001;24(4):738-42, <http://dx.doi.org/10.2337/diacare.24.4.738>.
14. Pfützner A, Marx N, Lübgen G, Langenfeld M, Walcher D, Konrad T, et al. Improvement of cardiovascular risk markers by pioglitazone is independent from glycemic control: results from the pioneer study. *J Am Coll Cardiol.* 2005;45(12):1925-31, <http://dx.doi.org/10.1016/j.jacc.2005.03.041>.
15. Yener S, Comlekci A, Akinci B, Demir T, Yuksel F, Ozcan MA, et al. Soluble CD40 ligand, plasminogen activator inhibitor-1 and thrombin-activatable fibrinolysis inhibitor-1-antigen in normotensive type 2 diabetic subjects without diabetic complications. Effects of metformin and rosiglitazone. *Med Princ Pract.* 2009;18(4):266-71, <http://dx.doi.org/10.1159/000215722>.

16. Rizzo MR, Barbieri M, Grella R, Passariello N, Paolisso G. Repaglinide has more beneficial effect on cardiovascular risk factors than glimepiride: data from meal-test study. *Diabetes Metab.* 2005;31(3 Pt 1):255-60, [http://dx.doi.org/10.1016/S1262-3636\(07\)70192-1](http://dx.doi.org/10.1016/S1262-3636(07)70192-1).
17. Derosa G, Franzetti I, Gadaleta G, Ciccarelli L, Fogari R. Metabolic variations with oral antidiabetic drugs in patients with Type 2 diabetes: comparison between glimepiride and metformin. *Diabetes Nutr Metab.* 2004;17:143-50.
18. Derosa G, Mugellini A, Ciccarelli L, Crescenzi G, Fogari R. Comparison between repaglinide and glimepiride in patients with type 2 diabetes mellitus: a one-year, randomized, double-blind assessment of metabolic parameters and cardiovascular risk factors. *Clin Ther.* 2003;25(2):472-84, [http://dx.doi.org/10.1016/S0149-2918\(03\)80090-5](http://dx.doi.org/10.1016/S0149-2918(03)80090-5).
19. Luis BJ, Bugos C, Dirnberger G, Atherton T. Efficacy and safety profile of glimepiride in Mexican American Patients with type 2 diabetes mellitus: a randomized, placebo-controlled study. *Clin Ther.* 2003;25(1):194-209, [http://dx.doi.org/10.1016/S0149-2918\(03\)90025-7](http://dx.doi.org/10.1016/S0149-2918(03)90025-7).
20. Corretti MC, Anderson TJ, Benjamin EJ, Celermajer D, Charbonneau F, Creager MA, et al. Guidelines for the ultrasound assessment of endothelial-dependent flow-mediated vasodilation of the brachial artery: a report of the International Brachial Artery Reactivity Task Force. *J Am Coll Cardiol.* 2002;39(2):257-65, [http://dx.doi.org/10.1016/S0735-1097\(01\)01746-6](http://dx.doi.org/10.1016/S0735-1097(01)01746-6).
21. Latt JA, Turner K. Evaluations of Trinder's glucose oxidase method for measuring glucose in serum and urine. *Clin Chem.* 1975;21(12):1754-60.
22. Trivelli LA, Ranney HM, Lai HT. Hemoglobin components in patients with diabetes mellitus. *New Engl J Med.* 1971 284:353-57.
23. Warnick GR. Enzymatic methods for quantification of lipoprotein lipids. *Methods Enzymol.* 1986;129:101-23, [http://dx.doi.org/10.1016/0076-6879\(86\)29064-3](http://dx.doi.org/10.1016/0076-6879(86)29064-3).
24. Warnick GR, Knopp RH, Fitzpatrick V, Brauson L. Estimating low-density lipoprotein cholesterol by the Friedewald equation is adequate for classifying patients on the basis of nationally recommended cutpoints. *Clin Chem.* 1990;36(1):15-9.
25. Bouloux P, Perret D, Besser GM. Methodological considerations in the determination of plasma catecholamines by high-performance liquid chromatography with electrochemical detection. *Ann Clin Biochem.* 1985;22(Pt 2):194-203.
26. Clauss A. Rapid physiological coagulation method for determination of fibrinogen. *Acta Haematol.* 1957;17(4):237-46, <http://dx.doi.org/10.1159/000205234>.
27. Born GVR. Quantitative investigation into the aggregation of blood platelets. *J Physiol.* 1962;162:67-8.
28. Singer JM, Andrade DF. Analysis of longitudinal data. In *Handbook of Statistics. Volume 18: Bio-Environmental and Public Health Statistics.* eds. P.K. Sen and C.R. Rao. Amsterdam: North Holland.2000;115-60.
29. Neter J, Kutner MH, Nachtsheim CJ, Wasserman W. In Richard D. Irwing ed. *Applied Linear Statistical Models*, 4. Ed. Illinois. 1996:1408p.
30. Meaney E, Vela A, Samaniego V, Meaney A, Asbún J, Zempoalteca JC, et al. Metformin, arterial function, intima-media thickness and nitroxidation in metabolic syndrome: the mefisto study. *Clin Exp Pharmacol Physiol.* 2008;35(8):895-903, <http://dx.doi.org/10.1111/j.1440-1681.2008.04920.x>.
31. Stocker DJ, Taylor AJ, Langley RW, Jezior MR, Vigersky RA. A randomized trial of the effects of rosiglitazone and metformin on inflammation and subclinical atherosclerosis in patients with type 2 diabetes. *Am Heart J.* 2007;153(3):445.e1-6, <http://dx.doi.org/10.1016/j.ahj.2006.11.005>.
32. Huang Z, Lei MX, Liu L, Tang QB. Effects of rosiglitazone on the IMTC and serum MMP-9 levels in newly diagnosed type 2 diabetic patients. *Zhong Nan Da Xue Xue Bao Yi Xue Ban.* 2006;31(3):367-72.
33. Stakos DA, Schuster DP, Sparks EA, Wooley CF, Osei K, Boudoulas H. Long term cardiovascular effects of oral antidiabetic agents in non-diabetic patients with insulin resistance: double blind, prospective, randomized study. *Heart.* 2005;91(5):589-94, <http://dx.doi.org/10.1136/hrt.2003.027722>.
34. Clark HE, Matthews DR. The effect of glimepiride on pancreatic beta - cell function under hyperglycaemic clamp and hyperinsulinaemic, euglycaemic clamp conditions in non. insulin. dependent diabetes mellitus. *Horm Metab Res.* 1996;28(9):445-60.
35. Inoguchi T, Umeda F, Kakimoto M, Sako Y, Ishii H, Noda K, et al. Chronic sulfonylurea treatment and hyperglycemia aggravate disproportionately elevated plasma proinsulin levels in patients with 2 diabetes. *Endocr J.* 2000;47(6):763-70, <http://dx.doi.org/10.1507/endocrj.47.763>.
36. Reitman ML, Schadt EE. Pharmacogenetics of metformin response: a step in the path toward personalized medicine. *J Clin Invest.* 2007; 117(5):1226-9, <http://dx.doi.org/10.1172/JCI32133>.
37. Bansal P, Wang Q. Insulin as a physiological modulator of glucagon secretion. *Am J Physiol Endocrinol Metab.* 2008;295(4):E751-E761, <http://dx.doi.org/10.1152/ajpendo.90295.2008>.
38. Grant PJ. Beneficial effects of metformin on haemostasis and vascular function in man. *Diabetes Metab.* 2003;29(4 Pt):6S44-52, [http://dx.doi.org/10.1016/S1262-3636\(03\)72787-6](http://dx.doi.org/10.1016/S1262-3636(03)72787-6).
39. Wing LR, Hawksworth GM, Bennett B, Booth NA. Clearance of t-PA, PAI-1, and t-PA-PAI-1 complex in an isolated perfused rat liver system. *J Lab Clin Med.* 1991;117(2):109-14.
40. Roussel R, Travert F, Pasquet B, Wilson PW, Smith SC Jr, Goto S, et al. For the Reduction of Atherothrombosis for Continued Health (REACH) Registry Investigators. Metformin Use and Mortality Among Patients With Diabetes and Atherothrombosis. *Arch Intern Med.* 2010;170(21): 1892-9.