

Controlled medial branch anesthetic block in the diagnosis of chronic lumbar facet joint pain: the value of a three-month follow-up

Ivan Dias da Rocha, Alexandre Fogaça Cristante, Raphael Martus Marcon, Reginaldo Perilo Oliveira, Olavo Biraghi Letaif, Tarcisio Eloy Pessoa de Barros Filho

Instituto de Ortopedia e Traumatologia, Hospital das Clínicas da Faculdade de Medicina da Universidade de São Paulo (IOT-HCFMUSP), Spine Surgery Division, São Paulo/SP, Brazil.

OBJECTIVES: To verify the incidence of facetary and low back pain after a controlled medial branch anesthetic block in a three-month follow-up and to verify the correlation between the positive results and the demographic variables.

METHODS: Patients with chronic lumbar pain underwent a sham blockade (with a saline injection) and then a controlled medial branch block. Their symptoms were evaluated before and after the sham injection and after the real controlled medial branch block; the symptoms were reevaluated after one day and one week, as well as after one, two and three months using the visual analog scale. We searched for an association between the positive results and the demographic characteristics of the patients.

RESULTS: A total of 104 controlled medial branch blocks were performed and 54 patients (52%) demonstrated >50% improvements in pain after the blockade. After three months, lumbar pain returned in only 18 individuals, with visual analogue scale scores >4. Therefore, these patients were diagnosed with chronic facet low back pain. The three-months of follow-up after the controlled medial branch block excluded 36 patients (67%) with false positive results. The results of the controlled medial branch block were not correlated to sex, age, pain duration or work disability but were correlated with patient age ($p<0.05$).

CONCLUSION: Patient diagnosis with a controlled medial branch block proved to be effective but was not associated with any demographic variables. A three-month follow-up is required to avoid a high number of false positives.

KEYWORDS: Low Back Pain; Zygapophyseal Joint; Anesthesia, Local; Nerve Block; Lidocaine; Diagnosis; Prevalence; Prospective Studies; False Positive Reactions; Spine.

Rocha ID, Cristante AF, Marcon RM, Oliveira RP, Letaif OB, Barros Filho TE. Controlled medial branch anesthetic block in the diagnosis of chronic lumbar facet joint pain: the value of a three-month follow-up. *Clinics*. 2014;69(8):529-534.

Received for publication on January 20, 2014; First review completed on February 24, 2014; Accepted for publication on February 24, 2014

E-mail: idr_77@hotmail.com

Tel.: 55 11 2645-3698

INTRODUCTION

The American College of Physicians and the American Pain Society recommend that patients with low back pain should be classified into one of three pain categories to allow for a proper clinical approach: back pain potentially associated with radiculopathy or spinal stenosis, back pain potentially associated with another specific spinal cause or nonspecific low back pain. Although imaging is recommended in cases with specific low back pain (i.e., when

severe or progressive neurologic deficits are present or when serious underlying conditions are suspected on the basis of history and physical examination), clinicians should not routinely obtain imaging or other diagnostic tests in patients with nonspecific low back pain (1); such patients can comprise 85% of all patients with lumbar pain complaints (2). Muscle contractions or myofascial syndromes, intervertebral disc or facet joint problems can cause the pain (1).

Between 8% and 12% of all patients with lumbar pain comprise chronic cases, with complaints lasting longer than three months (3,4). Facet joint disturbances can be responsible for 10% to 50% of all cases of chronic lumbar pain (5-8). However, clinical history or physical examination cannot identify facet joint alterations as the origin of pain (7,9-13), nor does imaging (e.g., radiography, computed tomography [CT] or magnetic resonance imaging [MRI]) (7,14-16). The only tool to identify facet joint alterations as the cause of

Copyright © 2014 CLINICS – This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

No potential conflict of interest was reported.

DOI: 10.6061/clinics/2014(08)05



pain is the verification of an analgesic response to anesthetic injections into the zygapophyseal joints or at their nerve supplies (17-20) and medial dorsal branch blocks are easier to perform (21). The diagnostic power of the blockade is based on the assumption that anesthetizing the facet joint or the capsule containing the innervations would result in pain relief. A positive result (i.e., pain relief) would mean that the facet joint is the site from which the pain originates. The technique of medial dorsal branch block consists of blocking each of the medial branches that innervate a facet above and a facet below their corresponding roots and also blocking the multifidus and interspinous muscles in the region of the corresponding dermatome (22,23).

However, several studies (20,24-26) have documented a high rate (varying from 25% to 40%) of false positive diagnoses in controlled medial branch blocks (CMBs) depending on the use of different anesthetics or saline solution as the control. There are many reasons for the high rates of false positives (e.g., the placebo response, use of sedation during CMBs, liberal use of anesthesia of the skin and subcutaneous tissue, use of large volumes of anesthetic, causing dispersion of anesthetics to other structures) (27). To minimize false positives, the following solutions are recommended: using saline controls (7,18,28,29) to avoid confusion with myofascial pain (24), reducing the amount of anesthetic applied to 0.5 ml at each point to avoid or reduce anesthesia of the skin as much as possible and avoiding the use of sedation for the procedure (7). No study, however, has investigated CMB after a three-month follow-up to verify whether the results are the same after such a period.

Therefore, the objective of this study was to determine the prevalence of low back pain after CMB immediately after the blocks and also after three months. The hypothesis to be verified here is that three-months of follow-up can modify the CMB results, thus, evidencing false-positives and altering therapeutic planning.

METHODS

Design, setting and ethics

This prospective, controlled, diagnostic study was performed as the first phase of a randomized controlled clinical trial conducted by our team to analyze the effect of neurolysis in chronic facetary lumbar pain (registered in Clinicaltrials.gov under the number NCT01367860). This diagnostic study, which concerns the value of a three-month follow-up in diagnosing facetary pain in patients undergoing controlled medial branch block (CMBs), was approved by the local ethics committee (under protocol 337/10); all participants provided signed informed consent. The patients were informed that the procedure would be performed without topical analgesia and that there was a small risk of allergy to the injected lidocaine. The study was conducted from March 2011 to January 2013.

Sample and participant recruitment

The participants in this study were patients with lumbar back pain being treated in public primary care settings in São Paulo city; they were referred to our specialized center or informed by relatives or acquaintances who had learned of the study and contacted our screening center. The study was open to anyone with lumbar back pain, regardless of the region of residence or registration at our hospital. All participants called the center by telephone and spoke with a

nurse who applied the triage criteria: age (18 to 76 years old), lumbar back pain for three months or more without a specific diagnosis and the ability to visit the hospital for clinical evaluation. The nurse also collected certain demographic data, such as educational level and marriage status.

Following this assessment, an orthopedic surgeon (IDR) conducted a clinical evaluation, verifying the presence of nonspecific chronic back pain and using a visual analog scale (VAS) to evaluate pain intensity. Pain with a score ≥ 4 was a criterion for study inclusion.

The following exclusion criteria were applied: pain irradiating to the limbs (sciatica), previous lumbar spine surgery, pregnancy, history of drug abuse, history of psychiatric problems, rheumatologic diseases, fibromyalgia or lidocaine allergy, as well as the presence of fever or neurologic abnormalities in the physical exam. The patients who did not sign the consent form were also excluded.

Interventions and outcome

For CMB, the patient was positioned in the prone position and monitored for vital signs. The participant was not sedated and did not receive any analgesic/anesthetic in the skin. Once positioned, the patient was asked to note (with a pencil) the intensity of pain using the VAS.

After the VAS was registered, the patient was warned that the test would begin with one injection and was asked to indicate the levels/sites of the most intense back pain. Saline solution (1 ml) was injected at these sites; 10 minutes later, the patient was again asked to indicate the pain intensity on the VAS. If the pain was improved by 50%, a placebo effect was detected and the patient was immediately excluded from the study and did not undergo CMB. If the pain was not improved, CMB was then applied. CMB was performed with a 25-G needle, 90 mm long, through the skin until reaching the medial branch nerves in L2, L3, L4 and L5 in both sides (Figure 1). A dose of 0.5 ml of lidocaine at 2%, without epinephrine, was used in all cases. All procedures were performed by the same surgeon (IDR).

After 10 minutes, the patient was again asked to note the pain intensity using VAS. If the pain improved by only




Figure 1 - A radiograph in the anterior-posterior view during the controlled medial branch block procedure in one patient, with needles positioned in the medial branches of L3, L4 and L5 (left).



≤50%, the patient was excluded from the study and was referred for MRI to investigate the origin of the symptom. If the pain improved by >50% using VAS, the patient was included in the study and followed up.

The following day, the patient again visited the center for pain evaluation. The VAS analysis was repeated after one week, one month and three months. The VAS score was expected to be <4 in these evaluations; a score ≥4 would be an indication for neurolysis and the patient would then be included in the above mentioned clinical trial study. All VAS scores were recorded for statistical analysis.

Statistical analysis

At the end of the data collection, the patients’ VAS data were tested for normal distribution with the Kolmogorov-Smirnov test using histogram subjective analysis. Given a normal distribution of the data, the averages and standard deviations were computed and subjected to parametric tests, including an analysis of variance (ANOVA) to verify the homogeneity of variance and the t-test for two independent samples. To analyze the association of dichotomous data, Fisher’s exact test and chi-squared tests were used. For these data, we used the Kruskal-Wallis analysis for three or more groups and the Wilcoxon test for the comparison of two groups. For all tests, an alpha error of 5% was established.

RESULTS

During the study period, 213 patients were screened by telephone at the center and 101 (47%) were excluded in this interview. The remaining 112 were clinically evaluated and 8 (7%) were further excluded: 3 because they did not agree with the study conditions and 5 because of the exclusion criteria (Figure 2).

Among the 104 remaining patients who were initially evaluated with chronic back pain, 50 (48%) were men. Regarding race, 87 were white (85%), 7 were black (5%), 9 (7%) mulattos and 1 was Asian. Most patients (63%) were married or had been married for some time (15%). The patient ages ranged from 29 to 75 years (average, 49.56 ± 10.97 years). Table 1 shows the educational levels of the patient cohort.

At the moment of study inclusion, the patients had been experiencing low back pain for 33.11 ± 39.85 months. Most patients (80%) were not using opioids for pain. Regarding work disability and social security benefits, 41 patients (39%) had already stopped working and were receiving government benefits. Among the 63 patients (61%) who were not receiving benefits, 21 (33%) felt they should receive benefits because of their pain.

The initial VAS score (before the first injection) was 75.72 ± 18.20 on average.




Figure 2 - Flowchart detailing the inclusion and exclusion criteria of patients in the diagnostic study.



Table 1 - Education levels among the patients with chronic back pain.

Studied for 8 years	35 (33.3%)
Studied for 8-10 years	19 (18.5%)
Completed high school (11 years)	31 (29.6%)
Completed college	19 (18.5%)

Among the 104 individuals subjected to controlled medial branch block (CMB), 17 (16.3%) felt improvement with the saline solution injection and were excluded from this study, as were 33 (31.7%) patients who felt $\leq 50\%$ improvement after the lidocaine injection. These 33 patients were referred for MRI and then diagnosed with degenerative diseases, except for one patient who had an osteoid osteoma in S1. Therefore, this study was completed with 54 patients undergoing CMB and all VAS evaluations.

The average VAS score immediately after CMB was 43.27 ± 31.23 (for the entire sample, $n=104$). The average VAS score was 67.78 ± 20.74 immediately after the CMBs among the 33 patients who did not respond well to the lidocaine injection (pain improvement $\leq 50\%$), whereas in the 54 well-responding patients (pain improvement $> 50\%$), the VAS score was on average 16.7 ± 12.80 . Figure 3 shows the patient distribution based on pain recurrence after CMB.

The average age of the well-responding patients ($n=41$) was 33.29 ± 44.53 and the average age of the non-responding patients ($n=33$) was 29.82 ± 27.82 ($p < 0.05$).

Of the 54 patients, 36 (67%) maintained pain improvement after three months, with a VAS score of 25.40 ± 9.48 . The remaining 18 (33%) experienced increases in pain, with a VAS score > 4 and these patients were diagnosed with chronic lumbar facetary pain. Twelve patients (67%) felt pain again in the first week and 6 (33%) felt pain again one month after the CMBs.

The investigation of correlations between VAS scores and demographic variables, including the duration of pain, showed no significant association between any variable and VAS scores ($p \geq 0.102$) or between the demographic variables and VAS scores before and after CMB ($p \geq 0.197$).

DISCUSSION

Imaging exams have been considered to be poor tools to identify chronic lumbar pain originating from the facet joints (30-33), as are a patient's clinical features and history




Figure 3 - Patient distribution at each evaluation time, according to their VAS scores ($n=54$).

(25,26). To the best of our knowledge, no controlled study has investigated the diagnosis of facet joint chronic lumbar pain using controlled medial branch anesthetic block (CMB) in an attempt to identify the prevalence of false positives in a three-month follow-up. Other studies have used CMB as a diagnostic tool for facet joint pain (34-37), but none have reevaluated the patients weeks after the blockade. We show here that it is important to reevaluate patients for weeks following the CMB procedure because when a patient begins to feel discomfort again after the effects of the lidocaine block have passed, it means that the source of pain was, in fact, the structure being anesthetized. However, as was the case of 67% of the patients ($n=36$) in our sample, when pain improvement is maintained, it likely indicates that the symptoms do not originate in the medial branch nerves, which are no longer anesthetized; most likely, the cause of pain is myofascial and has been resolved in another manner. The remaining 33% of patients ($n=18$) still had VAS scores > 4 in the three-month follow-up, confirming the diagnosis of chronic lumbar facetary pain and these patients began to experience pain again in the first week (12) or first month (6) after the CMBs.

Patients undergoing CMBs can feel relief for an average of up to 14 weeks (25,26). In our study, the 36 patients with improvement after three months most likely had myofascial pain and the facet joint origin of the pain was correctly removed after the final follow-up, allowing them to be treated properly. Other studies did not follow positive patients after such a period (34-37); in such a situation, these 36 patients could have been inappropriately included in the neurolysis clinical trial as if they indeed had pain originating in the facet joint.

False-positive rates can be as high as 38% in CMB using lidocaine only in patients with low back pain (24). The false positive rate calculated from the result of bupivacaine-controlled block 3-4 weeks after the first procedure was found to be 27% for the lumbar spine (6). Considering that using a controlled procedure is important, we opted to limit confounding factors in our research, for instance, avoiding the use of sedation (7), which can interfere with neuronal plasticity (38) and restricting the volume of lidocaine to 0.5 ml per injection site, thus eliminating a possible myofascial effect (39). More importantly, however, was the use of a controlled block with saline solution (37,40) in a sham procedure that we were able to implement at the time of surgery, thereby eliminating the need to utilize the operating room, staff and materials a second time (with the associated costs). The placebo and lidocaine injections were 10 minutes apart, which allowed the immediate exclusion of patients feeling pain improvement with the saline solution.

Our study is most likely representative of the population (41) because the research was open to the community, without any restrictions regarding the place of residence or need for prior registration at our hospital. The patients were referred from primary care units from across the city or spontaneously reached us by telephone after being informed of the project by health professionals or acquaintances. In fact, as shown in other studies (6,25,26,42,43), we found no correlation of the results with demographic data, such as age, years of schooling or social security benefit.

One of the limitations in the present study is the fact that we did not evaluate depression symptoms, which could possibly interfere with chronic pain results. This evaluation could have been accomplished with scales such as the



Zung-DV (Zung Self Rating Depression Scale) or MPI-DLV (Pain-Related Psychosocial and Behavioral Aspects). Nevertheless, van Wijk et al. (36) used these measures and observed no differences in depression prevalence between treated and placebo groups of patients undergoing radio-frequency denervation. Another study limitation is that, although we recorded the opioid use by our chronic low back patients, we did not perform a quantitative or qualitative analysis on this or any other analgesic. A recent study has shown, however, that pain sensitivity in chronic low back pain patients is not altered by chronic opioid intake (44).

In conclusion, the diagnosis of facet joint lumbar pain through CMB was proven to be effective but was not associated with demographic variables. A three-month follow-up is important to discard myofascial pain cases, which accounted for 67% of patients undergoing CMB.

AUTHOR CONTRIBUTIONS

Rocha ID and Cristante AF were responsible for study design, experimental study planning, manuscript writing, final review of the manuscript, data collection, analysis and interpretation. Marcon RM, Oliveira RP, and Letaif OB were responsible for data analysis and interpretation and final review of the manuscript. Barros TE was responsible for study design, experimental study planning, manuscript writing and final review of the manuscript.

REFERENCES

- Chou R, Qaseem A, Snow V, Casey D, Cross JT Jr, Shekelle P, et al. Diagnosis and treatment of low back pain: a joint clinical practice guideline from the American College of Physicians and the American Pain Society. *Ann Intern Med.* 2007;147(7):478-91, <http://dx.doi.org/10.7326/0003-4819-147-7-200710020-00006>.
- van Tulder MW, Assendelft WJ, Koes BW, Bouter LM. Spinal radiographic findings and nonspecific low back pain. A systematic review of observational studies. *Spine (Phila Pa 1976).* 1997;22(4):427-34, <http://dx.doi.org/10.1097/00007632-199702150-00015>.
- Carragee EJ. Clinical practice. Persistent low back pain. *N Engl J Med.* 2005;352(18):1891-8.
- Hart LG, Deyo RA, Cherkin DC. Physician office visits for low back pain. Frequency, clinical evaluation, and treatment patterns from a U.S. national survey. *Spine (Phila Pa 1976).* 1995;20(1):11-9.
- Schwarzer AC, Aprill CN, Derby R, Fortin J, Kine G, Bogduk N. The relative contributions of the disc and zygapophysial joint in chronic low back pain. *Spine (Phila Pa 1976).* 1994;19(7):801-6, <http://dx.doi.org/10.1097/00007632-199404000-00013>.
- Manchikanti L, Boswell MV, Singh V, Pampati V, Damron KS, Beyer CD. Prevalence of facet joint pain in chronic spinal pain of cervical, thoracic, and lumbar regions. *BMC Musculoskelet Disord.* 2004;5:15, <http://dx.doi.org/10.1186/1471-2474-5-15>.
- Cohen SP, Raja SN. Pathogenesis, diagnosis, and treatment of lumbar zygapophysial (facet) joint pain. *Anesthesiology.* 2007;106(3):591-614, <http://dx.doi.org/10.1097/0000542-200703000-00024>.
- Freburger JK, Holmes GM, Agans RP, Jackman AM, Darter JD, Wallace AS, et al. The rising prevalence of chronic low back pain. *Arch Intern Med.* 2009;169(3):251-8, <http://dx.doi.org/10.1001/archinternmed.2008.543>.
- Hirsch C, Ingelmark BE, Miller M. The anatomical basis for low back pain. Studies on the presence of sensory nerve endings in ligamentous, capsular and intervertebral disc structures in the human lumbar spine. *Acta Orthop Scand.* 1963;33:1-17.
- McCall IW, Park WM, O'Brien JP. Induced pain referral from posterior lumbar elements in normal subjects. *Spine (Phila Pa 1976).* 1979;4(5):441-6, <http://dx.doi.org/10.1097/00007632-197909000-00009>.
- Fairbank JC, Park WM, McCall IW, O'Brien JP. Apophyseal injection of local anesthetic as a diagnostic aid in primary low-back pain syndromes. *Spine (Phila Pa 1976).* 1981;6(6):598-605, <http://dx.doi.org/10.1097/00007632-198111000-00011>.
- Lippitt AB. The facet joint and its role in spine pain. Management with facet joint injections. *Spine (Phila Pa 1976).* 1984;9(7):746-50, <http://dx.doi.org/10.1097/00007632-198410000-00016>.
- Cohen SP, Huang JH, Brummett C. Facet joint pain—advances in patient selection and treatment. *Nat Rev Rheumatol.* 2013;9(2):101-16.
- Jackson RP, Jacobs RR, Montesano PX. 1988 Volvo award in clinical sciences. Facet joint injection in low-back pain. A prospective statistical study. *Spine (Phila Pa 1976).* 1988;13(9):966-71, <http://dx.doi.org/10.1097/00007632-198809000-00002>.
- Jensen MC, Brant-Zawadzki MN, Obuchowski N, Modic MT, Malkasian D, Ross JS. Magnetic resonance imaging of the lumbar spine in people without back pain. *N Engl J Med.* 1994;331(2):69-73.
- Schwarzer AC, Wang SC, O'Driscoll D, Harrington T, Bogduk N, Laurent R. The ability of computed tomography to identify a painful zygapophysial joint in patients with chronic low back pain. *Spine (Phila Pa 1976).* 1995B20(8):907-12, <http://dx.doi.org/10.1097/00007632-199504150-00005>.
- Dreyer SJ, Dreyfuss PH. Low back pain and the zygapophysial (facet) joints. *Arch Phys Med Rehabil.* 1996;77(3):290-300, [http://dx.doi.org/10.1016/S0003-9993\(96\)90115-X](http://dx.doi.org/10.1016/S0003-9993(96)90115-X).
- North RB, Kidd DH, Zahurak M, Piantadosi S. Specificity of diagnostic nerve blocks: a prospective, randomized study of sciatica due to lumbosacral spine disease. *Pain.* 1996;65(1):77-85, [http://dx.doi.org/10.1016/0304-3959\(95\)00170-0](http://dx.doi.org/10.1016/0304-3959(95)00170-0).
- Kaplan M, Dreyfuss P, Halbrook B, Bogduk N. The ability of lumbar medial branch blocks to anesthetize the zygapophysial joint. A physiologic challenge. *Spine (Phila Pa 1976).* 1998;23(17):1847-52, <http://dx.doi.org/10.1097/00007632-199809010-00008>.
- Dreyfuss PH, Dreyer SJ, NASS. Lumbar zygapophysial (facet) joint injections. *Spine J.* 2003;3(3 Suppl):50S-59S, [http://dx.doi.org/10.1016/S1529-9430\(02\)00450-3](http://dx.doi.org/10.1016/S1529-9430(02)00450-3).
- Bogduk N, Long DM. The anatomy of the so-called "articular nerves" and their relationship to facet denervation in the treatment of low-back pain. *J Neurosurg.* 1979;51(2):172-7.
- Pedersen HE, Blunck CF, Gardner E. The anatomy of lumbosacral posterior rami and meningeal branches of spinal nerve (sinu-vertebral nerves); with an experimental study of their functions. *J Bone Joint Surg Am.* 1956;38-A(2):377-91.
- Bogduk N. The innervation of the lumbar spine. *Spine (Phila Pa 1976).* 1983;8(3):286-93, <http://dx.doi.org/10.1097/00007632-198304000-00009>.
- Schwarzer AC, Aprill CN, Derby R, Fortin J, Kine G, Bogduk N. The false-positive rate of uncontrolled diagnostic blocks of the lumbar zygapophysial joints. *Pain.* 1994B58(2):195-200, [http://dx.doi.org/10.1016/0304-3959\(94\)90199-6](http://dx.doi.org/10.1016/0304-3959(94)90199-6).
- Manchikanti L, Pampati V, Fellows B, Baha AG. The inability of the clinical picture to characterize pain from facet joints. *Pain Physician.* 2000;3(2):158-66.
- Manchikanti L, Pampati V, Fellows B, Bakht CE. The diagnostic validity and therapeutic value of lumbar facet joint nerve blocks with or without adjuvant agents. *Curr Rev Pain* 2000;4(5):337-44, <http://dx.doi.org/10.1007/s11916-000-0016-4>.
- Hogan QH, Abram SE. Neural blockade for diagnosis and prognosis. A review. *Anesthesiology.* 1997;86(1):216-41, <http://dx.doi.org/10.1097/0000542-199701000-00026>.
- Lord SM, Barnsley L, Bogduk N. The utility of comparative local anesthetic blocks versus placebo-controlled blocks for the diagnosis of cervical zygapophysial joint pain. *Clin J Pain.* 1995;11(3):208-13.
- Lord SM, Barnsley L, Wallis BJ, McDonald GJ, Bogduk N. Percutaneous radio-frequency neurotomy for chronic cervical zygapophysial joint pain. *N Engl J Med.* 1996;335(23):1721-6.
- Carrera GF. Lumbar facet joint injection in low back pain and sciatica: preliminary results. *Radiology.* 1980;137(3):665-7.
- Carrera GF, Williams AL. Current concepts in evaluation of the lumbar facet joints. *Crit Rev Diagn Imaging.* 1984;21(2):85-104.
- Helbig T, Lee CK. The lumbar facet syndrome. *Spine (Phila Pa 1976).* 1988;13(1):61-4, <http://dx.doi.org/10.1097/00007632-198801000-00015>.
- Dolan AL, Ryan PJ, Arden NK, Stratton R, Wedley JR, Hamann W, et al. The value of SPECT scans in identifying back pain likely to benefit from facet joint injection. *Br J Rheumatol.* 1996;35(12):1269-73.
- Gallagher J, Petriccione di Vadi PL, Wedley JR. Radiofrequency facet joint denervation in the treatment of low back pain: a prospective controlled double-blind study to assess its efficacy. *Pain Clinic.* 1994;7:193-8.
- van Kleef M, Barendse GA, Kessels A, Voets HM, Weber WE, de Lange S. Randomized trial of radiofrequency lumbar facet denervation for chronic low back pain. *Spine (Phila Pa 1976).* 1999;24(18):1937-42, <http://dx.doi.org/10.1097/00007632-199909150-00013>.
- van Wijk RM, Geurts JW, Wynne HJ, Hammink E, Buskens E, Lousberg R, et al. Radiofrequency denervation of lumbar facet joints in the treatment of chronic low back pain: a randomized, double-blind, sham lesion-controlled trial. *Clin J Pain.* 2005;21(4):335-44.
- Nath S, Nath CA, Pettersson K. Percutaneous lumbar zygapophysial (Facet) joint neurotomy using radiofrequency current, in the management of chronic low back pain: a randomized double-blind trial. *Spine (Phila Pa 1976).* 2008;33(12):1291-7; discussion 1298, <http://dx.doi.org/10.1097/BRS.0b013e31817329f0>.
- Woolf CJ, Salter MW. Neuronal plasticity: increasing the gain in pain. *Science.* 2000;288(5472):1765-9, <http://dx.doi.org/10.1126/science.288.5472.1765>.
- Dreyfuss P, Schwarzer AC, Lau P, Bogduk N. Specificity of lumbar medial branch and L5 dorsal ramus blocks. A computed tomography



- study. *Spine (Phila Pa 1976)*. 1997;22(8):895-902, <http://dx.doi.org/10.1097/00007632-199704150-00013>.
40. Leclaire R, Fortin L, Lambert R, Bergeron YM, Rossignol M. Radiofrequency facet joint denervation in the treatment of low back pain: a placebo-controlled clinical trial to assess efficacy. *Spine (Phila Pa 1976)*. 2001;26(13):1411-6; discussion 1417, <http://dx.doi.org/10.1097/00007632-200107010-00003>.
 41. Hulley SB, Newman TB, Cummings SR. Choosing the study subjects: specification, sampling, and recruitment. In: Hulley SB, Cummings S, Browner WS, Grady DG, Newman TB. *Designing clinical research*. Philadelphia: Lippincott Williams & Wilkins; 2007. p. 27-36.
 42. Manchikanti L, Pampati V, Fellows B, Bakhit CE. Prevalence of lumbar facet joint pain in chronic low back pain. *Pain Physician*. 1999;2(3):59-64.
 43. Sowa G. Facet-mediated pain. *Dis Mon*. 2005;51(1):18-33, <http://dx.doi.org/10.1016/j.disamonth.2004.12.005>.
 44. Wang H, Fischer C, Chen G, Weinsheimer N, Gantz S, Schiltenswolf M. Does long-term opioid therapy reduce pain sensitivity of patients with chronic low back pain? Evidence from quantitative sensory testing. *Pain Physician*. 2012;15(3 Suppl):ES135-43.