

INSELBERGS AS A SOURCE OF β DIVERSITY IN A VEGETATION MATRIX IN COQUEIRAL, MINAS GERAIS, BRAZIL

INSELBERG COMO FONTE DE DIVERSIDADE β EM UMA MATRIZ VEGETACIONAL, NO MUNICÍPIO DE COQUEIRAL - MG

Gisele Cristina de Oliveira Menino¹ Rubens Manoel dos Santos² Daniel Salgado Pifano³
Rosângela Alves Tristão Borém⁴ Carlos Alberto Melo de Almeida⁵ Daniel Quedes Domingos⁶
Aline Martins Moreira⁷

ABSTRACT

This study aimed to answer the following question: Does habitat heterogeneity affect the tree community? To answer that question, we evaluated the influence of the species composition and structure of the tree community of inselbergs on the tree community of the matrix where the inselberg is located. The correlations between environmental and vegetation gradients were analyzed by canonical correspondence analysis of a species composition matrix constructed from abundance data of 40 plots sampled in two areas, one composed of seasonal semideciduous forest (SSF) and another with riparian forests, a vegetation corridor, and inselbergs. A similarity dendrogram of the four areas was calculated using the Dice-Sorensen index and UPGMA linkage method. The species indicator analysis was used to evaluate which species were characteristic of each area sampled. The eigenvalues of the first two axes were high indicating high species turnover. The correlations between environmental variables and plots indicated the formation of three groups: the first one formed by the SSF patch plots with high clay content; the second group formed by the vegetation corridor and inselberg plots, which did not separate as sharply as the first group, with a small separation in organic matter content; and the third group, formed by the riparian forest plots, which are located in areas of highest soil fertility, with high content of calcium, sum of bases and base saturation. The results showed high species turnover among areas, indicating that even though the areas are geographically close, the occurrence of inselbergs increased habitat heterogeneity, directly affecting the variation in species composition and community structure among areas.

Keywords: conservation; species replacement; forest patches.

RESUMO

O presente trabalho objetivou responder a seguinte pergunta: A heterogeneidade ambiental influencia a comunidade arbórea? Para isso avaliou-se a interferência da composição e estrutura da comunidade arbórea

- 1 Bióloga, Dr^a., Professora do Instituto Federal de Educação, Ciência e Tecnologia Goiano, Câmpus Rio Verde, Caixa Postal 66, CEP 75901-970, Rio Verde (GO). giselecristina.biologa@yahoo.com.br
- 2 Biólogo, Dr., Professor Adjunto II do Departamento de Ciências Florestais, Universidade Federal de Lavras, Campus Universitário, CEP 37200-000, Lavras (MG). rubensmanoel@dcf.ufla.br
- 3 Engenheiro Florestal, Dr., Professor Adjunto I do Colegiado de Ciências Biológicas, Universidade Federal do Vale do São Francisco, Campus de Ciências Agrárias. Rod. BR 407 Km 12 Lote 543 Projeto de Irrigação Senador Nilo Coelho, s/n, CI, CEP 56300-990, Petrolina (PE). daniel.pifano@univasf.edu.br
- 4 Engenheira Florestal, Dr^a., Professora Associado II do Departamento de Biologia, Universidade Federal de Lavras, CEP 37200-000, Lavras (MG). tristao@dbi.ufla.br
- 5 Biólogo, Graduando em Engenharia Agrônômica, Universidade Federal do Paraná, Setor de Ciências Agrárias, Rua dos Funcionários, 1540 - Juvevê, CEP 80035-050, Curitiba (PR). carlosgeas@yahoo.com.br
- 6 Biólogo, Departamento de Biologia, Universidade Federal de Lavras, CEP 37200-000, Lavras (MG). danielquedes@yahoo.com.br
- 7 Bióloga, Mestranda do Programa de Pós-graduação em Engenharia Florestal, Universidade Federal de Lavras, Caixa Postal 3037, CEP 37200-000, Lavras (MG). aline.martinsmoreira@posgrad.ufla.br

Recebido para publicação em 9/07/2012 e aceito em 26/11/2013

localizada em *inselberg* sobre a comunidade arbórea da matriz na qual o *inselberg* está inserido. Para isso fez-se uma CCA (análise de correspondência canônica) sobre uma matriz de abundância das espécies de 40 parcelas amostradas em dois fragmentos, um de Floresta Estacional Semidecidual (FES) e outro constituído por mata ciliar, corredor e *inselbergs*. Foi construído um dendrograma de similaridade para as quatro áreas utilizando o índice de Dice-Sorensen e o método de ligação UPGMA. A análise de espécies indicadoras foi usada para avaliar quais espécies foram características de cada área amostrada. Os autovalores dos dois primeiros eixos foram altos, evidenciando uma forte substituição das espécies. As correlações entre variáveis ambientais e parcelas indicaram a formação de três grupos: o primeiro formado pelas parcelas do fragmento FES, com alto teor de argila; o segundo grupo formado pelas parcelas do corredor e dos *inselbergs*, que não se separaram tão nitidamente como o primeiro grupo, apresentando uma pequena separação quanto ao teor de matéria orgânica; e o terceiro grupo, formado pelas parcelas da mata ciliar, com alto teor de magnésio, cálcio, soma de bases e saturação de bases, sendo solos de maior fertilidade. Os resultados evidenciaram alta substituição de espécies

entre as áreas, indicando que, apesar de ser bem próximas, a presença do *inselberg* aumentou a heterogeneidade ambiental, o que afeta diretamente a variação, tanto na composição de espécies quanto na estrutura entre as áreas.

Palavras-chave: conservação; substituição de espécies; fragmentos florestais.

INTRODUCTION

The conservation of biodiversity represents a great challenge due to the high level of anthropogenic disturbances of natural ecosystems (VIANA and PINHEIRO, 1998). There is an increasing trend to focus conservation efforts on ecotone areas, marginal ecosystems, and “fragments”, i.e., sub-units inserted in or associated with large biogeographical units. Stressing the need for specific conservation policies in areas with rocky outcrops, regardless of biome or vegetation type in which they are inserted (SCARANO, 2007). However, studies on plant communities in these environments are rare (POREMBSKI et al., 1998) and the vegetation on these outcrops is still quite unknown (SCARANO, 2007).

Inselbergs are monolithic mountains or groups of mountains that appear abruptly amidst the landscape, consisting mainly of granitic or gneissic rocks (POREMBSKI et al., 1998; POREMBSKI and BARTHLOTT, 2000). They naturally present unequal and spatially distributed resources (POREMBSKI and BARTHLOTT, 2000) which determine the heterogeneity (relative abundance of rocks and vegetation), structural complexity (shape and composition of rocks), and quality (abundance of rocks and vegetation resources) (MICHAEL et al., 2010) of the habitats. Moreover, the complexity of gradients, mainly soil depth and water availability, determine vegetation structure (GROGER and HUBER, 2007). The soils are shallow and characteristically nutrient-poor and

vary according to the rock with which they are associated (BENITES et al., 2007).

The vegetation of inselbergs is characterized as “a mosaic of marginal habitats” where a great number of taxa find adequate niches in an extremely condensed space (GRÖGER and HUBER, 2007). Although inselbergs present an apparently monotonous vegetation, they have a wide diversity of pedoenvironments in addition to an associated vegetation mosaic determined in great extent by the local relief and micro-environmental aspects (BENITES et al., 2007). These characteristics make inselbergs a center of diversity for plants well adapted to extreme conditions, particularly water shortage and low nutrient availability (POREMBSKI, 2007).

Inselbergs increase habitat heterogeneity. Some studies relate habitat variation to changes in species composition of communities, indicating that a greater variety of habitats and environmental conditions promote greater variation in species composition (MCINTYRE et al., 2001; BUCKLEY and JETZ, 2008; JANKOWSKI et al., 2009), which increases beta diversity. Because beta diversity reflects the dissimilarities in species composition between two or more areas, it reflects the differences in species composition at the landscape or habitat scale (WHITTAKER et al., 2001).

Although a significant number of inselbergs is located in conservation units (MMA 2002), management plans which take into account the particularities of those environments are still lacking. Moreover, inselbergs have been broadly disregarded from the biological standpoint (POREMBSKI,

2007). As a consequence of their adverse and unfavorable conditions, the vegetation of inselbergs is clearly influenced by its surroundings. The extent of the island character depends on the type of vegetation and floristic differentiation between the inselberg and its surrounding matrix (POREMBSKI, 2007), because there are strong relationships between inselbergs and the neighboring vegetation (SCARANO, 2007). Thus, phytosociology studies allow better knowledge of the composition and structure of the plant community, contributing towards better understanding of biogeographical patterns (FINA and MONTEIRO, 2009).

Most inselbergs studied have a small-sized vegetation, mostly composed of herbaceous and shrub species (FRANÇA et al., 1997; POREMBSKI and BARTHLOTT, 2000; CAIAFA and SILVA, 2007; RIBEIRO et al., 2007). Thus, studies in inselbergs with arboreal vegetation in the south of Minas Gerais state, Brazil are of fundamental importance for conservation efforts.

Thus, this study aimed to answer the following question: Does habitat heterogeneity affect the tree community? This hypothesis was tested by assessing: (i) the influence of the species composition of the tree community of an inselberg on the tree community of the matrix where the inselberg is located; (ii) the influence of the tree community structure of the inselberg on the tree community of the matrix where the inselberg is located. The evaluation of the influence of habitat heterogeneity on tree community enables the development of more effective conservation policies.

MATERIAL AND METHODS

Study area

The sample patches were located in the

municipality of Coqueiral, Minas Gerais state, Brazil (21°09'19" S and 45°28'17" W; 810–840 m altitude; Fig. 1). The climate is Cwb or mesothermal according to the Köppen classification, with mild summers and dry winters. The average annual rainfall is 1,493 mm and the average annual temperature is 19.3 °C (VILELA and RAMALHO, 1979). The first area is a 7.5 ha patch characterized as Seasonal Semideciduous Forest. In this area, 20 20 x 20 m plots, distant 20 m apart, were established in three transects parallel to the longest axis of the fragment. The second area is a mosaic composed of riparian forest, granitic-gneissic inselberg, and a vegetation corridor which accompanies the interior of the draining line over the inselberg linking the riparian forest to the inselberg. The inselberg plots were established in areas of litter deposition in rock crevices with shallow soils. Twenty plots, distant 20 m apart, were also established in this area: six in the riparian forest, nine in the vegetation corridor, and five in the inselberg. Inselberg plots only were 20 x 20 m, due to the more homogeneous, small-sized environment, whereas in other microenvironments plots were 10 x 40 m to maximize the greatest sampling intensity in the direction of least variation within each sampling unit, and consequently to sample the variation occurring in these environments. In each plot, all woody individuals with cbh (circumference at breast height) ≥ 15.7 cm were sampled, plant height was recorded, and the species were identified. All botanical material collected was stored in the ESAL herbarium at the Universidade Federal de Lavras. For soil characterization, a 500 g surface soil sample (0–20 cm depth) was collected in each plot. Each composite sample was obtained by mixing and homogenizing five subsamples randomly collected in each plot. The chemical and textural analyses of the samples were done in the Soil Analysis Laboratory at the Universidade Federal de



FIGURE 1: Forest patches in Coqueiral, Minas Gerais, Brazil.

FIGURA 1: Fragmentos localizados no município de Coqueiral - MG.

Lavras, following the EMBRAPA protocol (1997). The soil variables determined were: pH; contents of potassium, phosphorus-Mehlich, remaining-phosphorus, Calcium, Magnesium, Aluminum and Hydrogen Aluminum; sum of bases, base saturation; organic matter; aluminum saturation; effective cation exchange capacity and cation-holding capacity of the soil and contents of sand, silt, and clay.

Data analysis

To describe tree community structure, the classical quantitative descriptors were calculated per species: absolute and relative density, absolute and relative dominance, absolute and relative frequency, and importance value index (IVI) (MUELLER-DOMBOIS and ELLENBERG, 1974), in addition to the Shannon diversity index (H') and Pielou's equitability index (J) (BROWER and ZAR, 1984).

Tree density and basal area per plot were compared using analysis of variance (Anova) to test for differences in structure parameters among the microenvironments associated with the Tukey's test after testing for normality of the data using the D'Agostino test (ZAR, 1996). A species accumulation curve for the specific richness observed (Mao Tau) in each area was calculated after 500 randomizations using EstimateS 9.0 (COLWELL, 2013).

The correlations between environmental and vegetation gradients were analyzed by canonical correspondence analysis (CCA) (TER BRAAK, 1987) using PC-ORD for Windows version 4.14 software (MCCUNE and MEFFORD, 1999). A species composition matrix was constructed from abundance data per plot. According to the recommendations by TER BRAAK (1995), abundance values (a) were $\ln(a+1)$ transformed to compensate for the deviations caused by outliers.

The environmental variable matrix included one categorical variable that was used to define the four microenvironments and seventeen quantitative edaphic variables. Nevertheless, after a preliminary CCA was calculated, 12 quantitative variables were weakly correlated or redundant and excluded from the final analysis. Thus, the final CCA was calculated with the categorical variable and the six most representative variables that strongly correlated with the ordination axis: calcium (Ca), aluminum (Al), sum of bases (SB), base saturation (V), organic matter (MO), and clay. The Monte

Carlo permutation test was done (TER BRAAK, 1988) to test the significance of the model.

A similarity dendrogram of the four areas was calculated using the Dice-Sorensen index and UPGMA linkage method. We also calculated the cophenetic correlation coefficient, which measures the level of cophenetic correlation between the original similarity matrix and the calculated matrix. Values equal to or higher than 0.7 are considered satisfactory as representative of the original matrix (SHEPHERD, 2006). The similarity dendrogram and cophenetic correlation coefficient were calculated using Past software.

The species indicator analysis (DUFRENÉ and LEGENDRE, 1997) was used to evaluate which species were characteristic of each area sampled. The indicator value (ValInd) was compared with the results of 1,000 data randomizations using the Monte Carlo test. The calculations were done using PC-ORD for Windows version 4.14 software (MCCUNE and MEFFORD, 1999).

RESULTS

In the Seasonal Semideciduous Forest (SSF) patch 887 individuals distributed into 144 species were sampled. In the second fragment, consisting of three areas, 159 species were found: 82 in the riparian forest, 101 in the vegetation corridor, and 64 in the inselbergs, totaling 1,265 individuals, 286 sampled in the riparian forest, 601 in the corridor, and 378 in the inselbergs (Table 1). The basal area presented no statistical difference among the areas ($F_{0.05(1), 3.36} = 0.7877$; $p = 0.5114$), but there were significant differences in density ($F_{0.05(1), 3.36} = 97,3281$; $p < 0.0001$). The riparian forest differed from the corridor ($q = 4.7201$; $p < 0.05$), inselberg ($q = 6.0048$; $p < 0.01$) and of the SSF ($q = 11.4942$; $p < 0.01$). The SSF differed from the corridor ($q = 19.5272$; $p < 0.01$) and from the inselberg ($q = 0.179727$; $p < 0.01$). Only the corridor and inselberg did not differ as to density ($q = 2.0589$; $p > 0.05$). Comparisons were made using the mean values. These results show that community structure differed among the microenvironments, indicating that inselberg and vegetation corridor communities had a higher number of individuals of smaller CBH than riparian forest and SSF.

The species accumulation curve showed that, for the same number of individuals, the inselberg had the lowest species richness, followed by the vegetation corridor, riparian forest,

TABLE 1: Density (ind/ha), basal area (m²/ha), diversity - H' (nats/individual) and equability (J) in four microenvironments in the municipality of Coqueiral, Minas Gerais state, Brazil.

 TABELA 1: Densidade (ind/ha), Área basal (m²/ha), diversidade - H' (nats/indivíduo) e equabilidade (J) em quatro microambientes no município de Coqueiral - MG.

Area	Density (ind/ha)	Basal area (m ² /ha)	H' (nats/individual)	J
Riparian Forest	1191.67A	28.6A	3.88	0.88
Corridor	1669.44B	29.62A	4.03	0.87
Inselberg	1890.00B	22.22A	3.61	0.87
SSF Patch	1108.75C	29.33A	4.31	0.87

Where in: Parameters followed by the same letter in the column are not statistically different




FIGURE 2: Species accumulation curve after 500 randomizations of the number of individuals sampled in four microenvironments in the municipality of Coqueiral, Minas Gerais state, Brazil.

FIGURA 2: Curva de acumulação de espécies obtida de 500 randomizações do número de indivíduos amostrados em quatro microambientes no município de Coqueiral - MG.

and SSF (Fig. 2).

The most important species in the SSF patch were *Aspidosperma olivaceum* Müll.Arg., *Copaifera langsdorffii* Desf., *Aspidosperma australe* Müll.Arg., *Ocotea odorifera* (Vell.) Rohwer, *Tachigali rugosa* (Mart. ex Benth.) Zarucchi & Pipoly, *Amaioua intermedia* Mart. ex Schult. & Schult.f., *Mouriri glazioviana* Cogn., *Copaifera trapezifolia* Hayne, *Duguetia lanceolata* A.St.-Hil., and *Xylopia brasiliensis* Spreng. In the riparian forest, the most important species were *Platycamus regnellii* Benth., *Croton floribundus* Spreng., *Bathysa australis* (A.St.-Hil.) Benth. & Hook.f., *Cariniana legalis* (Mart.) Kuntze, and *Copaifera langsdorffii*; in the vegetation corridor *Callisthene major* Mart., *Copaifera langsdorffii*,

Tapirira obtusa (Benth.) J.D.Mitch., *Faramea nigrescens* Mart., and *Casearia arborea* (Rich.) Urb. were the most important species, whereas in the inselbergs, *Copaifera langsdorffii*, *Terminalia glabrescens* Mart., *Maytenus robusta* Reissek, *Platypodium elegans* Vogel, and *Myrciaria floribunda* (H.West ex Willd.) O.Berg were the most important species. Therefore, only *C. langsdorffii* was among the most important species in the four areas (Table 2).

The number of indicator species ranged among the areas: 17 in the riparian forest, 18 in the inselberg, 12 in the corridor, and nine in the SSF, which represents 20.73%, 28.12%, 11.88%, and 6.25% of the total number of species in each area, respectively.

The species with highest indicator values in the riparian forest were *Bauhinia longifolia* (Bong.) D.Dietr., *Croton floribundus*, *Machaerium hirtum* (Vell.) Stellfeld, *Celtis brasiliensis* (Gardn.) Planch., and *Senna macranthera* (Collad.) H.S.Irwin & Barneby. In the vegetation corridor, the species with highest indicator values were *Callisthene major*, *Faramea nigrescens*, *Siphoneugena densiflora* O.Berg, *Casearia arborea*, and *Myrsine umbellata* Mart., whereas the species *Maytenus robusta*, *Luehea divaricata* Mart., *Myrciaria floribunda*, *Eugenia cerasiflora* Miq., and *Zanthoxylum rhoifolium* Lam. were most indicative of inselbergs, and *Ocotea odorifera*, *Mouriri glazioviana*, *Duguetia lanceolata*, and *Aspidosperma olivaceum* were most indicative of the SSF patch (Table 3).

The eigenvalues of the first two axes were 0.465 and 0.369, respectively, and indicates high species turnover between the extremes of axis 1, which represents a soil quality gradient. Conversely, plots in axis 2 were separated by clay content. The

TABLE 2: Species of greatest importance value (IV) in the Riparian Forest, Inselberg, Vegetation Corridor, and Seasonal Semideciduous Forest Patch (SSF) in Coqueiral, Minas Gerais state, Brazil.

TABELA 2: Espécies de maior valor de importância (VI) da Mata ciliar, *Inselberg*, Corredor e Fragmento de Floresta Estacional Semidecidual (FES).

Environment/species	IV	Environment/species	IV
Riparian Forest		Corridor	
<i>Platycamus regnellii</i> Benth.	27.61	<i>Callisthene major</i> Mart.	23.33
<i>Croton floribundus</i> Spreng.	23.07	<i>Copaifera langsdorffii</i> Desf.	17.17
<i>Bathysa australis</i> (A.St.-Hil.) Benth. & Hook.f.	13.96	<i>Tapirira obtusa</i> (Benth.) J.D.Mitch.	15.59
<i>Cariniana legalis</i> (Mart.) Kuntze	13.58	<i>Farama nigrescens</i> Mart.	12.43
<i>Copaifera langsdorffii</i> Desf.	11.47	<i>Casearia arborea</i> (Rich.) Urb.	9.48
<i>Inga vera</i> Willd.	11.31	<i>Byrsonima laxiflora</i> Griseb.	8.95
<i>Metrodorea stipularis</i> Mart.	8.93	<i>Myrsine umbellata</i> Mart.	7.30
<i>Eugenia acutata</i> Miq.	8.33	<i>Siphoneugena densiflora</i> O.Berg	7.23
<i>Eugenia florida</i> DC.	7.07	<i>Protium spruceanum</i> (Benth.) Engl.	6.99
<i>Cecropia pachystachya</i> Trécul	6.94	<i>Terminalia glabrescens</i> Mart.	6.69
Environment/species	IV	Environment/species	IV
Inselberg		Seasonal Semideciduous forest	
<i>Copaifera langsdorffii</i> Desf.	24.03	<i>Aspidosperma olivaceum</i> Müll.Arg.	18.12
<i>Terminalia glabrescens</i> Mart.	20.42	<i>Copaifera langsdorffii</i> Desf.	16.40
<i>Maytenus robusta</i> Reissek	17.75	<i>Aspidosperma australe</i> Müll.Arg.	11.96
<i>Platyopodium elegans</i> Vogel	16.66	<i>Ocotea odorifera</i> (Vell.) Rohwer	11.09
<i>Myrciaria floribunda</i> (H.West ex Willd.) O.Berg	15.71	<i>Tachigali rugosa</i> (Mart. ex Benth.) Zarucchi & Pipoly	10.98
<i>Metrodorea stipularis</i> Mart.	10.60	<i>Amaioua intermedia</i> Mart. ex Schult. & Schult.f.	10.49
<i>Eugenia cerasiflora</i> Miq.	10.34	<i>Mouriri glazioviana</i> Cogn.	9.01
<i>Luehea divaricata</i> Mart.	9.68	<i>Copaifera trapezifolia</i> Hayne	7.84
<i>Eugenia acutata</i> Miq.	9.38	<i>Duguetia lanceolata</i> A.St.-Hil.	7.74
<i>Ixora brevifolia</i> Benth.	9.37	<i>Xylopia brasiliensis</i> Spreng.	7.05

correlation values between environmental variables and species was high (94% and 90%) and the Monte Carlo permutation test revealed a significant correlation between species abundance and environmental variables ($p < 0.005$). The proportion of the total variance explained by the first two axes was 12.2%, indicating that was a high level of 'noise', or remaining variance that was not explained by the environmental variables investigated. The correlations between environmental variables and plots indicated the formation of three groups (Fig. 3): the first one formed by the SSF patch plots with high clay content; the second group formed by the vegetation corridor and inselberg plots, which were not separated clearly, with a slight separation by organic matter content; and the third group, formed by the riparian forest plots, which are located in areas of highest soil fertility, with high content of calcium, sum of bases and base saturation.

Similarly to the CCA, the dendrogram showed a sharp distinction among the areas. The riparian forest was very dissimilar from the other

areas (approximately 63% dissimilarity), followed by the group composed of the SSF plots, and the vegetation corridor and inselberg plots, which did not separate clearly, showing that the inselbergs are more similar to the SSF patch than the riparian forest, even though they are geographically closer to the latter, indicating high beta diversity (Fig. 4). The cophenetic correlation coefficient was 0.73, indicating good reliability (low distortion) in the generation of the dendrogram from the raw data.

The differentiation between areas can also be observed by shared and exclusive species to each group. Only 12 species occurred in the four areas: *Albizia polycephala* (Benth.) Killip, *Casearia decandra* Jacq., *Casearia obliqua* Spreng., *Casearia sylvestris* Sw, *Copaifera langsdorffii*, *Cordia concolor* (Cham.) Kuntze, *Eugenia acutata* Miq., *Machaerium nyctitans* (Vell.) Benth., *Maytenus robusta*, *Myrcia splendens* (Sw.) DC., *Myrciaria floribunda*, and *Vochysia magnifica* Warm. In addition, a great number of species occurred in a single area: 58 in the SSF, seven in the inselberg,

TABLE 3: Indicator species of the riparian forest, inselberg, vegetation Corridor, and Seasonal Semideciduous Forest patch (SSF) in Coqueiral, Minas Gerais state, Brazil. VI - Indicator Value; p - significance.

 TABELA 3: Espécies indicadoras da Mata ciliar, *Inselberg*, Corredor e Fragmento de Floresta Estacional Semidecidual (FES). VI - valor de indicação; p - significância.

Environment/species	IV	p	Environment/species	IV	p
Riparian forest			Inselberg		
<i>Bauhinia longifolia</i> (Bong.) D.Dietr	66.7	0.0002	<i>Maytenus robusta</i> Reissek	86.9	0.0002
<i>Croton floribundus</i> Spreng.	59.3	0.0052	<i>Luehea divaricata</i> Mart.	80	0.0002
<i>Machaerium hirtum</i> (Vell.) Stellfeld	50	0.0022	<i>Myrciaria floribunda</i> (H.West ex Willd.) O.Berg	77.4	0.0002
<i>Celtis brasiliensis</i> (Gardn.) Planch.	50	0.0036	<i>Eugenia cerasiflora</i> Miq.	76.7	0.0002
<i>Senna macranthera</i> (Collad.) H.S.Irwin & Barneby	50	0.0036	<i>Zanthoxylum rhoifolium</i> Lam.	72.7	0.0004
<i>Cariniana legalis</i> (Mart.) Kuntze	50	0.0042	<i>Terminalia glabrescens</i> Mart.	52.9	0.0078
<i>Guapira hirsuta</i> (Choisy) Lundell	50	0.0042	<i>Luehea grandiflora</i> Mart. & Zucc.	48.7	0.0046
<i>Inga vera</i> Willd.	48.8	0.0052	<i>Cordia concolor</i> (Cham.) Kuntze	47.5	0.0206
<i>Cupania zanthoxyloides</i> Cambess.	44.2	0.0138	<i>Zanthoxylum fagara</i> (L.) Sarg.	47	0.0026
<i>Bathysa australis</i> (A.St.-Hil.) Benth. & Hook.f.	42.9	0.0362	<i>Ixora brevifolia</i> Benth.	47	0.0084
<i>Cabrlea canjerana</i> (Vell.) Mart.	41.7	0.0148	<i>Platypodium elegans</i> Vogel	47	0.01
<i>Endlicheria paniculata</i> (Spreng.) J.F.Macbr.	40.9	0.012	<i>Eriotheca candolleana</i> (K.Schum.) A.Robyns	45.5	0.0068
<i>Rollinia sylvatica</i> (A.St.-Hil.) Mart.	38.5	0.0128	<i>Vitex polygama</i> Cham.	43.2	0.0104
<i>Platycamus regnellii</i> Benth.	36.8	0.0156	<i>Casearia sylvestris</i> Sw.	43	0.0312
<i>Galipea jasminiflora</i> (A.St.-Hil.) Engl.	33.8	0.0198	<i>Myrciaria tenella</i> (DC.) O.Berg	38.9	0.0328
<i>Maclura tinctoria</i> (L.) D.Don ex Steud.	33.3	0.0328	<i>Casearia decandra</i> Jacq.	37.3	0.0476
<i>Ceiba speciosa</i> (A.St.-Hil.) Ravenna	33.3	0.0336	<i>Handroanthus serratifolius</i> (Vahl) S.O.Grose	36	0.035
			<i>Cedrela fissilis</i> Vell.	31.3	0.0362
Environment/species	IV	p	Environment/species	IV	p
Corridor			SSF Patch		
<i>Callisthene major</i> Mart.	63.2	0.0006	<i>Ocotea odorifera</i> (Vell.) Rohwer	72.7	0.0002
<i>Faramea nigrescens</i> Mart.	61.7	0.0026	<i>Mouriri glazioviana</i> Cogn.	57.9	0.0042
<i>Siphoneugena densiflora</i> O.Berg	59.8	0.001	<i>Duguetia lanceolata</i> A.St.-Hil.	56.3	0.0026
<i>Casearia arborea</i> (Rich.) Urb.	56.7	0.0068	<i>Aspidosperma olivaceum</i> Müll.Arg.	55.2	0.007
<i>Myrsine umbellata</i> Mart.	54.6	0.0048	<i>Jacaranda macrantha</i> Cham.	55	0.0032
<i>Tapirira obtusa</i> (Benth.) J.D.Mitch.	51.4	0.0208	<i>Tapirira guianensis</i> Aubl.	55	0.0034
<i>Dalbergia nigra</i> (Vell.) Allemão ex Benth.	44.4	0.0084	<i>Xylopia brasiliensis</i> Spreng.	55	0.0048
<i>Vismia brasiliensis</i> Choisy	44.4	0.0104	<i>Tachigali rugosa</i> (Mart. ex Benth.) Zarucchi & Pipoly	47.2	0.0114
<i>Protium spruceanum</i> (Benth.) Engl.	42.9	0.0274	<i>Calypttranthes clusiiifolia</i> O.Berg	46	0.0158
<i>Eugenia sonderiana</i> O.Berg	33.3	0.0268			
<i>Euplassa legalis</i> (Vell.) I.M.Johnst.	33.3	0.0302			
<i>Dalbergia villosa</i> (Benth.) Benth.	29	0.0486			

15 in the vegetation corridor, and 21 in the riparian forest, which represents 46.54% of the species total, stressing the difference in floristic composition among the environments.

DISCUSSION

The higher biomass allocation in number of individuals at the expense of basal area in inselbergs may be related to the higher light levels (BERTANI et al., 2001), the harsh environmental conditions, shallow soils, poor nutrient availability, and low water-holding capacity (GRÖGER and HUBER, 2007). In addition, despite the high recruitment in

inselbergs, mortality caused by abiotic factors is also high (FENNER, 1987). The low diversity in outcrops compared to other forest types was also report by Cao and Zhang (1997), who compared rain forests, montane forests, limestone outcrop forests in China. According to the authors, the low diversity of outcrops is due to the harsh environmental conditions, because outcrops are dry habitats with shallow soils, few species are adapted to these environments. In our study, the vegetation corridor had higher species richness than the inselberg, and this result is probably due to the fact that the former connects the riparian forest to the inselberg and thus has species from both formations. However,




FIGURE 3: Canonical correspondence analysis ordination diagram based on species density distributions of 40 plots (1–6: riparian forest; 7–15: vegetation corridor; 16–20: inselberg; 21–40: Seasonal Semideciduous Forest) in the municipality of Coqueiral, Minas Gerais, Brazil. The edaphic variables are represented by vectors.

FIGURA 3: Diagrama de ordenação de 40 parcelas: (1-6) mata ciliar; (7-15) corredor; (16-20) *inselberg*; (21-40) Fragmento de Floresta Estacional Semidecidual, no município de Coqueiral, pela análise de correspondência canônica, baseada na distribuição da densidade das espécies. As variáveis edáficas estão representadas por vetores.




FIGURE 4: Floristic similarity dendrogram of four microenvironments sampled in the municipality of Coqueiral, Minas Gerais, Brazil.

FIGURA 4: Dendrograma de similaridade florística de quatro microambientes amostrados no município de Coqueiral - MG.

the high equability observed in all environments demonstrates the lack of ecological dominance, indicating that the species are evenly represented in each studied area (KANIESKI et al., 2010).

Environmental heterogeneity results in high species turnover, which increases β diversity, (POREMBSKI, 2007). Habitat heterogeneity is the factor most widely associated with the determination of beta diversity (MORENO and HALFFTER, 2001), because beta diversity results from the species responses to variations between sites and, thus can be understood as the degree of habitat specialization of (SHMIDA and WILSON, 1985; JANKOWSKI et al., 2009; MELO et al. 2009). *Copaifera langsdorffii* was the only species with a high importance value in all areas, showing that adaptation varies even in geographically close areas, because plant species differ in their nutrient requirements and microclimate (DENSLOW, 1980). *Copaifera langsdorffii* is distributed throughout the Brazilian territory in open areas and forest formations with well drained soils (FREITAS and OLIVEIRA, 2002; FAGUNDES et al., 2007; VEIGA JUNIOR and PINTO, 2002). The species *Terminalia glabrescens*, which had the second IV in the inselberg is associated with well drained soils (FAGUNDES et al., 2007).

Some species were among the most important in each environment and also had the highest indicator values such as: *Ocotea odorifera* (SSF plots); *Croton floribundus* (riparian forest); *Callisthene major*, *Casearia arborea*, and *Faramea nigrescens* (vegetation corridor); and *Maytenus robusta* and *Myrciaria floribunda* (inselberg). *Ocotea odorifera* is listed as endangered in IBAMA Red List (IBAMA, 1992) and is among the most important tree species in ombrophilous forests (NEGRELLI and LEUCHTENBERG, 2001). In addition, *C. major* is more often found in well-drained environments (CARVALHO et al., 2005; IMAÑA-ENCINAS et al., 2007). Conversely, *M. robusta* is a heliophyte occurring in more open vegetation (LORENZI, 1998), which explains its higher indicator values in the inselberg.

Several studies have shown that the vegetation mosaic of forest patches is related to the physicochemical characteristics of soil (OLIVEIRA FILHO et al., 1994; RODRIGUES et al., 2007; CAMARGOS et al., 2008), including a few studies on rocky outcrops (CONCEIÇÃO et al., 2007; GRÖGER and HUBER 2007). Groger and Huber (2007) found that soil depth gradients and water

availability were the main factors determining the floristic composition on a rocky outcrop in the Guayana Escudo. Conceição et al. (2007) compared four areas in Chapada Diamantina, Brazil and found that the structure and floristic composition were related to soil characteristics rather than the distance between the sites, indicating that the variation in soil characteristics affected the plant community studied.

The value for axis 1 can be considered high (TER BRAAK, 1987; FELFILI and REZENDE, 2003) and the low variance explained by the first two axes is a common feature of vegetation data, which does not affect the significance of the species-environment relationships (TER BRAAK, 1987). Clay, which correlated mostly with SSF plots, is responsible for the adsorption of both cations and anions and holding bases and water, causing the soils to be wet, muddy, and more compressed (RIZZINI, 1997). However, calcium and base saturation, which correlated mostly with riparian forest plots, are indicative of low acidity (CAMARGOS et al., 2008).

The greatest dissimilarity of the riparian forest to the vegetation corridor, inselberg, and semideciduous forest patch was probably due to the stronger floristic links of riparian forests with wetter forests especially regarding high soil moisture content (OLIVEIRA FILHO and RATTER, 2009). These results also indicate that even communities situated in close areas can be floristically and structurally different (FERREIRA JUNIOR et al., 2008).

CONCLUSION

The harsh environmental conditions in the inselberg, which are conditioned by the shallow soils, affect its species richness and community structure. Inselbergs had the lowest species richness and a greater number of individuals of smaller diameter, resulting in greater differences in species composition and community structure between the inselberg and the surrounding matrix and consequently higher β diversity. The results showed high species turnover among areas, indicating that even though the areas are geographically close, the occurrence of inselbergs directly affects the variation in species composition and community structure among areas.

ACKNOWLEDGEMENTS

We are thankful to Minas Gerais State Research Foundation (FAPEMIG), to Universidade Federal de Lavras (UFLA) for logistic support and Coordination for the Improvement of Higher Education Personnel (CAPES) for providing a research fellowship to the first author.

REFERENCES

- BENITES, V. M. et al. Soils associated with rock outcrops in the Brazilian mountain ranges Mantiqueira and Espinhaço. **Revista Brasileira de Botânica**, São Paulo, v. 30, n. 4, p.569-577, out.-dez. 2007.
- BERTANI, D. F. et al. Análise temporal da heterogeneidade florística e estrutural em uma floresta ribeirinha. **Revista Brasileira de Botânica**, São Paulo, v. 24, n. 1, p.11-23, mar. 2001.
- BROWER, J. E.; ZAR, J. H. **Field and laboratory methods for general ecology**. Dubuque, W.M.C. Brow, 1984.
- BUCKLEY, L. B.; JETZ, W. Linking global turnover of species and environments. *Proceedings of The National Academy of Sciences of the USA*, v. 105, p. 17836-17841, 2008.
- CAIAFA, A.; SILVA, A. F. Structural analysis of vegetation on a highland granitic rock outcrop in Southeast Brazil. **Revista Brasileira de Botânica**, São Paulo, v. 30, n.4, p.657-664, out.-dez. 2007.
- CAMARGOS, V. L. et al. Influência de fatores edáficos sobre variações florísticas na floresta Estacional Semidecídua no entorno da Lagoa Carioca, Parque Estadual do Rio doce, MG, Brasil. **Acta Botanica Brasilica**, São Paulo, v. 22, n. 1, p. 75-84, 2008.
- CAO, M.; ZHANG, J. Tree species diversity of tropical forest vegetation in Xishuangbanna, SW China. **Biodiversity and Conservation**, Amsterdam, v. 6, p. 995-1006, 1997.
- CARVALHO, D.A. et al. Distribuição de espécies arbóreo-arbustivas ao longo de um gradiente de solos e topografia em um trecho de floresta ripária do Rio São Francisco em Três Marias, MG, Brasil. **Revista Brasileira de Botânica**, São Paulo, v. 28, p. 329-345, 2005.
- COLWELL, R. K. EstimateS: Statistical estimation of species richness and shared species from samples. Version 9. Persistent URL <purl.oclc.org/estimates>.
- CONCEIÇÃO, A. A.; PIRANI, J. R.; MEIRELLES, S. T. Floristics, structure and soil of insular vegetation in four quartzite-sandstone outcrops of “Chapada Diamantina”, Northeast Brazil. **Revista Brasileira de Botânica**, São Paulo, v. 30, n.4, p. 641-656, 2007.
- DENSLOW, J. S. Gap partitioning among tropical rainforest trees. **Biotropica**, Marburg, v. 12, p. 47-51, 1980.
- DUFRÊNE, M.; LEGENDRE, P. Species assemblages and indicator species: the need for a flexible asymmetrical approach. **Ecological Monographs**, Washington, v. 67, p. 345-366, 1997.
- EMBRAPA – Empresa Brasileira de Pesquisa Agropecuária. **Manual de métodos de análise de solo**. Rio de Janeiro, Empresa Brasileira de Pesquisa Agropecuária, Centro Nacional de Pesquisa de Solos, 1997.
- FAGUNDES, L. M. et al. Florística e estrutura do estrato arbóreo de dois fragmentos de florestas decíduas às margens do rio Grande, em Alpinópolis e Passos, MG, Brasil. **Acta Botanica Brasilica**, São Paulo, v. 21, n. 1, p. 65-78, 2007.
- FELFILI, J. M.; REZENDE, R. P. **Conceitos e métodos em fitossociologia**. Brasília: Universidade de Brasília, Departamento de Engenharia Florestal, Comunicações técnicas florestais 5, 2003, 68p.
- FENNER, M. Seedlings. **New Phytologist**, Oak Ridge, v. 106, p. 35-47, 1987.
- FERREIRA-JÚNIOR, E. V. et al. Composição, diversidade e similaridade florística de uma floresta tropical semidecídua submontana em Marcelândia – MT. **Acta amazônica**, Manaus, v. 38, n. 4, p. 673-680, 2008.
- FINA, B. G.; MONTEIRO, R. Estudo da estrutura da comunidade arbustivo-arbórea de uma área de cerradão, município de Pirassununga (SP). **Neotropical Biology and Conservation**, São Leopoldo, v. 4, n. 1, p. 40-48, jan.-abr. 2009.
- FRANÇA, F.; MELO, E.; SANTOS, C. C. Flora de Inselbergs da região de Milagres, Bahia, Brasil: I. Caracterização da vegetação e lista de espécies de dois inselbergs. **Sitientibus**, Feira de Santana, n. 17, p. 163-184, jul.-dez. 1997.
- FREITAS, C. V.; OLIVEIRA, P. E. Biologia reprodutiva de *Copaifera langsdorffii* Desf. (Leguminosae, Caesalpinioideae). **Revista Brasileira de Botânica**, São Paulo, v. 25, n. 3, p. 311-321, set. 2002.
- GROGER, A.; HUBER, O. Rock outcrop habitats in the Venezuelan Guayana lowlands: their main vegetation types and floristic components. **Revista Brasileira de Botânica**, São Paulo, v. 30, n. 4,

p. 599-609, out.-dez. 2007.

IBAMA. **Lista Oficial das Espécies da Flora Brasileira Ameaçadas de Extinção**. Diário Oficial. Portaria 006/92-N de 15 de janeiro de 1992.

IMAÑA-ENCINAS J.; MACED, L. A.; PAULA, J. E. Florística e fitossociologia de um trecho da Floresta Estacional Semidecidual na área do ecomuseu do cerrado, em Pirenópolis – Goiás. **Cerne**, Lavras, v. 13, n. 3, p. 308-320, jul.-set. 2007.

JANKOWSKI, J. E. et al. Beta diversity along environmental gradients: implications of habitat specialization in tropical montane landscapes. **Journal of Animal Ecology**, London, v. 78, p. 315-327, 2009.

KANIESKI, M. R.; ARAÚJO, A. C. B.; LONGHI, S. J. Quantificação da diversidade em Floresta Ombrófila Mista por meio de diferentes Índices Alfa. **Scientia Forestalis**, Piracicaba, v. 38, n. 88, p. 567-577, dez. 2010.

LORENZI, H. **Árvores brasileiras**. Editora Plantarum, Nova Odessa, 1998, 222 p.

MCCUNE, B.; MEFFORD, M. J. **PC-ORD version 4.0, multivariate analysis of ecological data**, Users guide. Glaneden Beach: MjM Software Design, 1999, 237 p.

MCINTYRE, N. E.; RANGO, J.; FAGAN, W. F.; FAETH, S. H. Ground arthropod community structure in a heterogeneous urban environment. **Landscape and Urban Planning**, v. 52, p. 257-274, 2001.

MELO, A. S.; RANGEL, T. F. L. V. B.; DINIZ-FILHO, J. A. F. Environmental drivers of beta-diversity patterns in New-World birds and mammals. **Ecography**, Denmark, v. 32, p. 226-236, 2009.

MICHAEL, D. R.; CUNNINGHAM, R. B.; LINDENMAYER, D. B. The social elite: Habitat heterogeneity, complexity and quality in granite inselbergs influence patterns of aggregation in *Egernia striolata* (Lygosominae: Scincidae). **Austral Ecology**, Malden, v. 35, n. 8, p. 862–870, 2010.

MINISTÉRIO DO MEIO AMBIENTE. **Biodiversidade Brasileira: Avaliação e Identificação de Áreas e Ações Prioritárias para Conservação, Utilização Sustentável e Repartição de Benefícios da Biodiversidade Brasileira**. Brasília, MMA/SBF, 2002.

MORENO, C. E.; HALFFTER, G. Spatial and temporal analysis of alfa, beta and gama diversities of bats in a fragmented landscape. **Biodiversity and Conservation**, Amsterdam, v.10, p. 367-382, 2001.

MUELLER-DOMBOIS, D.; ELLEMBERG, H. **Aims and Methods of Vegetation Ecology**, Bélgica. New York, John Wiley & Sons, 1974.

NEGRELLE, R. R. B.; LEUCHTENBERGER, R. Composição e estrutura do componente arbóreo de um remanescente de floresta ombrófila mista. **Floresta**, Curitiba, v. 31, 1974.

OLIVEIRA FILHO, A. T. et al. Effect of floating regime and understorey bomboos on the physiomy and tree species composition of a tropical semideciduous forest in southeastern Brazil. **Vegetatio**, v.113, p. 99-124, 1994.

OLIVEIRA FILHO, A. T.; RATTER, J. A. Padrões florísticos das Matas ciliares da Região do Cerrado e a Evolução das Paisagens do Brasil Central durante o Quaternário Tardio. In: RODRIGUES, R. R.; LEITÃO FILHO, H. F. (eds). **Matas ciliares: conservação e recuperação**: 73-89. Second edition. São Paulo, Editora da Universidade de São Paulo, Fapesp, 2009.

POREMBSKI, S. et al. Diversity and ecology of saxicolous vegetation mats on inselbergs in the Brazilian Atlantic Rainforest. **Biodiversity Research**, Vancouver, v. 4, p. 107-119, 1998.

POREMBSKI, S. Tropical inselbergs: habitat types, adaptative strategies and diversity patterns. **Revista Brasileira de Botânica**, São Paulo, v. 30, n. 4, p. 579-586, out.-dez. 2007.

POREMBSKI, S.; BARTHLOTT, W. **Inselbergs: Biotic Diversity of Isolated Rock Outcrops in Tropical and Temperate Regions**. Berlin: Ecological Studies 146, Springer-Verlag, 2000.

RIBEIRO, K.T.; MEDINA, B. M. O.; SCARANO, F. R. Species composition and biogeographic relations of the rock outcrop flora on the high plateau of Itatiaia, SE-Brazil. **Revista Brasileira de Botânica**, São Paulo, v. 30, n. 4, p. 623-639, 2007.

RIZZINI, C. T. **Tratado de fitogeografia do Brasil: aspectos ecológicos, sociológicos e florísticos**. Âmbito Cultural Edições Ltda, Rio de Janeiro, 1997.

RODRIGUES, L. A. et al. Efeitos de solos e topografia sobre a distribuição de espécies arbóreas em um fragmento de floresta estacional semidecidual, em Luminárias, MG. **Revista Árvore**, Viçosa, v. 31, n. 1, p. 25-35, 2007.

SCARANO, F.R. Rock outcrop vegetation in Brazil: a brief overview. **Revista Brasileira de Botânica**, São Paulo, v. 30, n. 4, p. 561-568, out.-dez. 2007.

SHMIDA, A.; WILSON, M. V. Biological determinants of species diversity. **Journal of**

- Biogeography**, Malden, v.12, p.1-20, 1985.
- SHEPHERD, G.J. **Fitopac 1.6**: Manual do usuário. Unicamp, 2006.
- TER BRAAK, C. J. F. The analysis of vegetation-environment relationship by canonical correspondence analysis. **Vegetatio**, Bélgica, v. 69, p. 69-77, 1987.
- TER BRAAK, C. J. F. **CANOCO- A FORTRAN program for canonical community ordination by (Partial) (Detrended) (Canonical) correspondence analysis and redundancy analysis, version 2.1**. Technical report LWA-88-2, TNO, Wageningen, Intitute of Applied Computer Science, 1988.
- TER BRAAK, C. J. F. Ordination. In: JONGMAN, R. H. G.; Ter Braak, C. J. F.; VAN TONGEREN, O. F. R. (eds). **Data analysis in community and landscape ecology**: 91-173. Cambridge University Press, Cambridge, 1995.
- VEIGA JUNIOR, V.F.; PINTO, A. C. O gênero *Copaifera* L. **Química Nova**, São Paulo, v. 25, n. 2, p. 273-286, 2002.
- VIANA, V. M.; PINHEIRO, L. A. F. V. Conservação da biodiversidade em fragmentos florestais. **Série Técnica IPEF**, Piracicaba, v. 12, n. 32, p. 25-42, dez. 1998.
- VILELA, E. A.; RAMALHO, M. A. P. Análise das temperaturas e precipitações pluviométricas de Lavras - MG. **Ciência e Prática**, Bebedouro, v. 3, p. 71-79, 1979.
- WHITTAKER, R. J.; WILLIS, K. J.; FIELD, R. Scale and species richness: towards a general, hierarchical theory of species diversity. **Journal of Biogeography**, Malden, v.28, p. 453-470, 2001.
- ZAR, J. H. **Biostatistical analysis**. 3a ed. Prentice Hall, New Jersey, 1996.