

## PERFORMANCE AND QUANTITATIVE AND QUALITATIVE CARCASS TRAITS IN PIGS FED ON CANOLA MEAL<sup>1</sup>

LINA MARÍA PEÑUELA SIERRA<sup>2</sup>, IVAN MOREIRA<sup>3</sup>, PAULO LEVI DE OLIVEIRA CARVALHO<sup>4\*</sup>, ADRIANA GOMEZ GALLEGOS<sup>3</sup>, LAURA MARCELA DIAZ HUEPA<sup>3</sup>, CLODOALDO DE LIMA COSTA FILHO<sup>3</sup>

**ABSTRACT** - Two experiments were performed for the purpose of determining the nutritional value and assess the performance and carcass traits in pigs fed on canola meal (*Brassica napus*). Experiment I consisted of a total digestibility trial conducted with 14 castrated commercial crossbred male pigs with  $60.69 \pm 4.26$  kg initial live weight. The treatments consisted of a control diet based on maize and soybean meal and another diet containing canola meal (control + 25% canola meal). Each pig was an experimental unit, totalling seven experimental units per diet. The digestible energy and the metabolizable energy of canola meal used was 2,999 and 2,730 kcal.kg<sup>-1</sup>, respectively. These values were used to formulate the diets of the performance experiment. Experiment II included 50 crossbred pigs with an average initial live weight of  $29.90 \pm 1.16$  kg and an average final live weight of  $60.33 \pm 3.38$  kg during the growing phase, and an average initial live weight of  $60.37 \pm 1.6$  kg and an average final live weight of  $90.37 \pm 3.19$  kg during the finishing phase. The experimental design consisted of randomized blocks, with four treatments (6, 12, 18, and 24% canola meal), 10 repetitions, and one pig per experimental unit. The control diet was formulated with 0% canola meal. The data were assessed using ANOVA test and the averages of the treatments were compared using Dunnett's test and regression analysis at a 5% probability level. The results suggest that it is possible to use up to 24% canola meal in the diets for pigs during the growing and finishing phases without affecting the performance and carcass traits.

**Keywords:** Protein food. Canola meal. Pigs.

## DESEMPENHO E CARACTERÍSTICAS QUANTITATIVAS E QUALITATIVAS DE CARÇAÇA DE SUÍNOS ALIMENTADOS COM FARELO DE CANOLA

**RESUMO** - Foram conduzidos dois experimentos com o objetivo de determinar o valor nutricional e avaliar o desempenho e características da carcaça de suínos alimentados com farelo de canola (*Brassica napus*). No Experimento I, foi conduzido um ensaio de digestibilidade total com 14 suínos mestiços de linhagem comercial, machos castrados, com  $60,69 \pm 4,26$  kg de PV inicial. O alimento teste farelo de canola (FC), substituiu em 25% a ração referência à base de milho e farelo de soja. Cada suíno constituiu uma unidade experimental, totalizando sete unidades experimentais por ração. Os valores de energia digestível (ED) e energia metabolizável (EM) na matéria natural para o FC foram de 2995 kcal/kg e 2796 kcal/kg, respectivamente. No Experimento II, foram utilizados 50 suínos mestiços, com peso vivo inicial de  $29,90 \pm 1,16$  kg e final de  $60,33 \pm 3,38$  kg, na fase de crescimento, e peso vivo inicial de  $60,37 \pm 1,6$  kg e final de  $90,37 \pm 3,19$  kg na fase de terminação. Foi utilizado o delineamento de blocos inteiramente casualizados, com quatro tratamentos, (6, 12, 18 e 24% de FC), com 10 repetições e um suíno por unidade experimental. Adicionalmente, foi formulada uma ração testemunha, contendo 0% de FC. Os dados foram submetidos à análise de variância e adicionalmente foi aplicado o Teste de Dunnett, e análise de regressão, a 5% de probabilidade. Os resultados sugerem que é possível incluir até 24% de FC nas dietas de suínos, em crescimento e terminação, sem prejudicar o desempenho e as características da carcaça.

**Palavras-chave:** Alimento protéico. Farelo de Canola. Suínos.

\*Corresponding author

<sup>1</sup>Received for publication in 02/18/2015; accepted in 02/18/2016.

Paper extracted from the doctoral thesis of the first author.

<sup>2</sup>Department of Veterinary and Animal Science, Universidad del Tolima, Ibagué, TOL, Colombia; linapenuela@gmail.com.

<sup>3</sup>Departament of Animal Science, Universidade Estadual de Maringá, PR, Brazil; imoreirai@gmail.com, adritonyv@yahoo.es, lemondedelaura28@hotmail.com, clodoaldocostafilho@yahoo.com.br.

<sup>4</sup>Department of Animal Science, Universidade Estadual do Oeste do Paraná, Marechal Cândido Rondon, PR, Brazil; paulolevi@yahoo.com.br.

## INTRODUCTION

Brazil is the fourth world producer and exporter of pigs and, in 2010, per capita pig consumption was 14.8 kg (ABIPECS, 2011). Soybean meal (*Glycine max*) is the most used protein source to formulate diets for pigs (CROMWELL, 2010). However, due to its high cost, it is necessary to evaluate other protein sources, such as canola (*Brassica napus*), cotton (*Gossypium herbaceum*), sunflower (*Helianthus annuus*), and peanut (*Arachis hypogaea*).

Canola is a rapeseed cultivar (*Brassica napus*) that contains low levels of glucosinolates in the seed ( $<3 \mu\text{g g}^{-1}$ ) and erucic acid in the oil ( $<2\%$ ) (Bell, 1993). In Brazil, its cultivation has been increasing as an option for the winter harvest and the production of oil rich in unsaturated fatty acids (SANTOS; BASSO, 1990).

Canola meal (CM) is a by-product of oil extraction and has a great content of sulphur amino acids, ether extract, fibres, calcium, phosphorus (Keith & Bell, 1991), and vitamins of the B complex (niacin, thiamine, riboflavin, folic acid, and biotin), compared with soybean meal (BELL, 1993). However, it has limiting factors, such as the production of metabolites of glucosinolates, trypsin inhibitors, phytic acid, phenolic compounds, and tannins (TESKEREDZIC et al., 1995).

The goal of the present study was to assess the nutritional value of CM in diets for growing-finishing pigs and its effects on performance and carcass traits.

## MATERIAL AND METHODS

The experiments were carried out in the swine sector of Iguatemi Experimental Farm, Agricultural Sciences Centre of the State University of Maringá, State of Paraná, Brazil (23°21' S, 52°04' W; 564 m altitude).

Were conducted two experiments. The first assessed the total digestibility of CM and the second assessed the performance of growing-finishing pigs and carcass traits. The CM was acquired in an agroindustrial firm of the State of Paraná, Brazil. The experimental procedures and the use of animals were previously approved by the Ethics Committee of the State University of Maringá, Approval Certificate No. 136/2010.

The digestibility experiment was carried out from March to April 2010 assessing 14 castrated commercial crossbred male pigs with initial live weight of  $60.69 \pm 4.26$  kg. The pigs were housed in individual metabolism crates under controlled environment. The digestibility trial lasted 15 days, ten for adaptation and five for collecting faeces and urine. The control diet consisted of maize (72.97%),

soybean meal (24.45%), salt (0.570%), calcium carbonate (0.635%), dicalcium phosphate (0.875%), and mineral-vitamin supplement (0.50%). These components were formulated according to the requirements proposed by Rostango et al. (2005). Were used two diets (treatments): control; and control (75%) with addition of CM (25%) performing seven repetitions per treatment.

Provision of diets and collection of faeces and urine were performed according to the method proposed by Sakomura and Rostagno (2007). During the collection period, feed intake was calculated based on the metabolic weight ( $\text{kg}^{0.75}$ ) of each pig and the average consumption recorded during the adaptation period (approximately seven days).

Feed was provided at 8 a.m. (55%) and 3 p.m. (45%). It was moistened with water (20%) to avoid waste, reduce exposure to dust, and improve consumption. After each meal, were provided 3 mL of water/ $\text{g}^{-1}$  feed.

To set the beginning and the end of the faeces collection procedure, were used 3% iron oxide ( $\text{Fe}_3\text{O}_2$ ) as faecal marker (SAKOMURA; ROSTAGNO, 2007) and faeces were collected once a day, packed in plastic bags and stored in a freezer ( $-18^\circ\text{C}$ ). Subsequently, each sample was homogenized and dried (approximately 350 g) in a forced-ventilation oven ( $55^\circ\text{C}$ ) and ground in a blade mill (1 mm sieve). Urine was collected in plastic buckets with 20 mL of HCl solution (1:1) to avoid bacterial growth and nutrient losses by volatilization. The analyses of feed and faeces were performed according to the procedures described by Silva and Queiroz (2002). Total energy values were determined by adiabatic calorimetry (Parr Instrument Compant, Illinois, USA). The digestibility coefficients of dry matter (DMDC), energy (EDC), protein (PDC), and organic matter (OMDC) were calculated according to the method proposed by Moreira et al. (1994).

Experiment II assessed the performance during the growing (30-60 kg) and finishing (60 to 90 kg) phases and was carried out from July 2010 to February 2011. During the experiment, the minimum average temperature was  $18.8 \pm 2.32^\circ\text{C}$  and the maximum  $29.1 \pm 4.11^\circ\text{C}$ . Were used 50 commercial crossbred pigs whose average initial live weight was  $29.90 \pm 1.16$  kg and the average finishing live weight was  $60.33 \pm 3.38$  kg during the growing phase. During the finishing phase, the average initial live weight was  $60.37 \pm 1.46$  and the average final live weight was  $90.37 \pm 3.19$  kg. The pigs were housed in brick sheds divided into two sectors, each with 10 pens ( $7.60 \text{ m}^2$  each) separated by a central corridor. Each pen had pacifier-type drinkers at the back and single feeders at the front, which provided free access to feed and water. The diets and water were provided *ad libitum* throughout the experiment. The treatments consisted of four diets (6, 12, 18, and 24% CM) and a control diet (0% CM). The diets (Tables 1

and 2) were formulated as recommended by Rostagno et al. (2005) for growing-finishing pigs. We assessed the ileal digestibility to determine the digestible amino acids of the CM and the results were used to formulate the diets. We determined the levels of protein, phosphorus, and calcium in maize and soybean meal samples at the Laboratory of Food and Animal Nutrition Analysis, State University of Maringá, Paraná, Brazil. The metabolizable energy values were those cited by Rostagno et al. (2005). The values of digestible

amino acids of maize and soybean meal were calculated using true amino acid digestibility coefficients reported by Rostagno et al. (2005).

The experimental design consisted of randomized blocks with four treatments (6, 12, 18, and 24% CM), 10 repetitions, and one pig per experimental unit. The pigs were weighed at the beginning and at the end of the experiment. We determined the daily feed intake, the daily weight gain, and calculated the feed:gain ratio of each pig.

**Table 1.** Centesimal and chemical composition of diets containing different levels of canola meal (CM) for growing pigs feeding (30 a 60 kg).

Item	Canola Meal levels (%)				
	0	6	12	18	24
Corn	76.57	72.10	67.62	63.15	58.68
Soybean meal	20.14	18.12	16.09	14.07	12.05
Canola meal	0.000	6.000	12.00	18.00	24.00
Dicalc. Phosp.	1.475	1.433	1.391	1.349	1.308
Limestone	0.484	0.437	0.391	0.345	0.298
Common salt	0.400	0.400	0.400	0.400	0.400
Soybean oil	0.225	0.837	1.449	2.062	2.675
Min. vit. premix <sup>†</sup>	0.300	0.300	0.300	0.300	0.300
Growth prom. <sup>††</sup>	0.005	0.005	0.005	0.005	0.005
L-Lys. HCL, 99%	0.278	0.268	0.257	0.246	0.236
DL-Meth. 99%	0.071	0.056	0.039	0.023	0.006
L-Thre. 98%	0.054	0.053	0.051	0.048	0.046
Calculated values <sup>†††</sup>					
Met. Eng. Kcal/kg	3230	3230	3230	3230	3230
Dig. lysine, %	0.89	0.89	0.89	0.89	0.89

<sup>†</sup> Vitamin and mineral premix for growing pigs (kg of product): vitamin A, 50 000 UI; D3, 10 000 UI; E, 160 mg; K3, 12 mg; B1, 12 mg; B2, 20 mg; B6, 12 mg; B12, 100 µg; pholic acid, 2.4 mg; niacin, 140 mg; pantothenic acid, 88 mg; biotin, 0.4 mg; choline, 1248 g; iron, 800 mg; copper, 800 mg; cobalt, 3.2 mg; manganese, 220 mg; zinc, 11.15 mg; selenium, 1.2 mg; Iodine, 7.2 mg.  
<sup>††</sup>Leucomycin 30%. <sup>†††</sup> Calculate based on Rostagno et al. (2005).

**Table 2.** Centesimal and chemical composition of diets containing different levels of canola meal (CM) for finishing pigs feeding (60-90 kg).

Item	Canola Meal levels (%)				
	0	6	12	18	24
Corn	81.58	77.59	75.35	73.11	70.88
Soybean meal	16.03	13.53	9.234	4.937	0.641
Canola meal	0.000	6.000	12.00	18.00	24.00
Dicalc. Phosp.	1.043	1.004	0.978	0.952	0.926
Limestone	0.445	0.398	0.355	0.312	0.269
Common salt	0.400	0.400	0.400	0.400	0.400
Soybean oil	0.105	0.700	1.222	1.745	2.268
Min. vit. premix <sup>†</sup>	0.150	0.150	0.150	0.150	0.150
Growth prom. <sup>††</sup>	0.005	0.005	0.005	0.005	0.005
L-Lys. HCL, 99%	0.199	0.194	0.242	0.289	0.337
DL-Meth. 99%	0.015	0.000	0.000	0.000	0.000
L-Thre. 98%	0.029	0.028	0.050	0.073	0.096
L-Tryptophan, 98%	0.004	0.005	0.014	0.024	0.034
Calculated values <sup>†††</sup>					
Met. Eng. Kcal/kg	3250	3250	3250	3250	3250
Dig. lysine, %	0.68	0.68	0.68	0.68	0.68

<sup>†</sup>Vitamin and mineral premix for finishing pigs, (kg of product): Vit A, 50.000 UI; Vit D3, 10.000 UI; Vit E, 160mg; Vit K3, 12 mg; Vit B1, 12 mg; Vit B2, 20 mg; Vit B6, 12 mg; Vit B12, 100 µg; pholic acid, 2.4 mg; niacin, 140 mg; pantothenic acid, 88mg; biotin, 0.4mg; choline, 1.248g; iron, 800mg; copper, 800mg; cobalt, 3.2mg; manganese, 220mg; zinc, 11.150mg; selenium, 1.2mg; iodine, 7.2mg. <sup>††</sup>Leucomycin 30%; <sup>†††</sup>Calculate based on Rostagno et al. (2005).

**Table 3.** Performance of growing (30-60 kg) and finishing (60-90 kg) pigs, fed on diets with canola meal.

Item	Canola Meal levels (%)					Media±EP <sup>†</sup>	CV <sup>††</sup>	Lin <sup>†††</sup>	Qua <sup>††††</sup>
	Growing								
	0	6	12	18	24				
DFI, kg	2.01	1.95	2.02	2.01	2.02	2.00±0.21	10.66	NS	NS
DWG, kg	0.89	0.90	0.89	0.97	0.92	0.92±0.11	11.87	NS	NS
Feed:gain	2.25	2.18	2.28	2.08	2.19	2.20±0.19	9.05	NS	NS
NUP, mg/dl	14.04	14.91	15.35	13.87	16.15	14.84±3.87	26.45	NS	NS
BT, mm	8.10	8.80	9.10	8.30	9.11	8.82±1.85	19.13	NS	NS
LD, mm	42.20	45.80	45.50	45.90	44.56	45.46±5.19	10.10	NS	NS
	Finishing								
DFI, kg	2.67	2.60	2.65	2.53	2.68	2.63±0.14	10.88	NS	NS
DWG, kg	0.99	0.95	0.94	0.87	0.93	0.94±0.14	16.08	NS	NS
Feed:gain	2.69	2.76	2.84	2.93	2.90	2.82±0.31	12.19	NS	NS
NUP, mg/dl	13.96	13.72	13.96	12.18	12.82	13.33±2.71	12.34	NS	NS
BT, mm	12.10	13.20	12.10	12.40	13.10	12.58±2.86	17.82	NS	NS
LD, mm	44.40	46.40	46.40	47.40	48.50	46.62±9.59	13.82	NS	NS

<sup>†</sup>Standard Error; <sup>††</sup>CV = Coefficient of variation; <sup>†††</sup>Linear Effect of Levels Canola Meal; <sup>††††</sup> Quadratic Effect of Levels Canola Meal; NS = non-significat; DFI = Daily feed intake; DWG = Daily weight gain; BT = Backfat thickness; LD = Loin depth.

However, Moreira et al. (1996) worked with growing-finishing pigs and concluded that there was a reduction in the performance of the pigs according to increased CM levels in the diets. These authors linked this behaviour to the fact that CM contains high levels of crude fibre that can impair the performance. Thacker and Newkirk (2005) worked with growing-finishing pigs (28.7 to 107.4 kg of live weight) and observed that there was a reduction in daily weight gain and feed:gain ratio during the growing phase when they completely replaced soybean meal with CM as a protein source. In the finishing phase, total replacement did not affect daily feed intake, daily weight gain, and feed:gain ratio.

On the other hand, Roth-Maier et al. (2004) observed an improvement trend in the yield variables of growing animals with increased levels of CM inclusion (0-26%). However, there was a linear decrease in daily gain weight as the CM level increased (0-17%). Gomes et al. (1998) observed that it was possible to include 15% of CM in the diet of finishing pigs, whereas Brand et al. (2001) concluded that the inclusion of 24% for pigs with 25 to 84 kg live weight did not affect the performance. The experiment carried out by Zanotto et al. (2009) showed a detrimental effect on the feed:gain ratio during the growing phase and a limitation in the daily feed intake during the finishing phase when high levels of CD (replacement level above 40% of soybean meal) were included to replace soybean meal.

Results obtained by Seneviratne et al. (2010) were similar to those of the present study. They assessed pigs fed on diets containing a maximum of 23% CD during the growing phase and found no differences. However, these authors reported that,

when they increased the level of CM inclusion in finishing pigs, the daily weight gain decreased linearly and feed:gain ratio worsened. With regard to the quantitative carcass traits (Table 4), none of the variables was influenced by the addition of CD ( $p \geq 0.05$ ).

Dunnett's test indicated that there was no difference ( $p \geq 0.05$ ) between the levels of CM inclusion and the control diet (0% CM) for most variables. However, for the variables 'hot carcass weight' and 'cold carcass weight', the values were lower ( $p \leq 0.05$ ) between inclusion levels of 12, 18, and 24% for hot carcass weight and between 18 and 24% for cold carcass weight compared with the control diet. This result is due to the lower weight of these animals at sacrifice. However, special attention should be given to these variables. Losses during cooling represent an important value related to the percentage of carcass water loss in the refrigerator (BRIDI; SILVA, 2009). Therefore, these results indicate that up to 24% of CM inclusion in the diets does not affect the quantitative carcass traits in pigs.

The results obtained in the present study are similar to those obtained by Gomes et al. (1998), Roth-Maier et al. (2004), and Thacker and Newkirk (2005), which did not show any effect of the CM level replacing soybean meal on the carcass traits assessed in growing-finishing pigs. Rojo et al. (2001) assessed the potential of partial (50%) or total (100%) replacement of soybean meal with CM and the effects on pigs sacrificed at 100 kg live weight. They reported that carcass yield, meat percentage, and backfat thickness were not affected by the treatments. However, Brand et al. (2001) assessed the carcasses of lightweight pigs (up to 85 kg live weight) including high CM levels. They found a

reduction in backfat thickness, increase in the percentage of carcass meat, and reduction in the carcass performance.

In addition, Zanotto et al. (2009) reported that the inclusion of CM in the diet resulted in carcasses

with lower meat yield. With the exception of colour (a\*), lightness (L\*), and tonality (b\*) of the longissimus muscle, none of the other qualitative meat traits (Table 5) were influenced by the inclusion of CM ( $p \geq 0.05$ ).

**Table 4.** Effect of diets with different levels of canola meal inclusion (CM) on quantitative carcass traits of pigs in finishing (60 to 90 kg).

Variables	Inclusion levels of MC (%)					Mean±SE <sup>†</sup>	CV <sup>††</sup>	Lin <sup>§</sup>	Qua <sup>¶</sup>
	0	6	12	18	24				
Fasting losses, %	3.98	3.66	4.06	4.08	4.02	3.96±1.37	29.46	NS	NS
Hot carcass weight, kg	72.93	71.89	71.30*	69.47*	70.65*	71.25±3.23	1.48	NS	NS
Hot carcass yield, %	81.81	82.47	82.20	81.80	82.27	82.11±1.57	1.51	NS	NS
Cold carcass weight, kg	70.03	69.41	68.76	67.11*	68.21*	68.70±2.99	1.78	NS	NS
Cold carcass yield, %	78.58	79.62	79.27	79.02	79.43	79.18±1.65	1.79	NS	NS
Loss yield, %	3.94	3.45	3.56	3.39	3.45	3.56±1.08	29.15	NS	NS
Loin depth, kg	32.23	30.51	31.36	31.43	31.29	31.36±1.74	5.40	NS	NS
Ham weight, kg	11.18	10.59	10.78	10.55	10.67	10.75±0.76	5.37	NS	NS
Backfat thickness, mm	11.80	13.10	11.60	12.80	12.40	12.34±3.20	22.14	NS	NS
Carcass length, cm	89.08	89.60	87.34	89.99	87.79	88.76±3.83	3.85	NS	NS
Longissimus area, cm <sup>2</sup>	38.75	39.83	37.95	37.75	38.70	38.60±4.28	11.13	NS	NS
Fat area, cm <sup>2</sup>	21.50	22.15	19.70	20.09	20.18	20.72±5.34	21.74	NS	NS
Lean meat in the carcass, kg	56.65	56.52	54.36	55.25	54.90	55.54±4.53	8.17	NS	NS
Lean meat in the carcass, %	77.76	78.60	76.51	79.52	77.71	78.02±6.23	7.98	NS	NS
Lean:fat ratio	0.57	0.56	0.54	0.43	0.54	0.53±0.18	33.32	NS	NS

<sup>†</sup> Standard Error; <sup>††</sup>CV = Coefficient of variation; <sup>§</sup> Linear Effect of Levels Canola Meal; <sup>¶</sup> Quadratic Effect of Levels Canola Meal; NS = non-significant; \* Significant ( $p \leq 0.05$ ) the Dunnett tests.

**Table 5.** Effect of diets with different levels of canola meal inclusion (CM) on qualitative traits of *Longissimus dorsi* muscle in pigs (60 to 90 kg).

Item	Inclusion levels of MC, %					Media±SE <sup>†</sup>	CV <sup>††</sup>	Lin <sup>§</sup>	Qua <sup>¶</sup>
	0	6	12	18	24				
pH, 45 min	6.24	6.18	6.31	6.50	6.44	6.33±0.35	4.28	0.04	NS
pH, 24 h	5.52	5.47	5.51	5.54	5.61	5.53±0.16	2.52	NS	NS
Drip loss, %	2.20	3.30	3.46	3.41	3.29	3.13±1.25	23.02	NS	NS
Marbling	1.90	1.70	2.30	1.80	2.00	1.94±0.71	34.92	NS	NS
Minolta a <sup>ac</sup>	7.82	7.27	6.53	6.85	6.07*	6.91±1.46	20.96	NS	NS
Minolta b <sup>ac</sup>	6.48	6.27	5.72	5.54	5.02*	5.80±1.16	19.09	0.02	NS
Minolta L <sup>ac</sup>	55.19	55.88	54.54	53.67*	51.80*	54.21±3.30	6.03	0.02	NS
Consistency	2.50	2.30	2.40	2.50	2.70	2.48±0.54	21.83	NS	NS
Color	2.00	1.80	1.80	2.10	2.10	1.96±0.49	22.26	NS	NS
FLT, % <sup>a</sup>	11.59	11.65	12.14	12.15	12.58	12.02±3.06	22.52	NS	NS
FLC, % <sup>b</sup>	27.69	29.70	30.08	27.30	30.09	28.97±4.68	13.78	NS	NS
Shear force, kgf/seg	2.71	2.75	2.80	3.08	2.77	2.82±0.64	20.11	NS	NS

<sup>†</sup> Standard error; <sup>††</sup>CV = Coeficiente de variación; <sup>§</sup> Linear Effect of Levels CM (Minolta b:  $Y=6.6145-0.0652583x$ ; Minolta L:  $Y=57.253-0.218756x$ ); <sup>¶</sup> Quadratic Effect of Levels CM; <sup>a</sup>FLT = Fluid loss from thawing; <sup>b</sup>FLC = Fluid loss from cooking; <sup>ac</sup> a\*: indicates the color of the flesh varies from red to green (green red high color indicates red, low color indicates green); b\*: indicates the flesh color varying from yellow to blue (b\* high indicates yellow color, b\* low indicates blue color); L\*: indicates the degree of brightness of the flesh (L\* = 0 dark meat, L = 100 white meat); NS = non-significant; \*Significant ( $p \leq 0.05$ ) the Dunnett tests.

There was a linear decrease ( $p \leq 0.05$ ) in lightness (L\*) and tonality (b\*) of the longissimus muscle as the CM level increased in the diet. Dunnett's test indicated differences ( $p \leq 0.05$ ) between the levels of CM inclusion and the control diet (0% CM) with respect to muscle colour.

With respect to lightness (L\*), the inclusion of 18 and 24% of CM exhibited difference ( $p \leq 0.05$ ) with respect to 0% CM. For the variables saturation

(a\*) and tonality (b\*), the inclusion of 24% CM showed differences ( $p \leq 0.05$ ) compared to 0%. Dunnett's test did not show difference ( $p \geq 0.05$ ) in the other qualitative muscle traits.

Rojo et al. (2001) reported that high canola levels darkens the meat. However, some authors (HERTZMANN et al., 1988; BELL et al., 1991; NRC, 1998) did not mention this risk when they used turnip paste with high levels of glucosinolates

and rich in erucic acid. As a result, the possibilities of generating dark meat with CM would be less.

## CONCLUSION

The addition of up to 24% CM did not interfere in the performance and quantitative and qualitative carcass traits in growing-finishing pigs. CM can be used as alternative protein source in the balancing of diets for pigs. However, it is important to conduct further studies to determine whether the cost-benefit ratio is adequate.

## REFERENCES

- ABIPECS - Associação brasileira da indústria produtora e exportadora de carne suína. **Estatística de produção**. Disponível em: <<http://www.abipecs.org.br/news/263/135/Produtores-de-suinos-apostam-no-mercado-interno-em-2011.html>>. Acesso em: 26 jul. 2011.
- BELL, J. M. Nutrients and toxicants in rapeseed meal: A review. **Journal of Animal Science**, Madison, v. 58, n. 4, p. 996-1010, 1984.
- BELL, J. M. Factors affecting the nutritional value of canola meal: a review. **Canadian Journal of Animal Science**, Saskatoon, v. 73, n. 4, p. 326-333, 1993.
- BELL, J. M.; KEITH, M. O.; HUTCHESON, D. S. Nutritional evaluation of very low glucosinolate canola meal. **Canadian Journal of Animal Science**, Saskatoon, v. 71, n. 2, p. 497-506, 1991.
- BRAND, T. S.; BRANDT, D. A.; CRUYWAGEN, C. W. Utilisation of growing-finishing pig diets containing high levels of solvent or expeller oil extracted canola meal. **New Zealand Journal of Agricultural Research**, Lahore, v. 44, n. 1, p. 31-35. 2001.
- BRIDI, A. M.; SILVA, C. A. **Avaliação da Carne Suína**, 2. ed. Londrina, PR: Midigraft, 2009. 120 p.
- Canola Council of Canada. **Canola Meal Feed Industry Guide**. 4. ed. Manitoba, Winnipeg: Canada Canola Council, 2009. 48 p.
- CAI, Y.; ZIMMERMAN, D. R.; EWAN, R. C. Diurnal variation in concentrations of plasma urea nitrogen and amino acids in pigs given free access to feed or fed twice daily. **Journal of Nutrition**, Ames, v. 124, n. 7, p. 1088-1093, 1994.
- COMA, J.; CARRION, D.; ZIMMERMAN, D. R. Use of plasma urea nitrogen as a rapid response criterion to determine the lysine requirement of pigs. **Journal of Animal Science**, Champaign, v. 73, n. 2, p. 472-481, 1995.
- CROMWELL, G. L. Feeding swine. In: KELLEMS, R. O.; CHURCH, D. C. **Livestock Feeds and Feeding**. Prentice-Hall, 6. ed. Englewood Cliffs, NJ: Prentice Hall, Inc. 2010. p. 244-284
- GOMES, P. C. et al. Uso do farelo de canola para suínos na fase de terminação. **Revista Brasileira de Zootecnia**, Viçosa, v. 27, n. 4, p. 749-753, 1998.
- HERTZMANN, C. L.; GORANSSON, R. H. Influence of fishmeal, rapeseed and rapeseed meal in the feed on the fatty acid composition and storage stability of porcine adipose tissue. **Meat Science**, Illinois, v. 23, n. 1, p. 53-64, 1988.
- KEITH, M. O.; BELL, J. M. Composition and digestibility of canola press cake as a feedstuff for use in swine diets. **Canadian Journal of Animal Science**, Saskatoon, v. 71, n. 3, p. 879-885, 1991.
- MOREIRA, I. et al. Utilização do farelo de canola na alimentação de suínos na fase total de crescimento e terminação (61 a 141 dias). **Revista Brasileira de Zootecnia**, Viçosa, v. 25, n. 4, p. 697-712, 1996.
- MOREIRA, I. et al. Determinação dos coeficientes de digestibilidade, valores energéticos e índices de controle de qualidade do milho e da soja integral processados pelo calor. **Revista Brasileira de Zootecnia**, Viçosa, v. 23, n. 4, p. 916-929, 1994.
- NARENDRAN, R. et al. Effect of different levels of tower rapeseed meal in corn – soybean meal based diets on growing - finishing pig performance. **Canadian Journal of Animal Science**, Saskatoon, v. 61, n. 1, p. 213-216, 1981.
- NATIONAL RESEARCH COUNCIL - NRC. **Nutrient Requirements of Swine**. 10 ed. Washington, DC. 1998.
- ROJO, G. A. et al. Canola meal as a protein supplement in diets for finishing pigs. **Revista Mexicana de Ciencias Pecuarias**, Tulancingo, v. 39, n. 2, p. 179-192, 2001.
- ROSTAGNO, H. S. et al. **Tabelas brasileiras para aves e suínos: composição de alimentos e exigências nutricionais**. 2. ed. Viçosa, MG: UFV, 2005. 186 p.
- ROSTAGNO, H. S. et al. **Tabelas brasileiras para aves e suínos: composição de alimentos e exigências nutricionais**. 3. ed. Viçosa, MG: UFV, 2011. 252 p.

ROTH-MAIER, D. A.; BÖHMER, B. M.; ROTH, F. X. Effects of feeding canola meal and sweet lupin (*L. luteus*, *L. angustifolius*) in amino acid balanced diets on growth performance and carcass characteristics of growing-finishing pigs. **Animal Research**, Munich, v. 54, n.1, p. 21-34, 2004.

SAKOMURA, N. K.; ROSTAGNO, H. S. **Métodos de pesquisa em nutrição de monogástricos**. Jaboticabal: Funep. 2007. 283 p.

SAMPAIO, I.B.M. **Estatística aplicada à experimentação animal**. Fundação de Ensino e Pesquisa em Medicina Veterinária e Zootecnia, Belo Horizonte, MG: UFMG, 1998. 221 p.

SANTOS, A. C.; BASSO, L. C. Composição química e qualidade nutricional do concentrado protéico de colza. **Arquivos de Biologia e Tecnologia**, Curitiba, v. 33, n. 4, p. 879-893, 1990.

SENEVIRATNE, R. W. et al. The nutritional value of expeller-pressed canola meal for grower-finisher pigs. **Journal Animal Science**, Champaign, v. 88, n. 6, p. 2073-2083, 2010.

SILVA, D. J.; QUEIROZ, A. C. **Análise de alimentos - métodos químicos e biológicos**. 3. ed. Viçosa, MG: UFV, 2002. 235 p.

SMITHARD, R. Full fat rapeseed for pig and poultry diets. **Feed Compounder**, Whitland, v., n. 1, p. 35-38, 1993.

SPRAGG, J. C.; MAILER, R. J. **Canola Meal Value Chain Quality Improvement**. Berwick, Victoria: AOF and Pork CRC 2007. 31 p.

TESKEREDZIC, Z. et al. Assessment of undephytinized and dephytinized rapeseed protein concentrate as sources of dietary protein for juvenile rainbow trout (*Oncorhynchus mykiss*). **Aquaculture**, Amsterdam, v. 132, n. 2, p. 261-277, 1995.

THACKER, P. A.; NEWKIRK, R. W. Performance of growing-finishing pigs fed barley-based diets containing toasted or non-toasted canola meal. **Canadian Journal of Animal Science**, Saskatoon, v. 85, n. 1, p. 53-59, 2005.

ZANOTTO, D. et al. Utilização do farelo de canola na alimentação de suínos em crescimento e terminação. **Archivos de Zootecnia**, Córdoba, v. 58, n. 224, p. 717-728, 2009.