

Dog parasite incidence and risk factors, from sampling after one-year interval, in Pinhais, Brazil

Incidência e fatores de risco de parasitas de cães, amostragem após um ano, Pinhais, Brasil

Camila Marinelli Martins¹; Cristiane da Conceição de Barros²; Daniele Bier¹; Ana Paula Marinho¹;
Jaqueline Maria Gonçalves Figueiredo¹; Juliano Leônidas Hoffmann³;
Marcelo Beltrão Molento¹; Alexander Welker Biondo^{1,4*}

¹Department of Veterinary Medicine, Federal University of Paraná – UFPR, Curitiba, PR, Brasil

²Department of Health Surveillance, Municipal Health Department, Pinhais, PR, Brasil

³Field Epidemiology Training Program – FETP/Brazil – EPISUS, Center for Strategic Information and Response in Health Surveillance, Department of Epidemiological Surveillance, Secretariat of Health Surveillance, Ministry of Health, Brasília, DF, Brasil

⁴Department of Veterinary Pathobiology, University of Illinois, Illinois, IL, USA

Received June 27, 2011

Accepted February 14, 2012

Abstract

Domestic animals in urban areas may serve as reservoirs for parasitic zoonoses. The aim of this study was to monitor the parasitic status of household dogs in an urban area of Pinhais, in the metropolitan region of Curitiba, Paraná State, Brazil, after a one-year period. In May 2009, fecal samples, skin scrapings and ticks were collected from 171 dogs. Questionnaires were applied to the owners (sex, age, environment and anthelmintic use). In May 2010, 26.3% (45/171) of the dogs were fecal samples reanalysed. From the fecal samples, 33.3% (57/171) in 2009 and 64.4% (29/45) in 2010 were positive. The parasite species most observed were, respectively in 2009 and 2010, *Ancylostoma* sp., 66.7 and 44.8%, and *Strongyloides stercoralis*, 26.3 and 3.4%. All the skin scrapings were negative, and no ticks or protozoa were found. There was no statistical association ($p > 0.05$) between positive fecal tests and age, sex or environment. In 2009 alone, dogs with a history of antiparasitic drug administration were 2.3 times more likely to be negative. A great number of replacement dogs was noticed one year later. Therefore, isolated antiparasitic treatment strategies may have no impact on parasite control, given the risk of introduction of new agents, thereby limiting the prevention strategies.

Keywords: Dog, parasitism, helminth, public health, zoonosis.

Resumo

Animais domésticos em áreas urbanas podem servir de reservatório para zoonoses parasitárias. O objetivo deste trabalho foi monitorar a situação parasitária de cães domiciliados, após um ano, em área urbana de Pinhais, região metropolitana de Curitiba, Estado do Paraná, Brasil. Em maio de 2009, foram coletadas amostras de fezes de 171 cães, realizados raspados cutâneos e pesquisa de carrapatos. Foi aplicado um questionário aos proprietários (sexo, idade, ambiente e uso de vermífugos). Em maio de 2010, 26,3% (45/171) dos cães tiveram as amostras de fezes analisadas novamente. Das amostras de fezes, 33,3% (57/171) em 2009 e 64,4% (29/45) em 2010, foram positivas. As espécies de parasitos mais frequentes em 2009 e 2010 foram, respectivamente, *Ancylostoma* sp. 66,7 e 44,8% e *Strongyloides stercoralis*, 26,3 e 3,4%. Todos os raspados cutâneos foram negativos e nenhum carrapato ou protozoário foi encontrado. Não houve associação estatística ($p > 0.05$) entre exame positivo e idade, sexo ou ambiente. Somente em 2009, cães com histórico de antiparasitários tiveram 2,3 vezes mais chance de serem negativos. Foi observada grande substituição dos cães após um ano. Dessa forma, estratégias isoladas de tratamento podem obter efeito nulo quanto ao controle de parasitas, haja vista o risco de introdução de novos agentes, limitando estratégias de prevenção dos mesmos.

Palavras-chave: Cão, parasitismo, helminto, saúde pública, zoonoses.

*Corresponding author: Alexander Welker Biondo

Departamento de Medicina Veterinária, Universidade Federal do Paraná – UFPR,
Rua dos Funcionários, 1540, Juvevê, CEP 80035-050, Curitiba, PR, Brasil

e-mail: abiondo@ufpr.br

Introduction

The excessive number of owned dogs hosting intestinal parasites that have access to the streets in Brazilian urban areas may expose local communities to contaminated feces in public areas (Robertson et al., 2000). This has been reported as a risk factor for zoonotic infections (CAPUANO; ROCHA, 2005). Dog endoparasites with occasional human infection include *Toxocara canis* and *Ancylostoma* sp., which are the causative agents of visceral and cutaneous larva migrans, respectively (BÄCHLI et al., 2004), and *Cryptosporidium parvum*, *Giardia lamblia* and *Dipylidium caninum*, which may cause gastrointestinal disorders (OLIVEIRA-SEQUEIRA et al., 2002).

Several studies conducted worldwide on dog endoparasites have shown varying results. The rate of infection ranged from 2.4 to 33.2% ($9.8 \pm 13.1\%$) to *Ancylostoma* sp. and 0.7 to 33.6% ($11.5 \pm 11.4\%$) to *Toxocara* sp. In surveys conducted in Australia, Switzerland, Italy, Spain and Greece (NABAIS, 2008). *Trichuris vulpis* infection was found to range from 15 to 20% in the United States (BLAGBURN et al., 1996). In South America, a few studies in Colombia, Venezuela, Argentina and Chile have shown similar widely ranging results (NABAIS, 2008; FONTANARROSA et al., 2006; LÓPEZ et al., 2006), with *Ancylostoma* spp. infection ranging from 1.8 to 61.1% ($22.4 \pm 26.3\%$) and *Toxocara* spp. from 2.5 to 55% ($19.9 \pm 23.7\%$).

A review performed by a study in Brazil verified the status of fecal parasitism in dogs in several cities in the southern region (Porto Alegre and Londrina) and the southeastern region (Viçosa, Uberlândia, São Paulo, Araçatuba, Espírito Santo do Pinhal, Botucatu and Monte Negro) (LABRUNA et al., 2006). The highest frequency observed in this study was for *Ancylostoma* spp. infection, ranging from 13.5 to 79.1%, followed by *Toxocara* spp., from 5.5 to 39.0%, *Trichuris vulpis*, from 0.0 to 70.0%, and *Dipylidium caninum*, from 0.0 to 8.3%. Another, other studies were conducted in the central-western region (Goiânia), southern region (Itapema, Porto Alegre, Curitiba and Rio Grande) and southeastern region (Praia Grande and Lavras), showing *Ancylostoma* spp. infection ranging from 9.2 to 93.7%, followed by *Trichuris vulpis*, from 3.25 to 32.5%, *Toxocara* spp., from 1.2 to 23%, and *Dipylidium caninum*, from 0.26 to 1.9% (SCAINI et al., 2003; LEITE et al., 2004; ALVES et al., 2005; BLAZIUS et al., 2005; CASTRO et al., 2005; GUIMARÃES et al., 2005; LORENZINI et al., 2007).

All these previous studies were conducted on single-sample prevalence of endoparasites in different dog populations. However, no survey has previously been conducted on parasitism within the same dog population over time. Accordingly, the aim of the present study was to monitor the parasitic status of dogs after a one-year interval in an urban area of Pinhais, in the metropolitan region of Curitiba, southern Brazil.

Material and Methods

The survey for the present study was conducted over a two-day period in May 2009 and again in May 2010, in the Bonilauri neighborhood of the Alto Tarumã district, city of Pinhais. The city is located at latitude $25^{\circ} 26' 41''$ and longitude $49^{\circ} 11' 33''$,

and is in the western part of the metropolitan region of Curitiba (Figure 1). Pinhais has 112,852 inhabitants (IBGE, 2010), and its human development index (HDI) is 0.815. This study received technical and official support from the Zoonosis Control Center (ZCC) of Pinhais and the fieldwork was done by 60 volunteers and ZCC personnel. Since the study was approved by the city's Health Department as part of its yearly activities, dog owners were officially informed house-to-house by health agents and were encouraged to participate in the survey.

The Bonilauri neighborhood is a homogeneous settlement of resource-poor people distributed in 380 households and was chosen as the study area. The area has an estimated population of 826 people and 297 dogs, with a dog-to-human ratio of 1:2.78 (PAMPUCH, 2008, personal communication). The number of sampled animals with 5% confidence level ($p < 0.05$) was 168 dogs (FLEISS, 1981). The houses were divided according to the number of blocks/streets and number of houses per square meter, to ensure homogeneous spatial distribution.

Permission forms were handed out to dog owners, who signed them before animal sampling was done. A questionnaire was used to obtain household and animal information (sex, age, street access, characteristics of the dog's living place and whether the dogs had been given antiparasitic drugs at least once). The information was stratified according to sex, age (up to year of age or over one year), street access (yes or no), quality of the dog's living place (excellent, good, regular or poor) and antiparasitic treatment (yes or no). All available animals belonging to the household underwent fecal sampling, skin scraping and tick collection. The sampling procedures were performed similarly in 2010, when the same animals were sought, based on the original registration forms.



Figure 1. Map of South America and the location of the cities of Pinhais and Curitiba, state of Paraná, Brazil.

Fecal samples were collected directly from the dogs' rectal ampulla and around 1 g of feces was placed into a commercial transportation tube containing 9% formalin (Paratest®, Diagnostek, São Paulo, Brazil). Parasite eggs were observed under an optical microscope and were identified in accordance with standard techniques (MONTEIRO, 2011). All the animals were examined for the presence of ticks and skin lesions and skin scrapings were placed into a glass tube for further analysis.

1. Statistical analysis

The results were analysed with Epi-info (Epi-Info version 3.5.2, Centers for Diseases Control and Prevention, Atlanta, USA). The odds ratio was calculated for each parameter, and descriptive and

chi-square analyses were performed at a confidence level of 5% ($p < 0.05$). Maps were created using the ArcView software.

Results

Out of the total number of fecal samples obtained in 2009, 57/171 (33.3%) was positive for at least one parasite. Among the parasites, 66.7% were identified as *Ancylostoma* spp., 26.3% as *Strongyloides stercoralis*, 14.1% as *Trichuris vulpis* and 10.5% as *Toxocara canis*. When coinfection was evaluated, 86% was found to be infected with only one, 8.8% with two and 5.3% with three helminth species simultaneously (Table 1). In 2010, 64.4% were positive for at least one parasite, and 58.6% of the samples were identified as positive for *Toxocara canis*, 44.8% for *Ancylostoma* spp.,

Table 1. Results stratified according to the dogs' sex, age, street access, living place and antiparasitic treatment, including positive and negative results from 2009 and 2010, in the Bonilauri area of Pinhais, Brazil.

	2009			2010		
	Positive n (%)	Negative n (%)	Total	Positive n (%)	Negative n (%)	Total
Males	36 (63.8)	56 (49.6)	92	14 (48.3)	7 (43.7)	21
Females	21 (36.8)	57 (50.4)	78	15 (51.7)	9 (56.2)	24
Not informed*			1			0
Total	57 (100)	113 (100)	171	29 (100)	16 (100)	45
OR/p		0.57/	0.09		0.83/	0.77
Age						
Up to one year	19 (35.2)	23 (23.2)	42	5 (19.2)	1 (6.2)	6
Over one year	35 (65.8)	76 (76.7)	111	21 (80.1)	15 (93.7)	36
Not informed*			19			3
Total	54 (100.0)	99 (100.0)	171	26 (100.0)	16 (100.0)	45
OR / p		0.56/	0.11		0.28/	0.24
Street access						
Yes	24 (42.1)	41 (36.3)	45	10 (38.5)	7 (43.7)	17
No	33 (57.9)	72 (63.7)	105	16 (61.5)	9 (56.2)	25
Not informed*			21			3
Total	57 (100.0)	113 (100.0)	171	26 (100.0)	16 (100.0)	45
OR / p		0.78/	0.46		1.24/	0.73
Living place						
Excellent and good	22 (40.7)	42 (38.2)	64	10 (35.7)	7 (46.6)	17
Regular and poor	32 (59.2)	68 (61.8)	100	18 (64.3)	8 (53.3)	26
Not informed*			7			2
Total	54 (100.0)	110 (100.0)	171	28 (100.0)	15 (100.0)	45
OR / p		0.89/	0.75		1.57/	0.48
Antiparasitic drugs						
Treated	24 (43.6)	70 (64.2)	94	19 (42.2)	11 (24.4)	30
Not treated	31 (56.4)	39 (35.8)	70	7 (15.5)	5 (11.1)	12
Not informed*			7			3
Total	55 (100.0)	109 (100.0)	171	26 (100.0)	16 (100.0)	45
OR / p		2.31/	0.01		0.81/	0.76
Comparative between years						
		1		2.66/		0.24

n = number of samples; *data not informed were not taken into consideration; OR = odds ratio; p = statistical significance at 5% level.

10.3% for *Toxascaris* spp., 6.9% for *Trichuris vulpis* and 6.9% for *Dipylidium caninum*. A total of 21/29 animals (72.4%) were infected with only one parasite, 24.2% with two and 3.4% with three species simultaneously.

No protozoan parasites were found in the samples, at either sampling time. Skin scrapings were obtained from 30 dogs in 2009 and 7 dogs in 2010, and all the results were negative. No ticks were found at either sampling time.

The information regarding gender, age, outdoor access and living conditions were not statically significantly different between the positive and negative animals. Males and females were equally likely to be negative in 2009, and although there was a tendency for males to be more infected ($p = 0.09$), no significant difference was found. Similarly, no gender differences were observed in 2010 ($p = 0.77$). In addition, no differences between endoparasites were found for any age ($p = 0.11$ and 0.24), for outdoor access ($p = 0.46$ and 0.73) or for living place ($p = 0.75$ and 0.48), in 2009 and 2010 respectively. Dogs that were infected in 2009 had a similar tendency to be infected in 2010 ($p = 0.24$), but dogs with no history of anthelmintic treatment in 2009 were statistically different ($p = 0.01$) and 2.3 times more likely to be infected than were treated animals (Table 1).

Out of the sampled households, 34.6% in 2009 and 67.6% in 2010 presented at least one dog with endoparasites. The positive and negative dogs per household in the 2009 and 2010 samplings are presented in Figure 2. A total of 171 dogs from 150 households were successfully sampled in 2009, but only 45 (26.3%) of the same dogs, from 37 households, were found in 2010 based on the previous registrations. The main reason for not finding the 126 animals that were missing in 2010 were as follows: dogs whose owners were not present at the house (30.2%); inadequate or

insufficient sample obtained (21.4%); dog deaths (17.5%); dogs with no signed permission (15.1%); owners who had moved out of the neighborhood (9.5%), lost dogs (8.7%), aggressive dogs (8.7%); and donated dogs (4.8%).

Discussion and Conclusion

The high rate of infected dogs at the two sampling times is within the range found in previous studies in different regions of Brazil (LEITE et al., 2004; ALVES et al., 2005; BLAZIUS et al., 2005; LORENZINI et al., 2007). *Ancylostoma* spp. was also one of the most prevalent parasite species in previous studies (LORENZINI et al., 2007; LEITE et al., 2004; CASTRO et al., 2005; GUIMARÃES et al., 2005; BLAZIUS et al., 2005). The second most prevalent parasite found in 2009 was *Strongyloides stercoralis*. This result is different from most of the results found in the literature in Brazil (LEITE et al., 2004; ALVES et al., 2005; BLAZIUS et al., 2005; LORENZINI et al., 2007), but the same as in a study in the city of Rio Grande (SCAINI et al., 2003). *Trichuris vulpis* eggs were found in both 2009 and 2010, and similar results were observed in São Paulo (OLIVEIRA-SEQUEIRA et al., 2002), Santa Catarina (BLAZIUS et al., 2005) and Curitiba (LEITE et al., 2004). *Toxocara canis* was the main helminth found in 2010, corresponding to 58.6% of the number of eggs found. However, other studies in Brazil have indicated widely ranging lower prevalence for this parasite, from 1 to 11% (OLIVEIRA-SEQUEIRA et al., 2002; SCAINI et al., 2003; LEITE et al., 2004; ALVES et al., 2005; BLAZIUS et al., 2005; CASTRO et al., 2005; LORENZINI et al., 2007). Only a single study has demonstrated similar prevalence, of 70% (GUIMARÃES et al., 2005).

No protozoan parasites were found at either sampling time, thus differing from other studies in Brazil that have detected protozoa. *Giardia* sp., *Cystoisospora* sp., *Sarcocystis* sp. and *Cryptosporidium parvum* have been the protozoa most frequently found (OLIVEIRA-SEQUEIRA et al., 2002; LEITE et al., 2004; ALVES et al., 2005; BLAZIUS et al., 2005; LORENZINI et al., 2007). Our data may reflect the low number of these parasites in this area and the environmental effect of the fall season. No animals presented diarrhea.

In urban areas, the most frequent tick species is *Rhipicephalus sanguineus* (RIBEIRO et al., 1997). The absence of adult ticks on the animals of the present study may have been due to the seasonality of this species, since the study was conducted in late fall, a period when *R. sanguineus* could be in the larval or nymph stages (SOUZA, 2008), living in small cracks in concrete or wood surfaces (DIPEOLU, 1982). Although fleas and lice are important regarding transmission of zoonotic diseases, only a small number of fleas (*Ctenocephalides* sp.) and no lice were found in the dogs of the present study.

Host characteristics relating to parasite burden, including gender, age, street access, living place and antiparasitic treatment, are still a matter of major controversy. Some studies, similarly to the present one, have not found any statistical significance or tendencies, either for males or for females (ALVES et al., 2005; LABRUNA et al., 2006). However, males had higher parasite burden in another study (FONTANARROSA et al., 2006). Most



Figure 2. Description of the Bonilauri area and the locations at which dogs were sampled for parasitological tests in 2009 and 2010. Filled squares correspond to positive animals in 2009 and empty squares correspond to negative animals in 2009. Filled circles correspond to positive animals in 2010 and empty circles correspond to negative animals in 2010.

other studies have shown that dogs younger than one year of age are more likely to be parasitized (OLIVEIRA-SEQUEIRA et al., 2002; ALVES et al., 2005; FONTANARROSA et al., 2006; LORENZINI et al., 2007), although a single previous study showed that there was no statistical difference regarding age, similarly to the present study (LABRUNA et al., 2006). We consider that this factor may be dependent on the intensity of parasite infection and the local conditions for the animals.

To the present authors' knowledge, only a single study has compared street access as a risk factor for parasite burden (LABRUNA et al., 2006). That study presents results similar to those of the present study, thus showing that both the indoor and the outdoor environments may have been contaminated with viable parasite eggs in these urban areas (CHAVES et al., 2006; PRADO et al., 2001).

A history of antiparasitic drug administration may be a significant single protection factor, as previously published (XAVIER, 2006), thus indicating that periodic use of anthelmintics may cause a reduction in helminth frequency in dogs (JORDAN et al., 1993; BUGG, 1999). To the present authors' knowledge, no study has previously been conducted on the same animals over a period of time. However, the same dogs in this area were found only 26.3% in 2010, while the other 73.7% were not found or had died or moved. Although this situation was accounted for through using a questionnaire at the beginning of the survey, the high turnover rate among the dogs in this area was surprising. The high death rate (17.5%) may reflect the short life expectancy of dogs in resource-poor areas, and we consider that any intervention will only be successful when the strategies focus on educating owners to take responsible care of their animals.

In conclusion, we have demonstrated that programs that monitor animal health status are effective, and that even dogs that did not have street access can become reinfected with parasite species that are important from a public health point of view. Annual application of antiparasitic drugs may not be sufficiently effective for parasite eradication, with regard to both the parasitic and the free-living stages, but it may significantly lengthen the time taken for reinfection to become established, thus leading to a safer environment. In addition, the high turnover among the dogs in this area may reflect an attitude towards human and animal health that is of great concern, since it may cause an influx of new pathogens that can harm local communities.

References

- Alves OF, Gomes AG, Silva AC. Ocorrência de enteroparasitos em cães do município de Goiânia, Goiás: Comparação de técnicas de diagnóstico. *Cienc Anim Bras* 2005; 6(2): 127-133.
- Bächli H, Minet JC, Gratzl O. Cerebral toxocaríasis: a possible cause of epileptic seizure in children. *Child Nerv Syst* 2004; 20(7): 468-472. PMID:15138789.
- Blagburn BL, Lindsay DS, Vaughan JL, Rippey NS, Wright JC, Lynn RC, et al. Prevalence of canine parasites based on fecal flotation. *Comp Contin Educ Vet Pract* 1996; 18: 483-509.
- Blazius RD, Emerick S, Prophiro JS, Romão PRT, Silva OS. Ocorrência de protozoários e helmintos em amostras de fezes de cães errantes da Cidade de Itapema, Santa Catarina. *Rev Soc Bras Med Trop* 2005; 38(1): 73-74. PMID:15717103. <http://dx.doi.org/10.1590/S0037-86822005000100018>
- Bugg RJ, Robertson ID, Elliot AD, Thompson RCA. Gastrointestinal parasites of urban dogs in Perth, Western Australia. *Vet J* 1999; 157(3): 295-301. PMID:10328840. <http://dx.doi.org/10.1053/tvjl.1998.0327>
- Capuano DM, Rocha GM. Environmental contamination by *Toxocara* sp. Eggs in Ribeirão Preto, São Paulo state, Brazil. *Rev Inst Med Trop São Paulo* 2005; 47(4): 223-226. <http://dx.doi.org/10.1590/S0036-46652005000400009>
- Castro JM, Santos SV, Monteiro NA. Contaminação de canteiros da orla marítima do Município de Praia Grande, São Paulo, por ovos de *Ancylostoma* e *Toxocara* em fezes de cães. *Rev Soc Bras Med Trop* 2005; 38(2): 199-201. PMID:15821802. <http://dx.doi.org/10.1590/S0037-86822005000200017>
- Chaves EMS, Vazquez L, Lopes K, Flores J, Oliveira L, Rizzi L, et al. Levantamento de Protozoonoses e Verminoses nas sete creches municipais de Uruguaiana, Rio Grande do Sul - Brasil. *RBAC* 2006; 38(1): 39-41.
- Dipeolu OO, Akinboade AO, Ogunji FO. Observations on the epidemiology of house infesting *Rhipicephalus sanguineus* in a household in Lagos, Nigeria. *Bull Anim Health Prod Afr* 1982; 30(1): 29-30. PMID:7186809.
- Fleiss JL. *Statistical Methods for rates and proportions*. Wiley-Interscience; 1981. 414 p.
- Fontanarrosa MF, Vezzani D, Basabe J, Eiras DF. An epidemiological study of gastrointestinal parasites of dogs from Southern Greater Buenos Aires (Argentina): age, gender, breed, mixed infections, and seasonal and spatial patterns. *Vet Parasitol* 2006; 136(3-4): 283-95. PMID:16364551. <http://dx.doi.org/10.1016/j.vetpar.2005.11.012>
- Guimarães AM, Alves EG, De Rezende GF, Rodrigues MC. Ovos de *Toxocara* sp. e larvas de *Ancylostoma* sp. em praça pública de Lavras, MG. *Rev Saúde Pública* 2005; 39(2): 293-295. PMID:15895151.
- Instituto Brasileiro de Geografia e Estatística – IBGE. *Censo 2010 Paraná* [online]. 2010 [cited 2011 Feb. 04]. Available from: http://www.ibge.com.br/home/estatistica/populacao/censo2010/tabelas_pdf/total_populacao_parana.pdf.
- Jordan HE, Mullins ST, Stebbins ME. Endoparasitism in dogs: 21,583 cases (1981-1990). *J Am Vet Med Assoc* 1993; 203(4): 547-549. PMID:8407513.
- Labruna MB, Pena HFJ, Souza SLP, Pinter A, Silva JCR, Ragozo AMA, et al. Prevalência de endoparasitas em cães da área urbana do município de Monte Negro, Rondônia. *Arq Inst Biol* 2006; 73(2): 183-193.
- Leite LC, Marinoni LP, Círio SM, Diniz JMF, Silva MAN, Luz E, et al. Endoparasitas em cães (*Canis familiaris*) na cidade de Curitiba, Paraná, Brasil. *Arch Vet Sci* 2004; 9(2): 95-99.
- López JD, Abarca KV, Paredes PM, Inzunza ET. Intestinal parasites in dogs and cats with gastrointestinal symptoms in Santiago, Chile. *Rev Méd Chil* 2006; 134(2): 193-200. PMID:16554927.
- Lorenzini G, Tasca T, Carli, GA. Prevalence of intestinal parasites in dogs and cats under veterinary care in Porto Alegre, Rio Grande do Sul, Brazil. *Braz J Vet Res Anim Sci* 2007; 44(2): 137-145.
- Monteiro SG. *Parasitologia na Medicina Veterinária*. Editora Roca; 2011.

- Nabais PMMD. *Controlo de helmintoses gastrointestinais em cães* [Dissertação]. Lisboa: Faculdade de Medicina Veterinária de Lisboa; 2008.
- Oliveira-Sequeira TCG, Amarante AF, Ferrari TB, Nunes LC. Prevalence of intestinal parasites in dogs from São Paulo State, Brazil. *Vet Parasitol* 2002; 103(1-2): 19-27. [http://dx.doi.org/10.1016/S0304-4017\(01\)00575-1](http://dx.doi.org/10.1016/S0304-4017(01)00575-1)
- Prado MS, Barreto ML, Strina A, Faria JA, Nobre AA, Jesus SR. Prevalence and intensity of infection by intestinal parasites in school-aged children in the City of Salvador (Bahia, Brazil). *Rev Soc Bras Med Trop* 2001; 34(1): 99-101. PMID:11340505.
- Ribeiro VLS, Weber MA, Fetzter LO, Vargas CRB. Espécies e prevalência das infestações por carrapatos em cães de rua da cidade de Porto Alegre, RS, Brasil. *Cienc Rural* 1997; 27(2): 285-289. <http://dx.doi.org/10.1590/S0103-84781997000200019>
- Robertson ID, Irwin PJ, Lymbery AJ, Thompson RCA. The role of companion animals in the emergence of parasitic zoonoses. *Int J Parasitol* 2000; 30: 1369-1377.
- Scaini CJ, Toledo RN, Lovatel R, Dionello MA, Gatti FA, Susin L, et al. Contaminação ambiental por ovos e larvas de helmintos em fezes de cães na área central do Balneário Cassino, Rio Grande do Sul. *Rev Soc Bras Med Trop* 2003; 36(5): 617-619. PMID:14576878.
- Souza BMPS. *Epidemiologia da Ehrlichia sp. e avaliação do seu potencial zoonótico no município de Salvador, Bahia* [Dissertação]. Salvador: Escola de Medicina Veterinária, Universidade Federal da Bahia; 2008.
- Xavier GA. *Prevalência de endoparasitos em cães de companhia em Pelotas-RS e risco zoonótico* [Monografia]. Pelotas: Instituto de Biologia da Universidade Federal de Pelotas; 2006.