

Caryospora peneireiroi n. sp. (Apicomplexa: Eimeriidae) in the common kestrel, *Falco tinnunculus* (Falconiformes: Falconidae), in mainland Portugal

Caryospora peneireiroi n. sp. (Apicomplexa: Eimeriidae) no peneireiro, *Falco tinnunculus* (Falconiformes: Falconidae), em Portugal Continental

Sergian Vianna Cardozo^{1,2}; Bruno Pereira Berto^{3*}; Inês Caetano⁴; Viviane Camara Maniero¹; Isabel Pereira da Fonseca⁵; Carlos Wilson Gomes Lopes⁶

¹ Programa de Pós-graduação em Biomedicina Translacional, Universidade do Grande Rio – UNIGRANRIO, Duque de Caxias, RJ, Brasil

² Programa de Pós-graduação em Ciência, Tecnologia e Inovação em Agropecuária, Universidade Federal Rural do Rio de Janeiro – UFRRJ, Seropédica, RJ, Brasil

³ Departamento de Biologia Animal, Instituto de Ciências Biológicas e da Saúde, Universidade Federal Rural do Rio de Janeiro – UFRRJ, Seropédica, RJ, Brasil

⁴ Centro de Recuperação de Animais Silvestres de Lisboa, Parque Florestal de Monsanto, Lisboa, Portugal

⁵ Centro Interdisciplinar de Investigação em Sanidade Animal, Faculdade de Medicina Veterinária, Universidade de Lisboa, Lisboa, Portugal

⁶ Departamento de Parasitologia Animal, Instituto de Veterinária, Universidade Federal Rural do Rio de Janeiro – UFRRJ, Seropédica, RJ, Brasil

Received March 7, 2016

Accepted April 7, 2016

Abstract

The common kestrel *Falco tinnunculus* Linnaeus, 1758, is a widespread raptor, native in Europe, Asia and Africa, and vagrant in the Americas. In the current work, 27 fecal samples were collected from common kestrels kept in the Lisbon Center for Wild Animal Recovery, located at Monsanto Forest Park, Lisbon, Portugal. Five (19%) of them were found to be passing an undescribed species of *Caryospora* in their feces. The oocysts of *Caryospora peneireiroi* n. sp. were ellipsoidal with a bilayered wall and measured $47.1 \times 37.6 \mu\text{m}$ with a shape index of 1.25. No micropyle, oocyst residuum or polar granule was present. The sporocysts were subspherical, measuring $25.1 \times 24.3 \mu\text{m}$. Stieda, sub-Stieda and para-Stieda bodies were absent. The sporocyst residuum was composed of many homogenous globules scattered throughout the periphery of the sporocyst. This is the fourth caryosporan species reported from *F. tinnunculus*.

Keywords: Coccidia, oocysts, morphology, taxonomy, conservation, raptors.

Resumo

O peneireiro *Falco tinnunculus* Linnaeus, 1758, é uma ave de rapina com vasta distribuição geográfica, nativa da Europa, Ásia e África, e errante nas Américas. No presente trabalho, 27 amostras de fezes foram coletadas de peneireiros mantidos no Centro de Recuperação de Animais Silvestres de Lisboa, localizado no Parque Florestal de Monsanto, Lisboa, Portugal. Cinco (19%) deles eliminaram uma espécie não descrita de *Caryospora* em suas fezes. Os oocistos de *Caryospora peneireiroi* n. sp. foram elipsoides, com parede de dupla camada, medindo $47,1 \times 37,6 \mu\text{m}$, e índice morfométrico de 1,25. A micrópila, resíduo do oocisto e grânulo polar não foram evidenciados. Os esporocistos foram subesféricos, medindo $25,1 \times 24,3 \mu\text{m}$. Corpos de Stieda, substieda e parastieda não se manifestaram. Resíduo do esporocisto foi composto de muitos glóbulos homogêneos espalhados por toda a periferia do esporocisto. Essa é a quarta espécie *Caryospora* descrita de *F. tinnunculus*.

Palavras-chave: Coccidia, oocistos, morfologia, taxonomia, conservação, aves de rapina.

*Corresponding author: Bruno Pereira Berto. Departamento de Biologia Animal, Instituto de Ciências Biológicas e da Saúde, Universidade Federal Rural do Rio de Janeiro – UFRRJ, Rodovia BR-465 Km 7, CEP 23897-000, Seropédica, RJ, Brasil. e-mail: bertobp@ufrj.br

Introduction

The common kestrel *Falco tinnunculus* Linnaeus, 1758, is a raptor observed in several environments, easily recognizable through its ability to hover while searching for its prey. In Portugal, this species was one of the most abundant in the 1940s; however, it suffered a sharp decline over the subsequent decades. Nowadays, although the population trend appears to be still decreasing, it is recognized that the population size is still extremely large (PALMA, 1985; Birdlife International, 2015).

Genus *Caryospora* Leger, 1904, is occasionally reported in raptors and reptiles, which act as its definitive hosts, and in rodents, which act as its intermediate hosts (UPTON et al., 1990; BERTO et al., 2014). *Caryospora* is the third largest genus in the family Eimeriidae. *Caryospora* spp. have now been reported worldwide, especially in raptors kept in captivity (UPTON et al., 1992; PAPAZAHARIADOU et al., 2001; MCALLISTER et al., 2013a, b). However, little is known about the distribution and importance of species of the genus *Caryospora* in free-living birds (YANG et al., 2014). The current work describes a new species of *Caryospora* from the common kestrels *F. tinnunculus* that were being kept for rehabilitation and reintroduction into the wild at the Lisbon Center for Wild Animal Recovery (Centro de Recuperação de Animais Silvestres de Lisboa, LxCRAS), Monsanto Forest Park, Lisbon, Portugal.

Materials and Methods

Twenty-seven fecal samples were collected from common kestrels that were being kept in individual cages on the premises of the Lisbon Center for Wild Animal Recovery (LxCRAS), located in Monsanto Forest Park, Lisbon, Portugal. The samples were collected immediately after defecation and were placed in plastic

vials containing 2.5% potassium dichromate ($K_2Cr_2O_7$) solution at 1:6 (v/v). In the laboratory, the samples were incubated at room temperature for 10 days or until day three when around 70% of the oocysts had sporulated. The oocysts were recovered by means of flotation in Sheather's sugar solution (specific gravity: 1.20). Morphological observations, line drawings, photomicrographs and measurements were made using an Olympus BX40 microscope equipped with a digital camera (Olympus DP10). The line drawings were edited using two software applications from CorelDRAW® (Corel Draw Graphics Suite, Version 11.0, Corel Corporation, Canada), specifically Corel DRAW and Corel PHOTO-PAINT. All measurements were made in micrometers and are given as the range followed by the mean in parentheses. The descriptions of oocysts and sporocysts followed the guidelines of Duszynski & Wilber (1997) and Berto et al. (2014), as follows: oocyst (O) length (L) and width (W) and their ranges and ratios (L/W); micropyle (M), oocyst residuum (OR), polar granule (PG) and sporocyst (SP) length (L) and width (W) and their ranges and ratios (L/W); and Stieda body (SB), sub-Stieda body (SSB), para-Stieda body (PSB), sporocyst residuum (SR), sporozoite (SZ), refractile body (RB) and nucleus (N).

Results

Caryospora peneireiroi n. sp.

(Figures 1a, b)

Description of sporulated oocyst: Oocyst shape: ellipsoidal; oocyst wall: bilayered, ~2.0 thick; outer layer smooth; L × W (n = 20) 47.1 × 37.6 (42-49 × 34-40); L/W 1.25 (1.2-1.4). M, OR, PG: all absent. Distinctive features of oocysts: large oocyst ellipsoidal lacking M, OR, and PG.




Figure 1. Oocysts of *Caryospora peneireiroi*, a new coccidium species recovered from the common kestrel *Falco tinnunculus*. (a) Composite line drawing; (b-e) Photomicrographs. Bars=10 µm.

Description of sporocyst and sporozoites: Sporocyst shape: subspherical; sporocyst wall: single-layered, ~0.8 thick, slightly rough to wrinkled; L × W (n = 20) 25.1 × 24.3 (24–27 × 24–25); L/W 1.03 (1.0–1.1); SB, SSB, PSB: all absent; SR: present; SR characteristics: composed of many homogenous globules scattered throughout the periphery of the SP. SZ: stout, L × W (n = 5) 17.0 × 4.2 (15–18 × 3–5) in situ, arranged parallel to one another in SP. Each SZ without striations discernible, but with one medium-sized spherical PRB, 3.0 (2–4) wide; single N in midpoint of body. Distinctive features of sporocyst: large size, presence of globules scattered throughout the periphery of the SP, and large SZ.

Taxonomic summary

Type host: common kestrel *Falco tinnunculus* Linnaeus, 1758 (Aves: Falconiformes: Falconidae).

Other hosts: None.

Type specimens: Phototypes and line drawings are deposited and available (<http://r1.ufrrj.br/lcc>) in the Parasitology Collection of the Coccidia and Coccidiosis Laboratory, at the Federal Rural University of Rio de Janeiro (UFRRJ), located in Seropédica, State of Rio de Janeiro, Brazil. Photographs of the type-host specimens (symbiotypes) are deposited in the same collection. The repository number is P-63/2015.

Type locality: Lisbon Center for Wild Animal Recovery (LxCRAS), Lisbon, Portugal (38°44'22.9" N, 9°11'02.3" W).

Prevalence: Five out of 27 (19%).

Sporulation: Exogenous. All oocysts were passed in the feces unsporulated and were fully sporulated by day 7 in $K_2Cr_2O_7$ solution at room temperature.

Prepatent and patent periods: Unknown.

Site of infection, definitive host: Unknown. Oocysts recovered from feces.

Site of infection, secondary host: Unknown.

Endogenous stages, definitive host: Unknown.

Endogenous stages, secondary host: Unknown.

Cross-transmission: None to date.

Pathology, definitive host: Unknown.

Pathology, secondary host: Unknown.

Etymology: The specific epitaph is derived from the common local name for the host, which is 'peneireiro'.

Remarks

Table 1 and Table 2 show some characteristic features of *Caryospora* spp. described from Accipitriformes and Falconiformes respectively. Of *Caryospora* species listed in Table 1, only *Caryospora aquilae* Volf, Koudela and Modrý, 2000, *Caryospora petersoni* McAllister, Duszynski and McKown, 2013, and *Caryospora hanebrinki* McAllister, Duszynski and McKown, 2013, have oocyst measurements similar to those of *C. peneireiroi*. However, *C. aquilae* can be distinguished through the smooth surface of the SP and smaller SZ, which has N not discernible. *Caryospora petersoni* and *C. hanebrinki*, in turn, can easily be distinguished through the smooth surface of the SP, SR forming a compact mass and SZ with striations. Of the species of *Caryospora* described from Falconiformes (Table 2), only *Caryospora megafalconis* Klüh, 1994, and *Caryospora biarmicus* Alyousif, Alfaleh and Al-Shawa, 2011, have oocyst measurements similar to those of *C. peneireiroi*. However, *C. megafalconis* can be distinguished through the smooth surface of the SP and SR forming a compact mass. *Caryospora biarmicus* can be distinguished through the presence of OR and smaller SP with smooth surface.

Discussion

The family Falconidae comprises 66 species of raptors distributed in 11 genera. The genus *Falco* comprises 39 species worldwide, while other genera are exclusively Neotropical (Birdlife International, 2015). The common kestrel *F. tinnunculus* Linnaeus, 1758, is a widespread kestrel species, found throughout Europe, Asia and Africa. In Europe, *F. tinnunculus* is migratory in winter, heading for southern Europe and sub-Saharan Africa (VILLAGE, 1990; Birdlife International, 2015).

Table 1. Comparative data of *Caryospora* spp. described from Accipitridae (Accipitriformes).

Species	Host	Locality	Reference	Oocyst		Sporocyst	
				Shape	Size (μm)	Shape	Size (μm)
<i>C. arcayae</i>	<i>Buteo magnirostris</i> ; <i>Buteo platypterus</i>	Venezuela; USA	Upton et al. (1990)	subspherical	32.9 × 29.4	subspherical	21.9 × 21.8
<i>C. uptoni</i>	<i>Buteo jamaicensis</i>	USA	Lindsay & Blagburn (1986, 1989)	subspherical	28.1 × 26.4	subspherical	18.2 × 17.9
<i>C. kansasensis</i>	<i>Buteo swainsoni</i>	USA	Upton et al. (1990)	ovoid	37.2 × 32.6	spherical	22.5
<i>C. lindsayi</i>	<i>B. jamaicensis</i>	USA	Upton et al. (1990)	subspherical	33.7 × 31.6	subspherical	19.2–22.0
<i>C. aquilae</i>	<i>Aquila chrysaetos</i>	Czech Republic	Volf et al. (2000)	subspherical to ellipsoidal	43.0 × 37.5	subspherical	23.8 × 23.3
<i>C. circi</i>	<i>Circus aeruginosus</i>	Czech Republic	Volf et al. (2000)	ovoid	24.5 × 21.8	subspherical	16.2 × 15.6
<i>C. hanebrinki</i>	<i>Haliaeetus leucocephalus</i>	USA	McAllister et al. (2013a)	ellipsoidal to ovoidal	48.1 × 42.1	spherical	24.8
<i>C. petersoni</i>	<i>Accipiter striatus</i>	USA	McAllister et al. (2013b)	subspherical	43.1 × 39.8	subspherical to spherical	23.4 × 23.3

Table 2. Comparative data of *Caryospora* spp. described from Falconidae (Falconiformes).

Species	Host	Locality	Reference	Oocyst		Sporocyst	
				Shape	Size (μm)	Shape	Size (μm)
<i>C. falconis</i>	<i>Falco peregrinus</i> ; <i>Falco subbuteo</i> ; <i>Falco tinnunculus</i>	Europe	Upton et al. (1990); Alfaleh et al. (2013)	subspherical	29.5 × 36.5	subspherical	21.0 × 23.0
<i>C. kutzeri</i>	<i>Falco biarmicus</i> ; <i>Falco cherrug</i> ; <i>Falco jugger</i> ; <i>Falco mexicanus</i>	Europe	Upton et al. (1990); Alfaleh et al. (2013)	subspherical	38.7 × 34.1	ovoid	24.6 × 21.0
<i>C. neofalconis</i>	<i>F. biarmicus</i> ; <i>F. mexicanus</i> ; <i>F. subbuteo</i> ; <i>F. tinnunculus</i> ; <i>F. peregrinus</i>	Europe; Mexico	Upton et al. (1990); Alfaleh et al. (2013); Santana-Sánchez et al. (2015)	subspherical	27.0 × 23.8	ovoid	18.8 × 14.8
<i>C. boeri</i>	<i>F. tinnunculus</i>	Europe	Alfaleh et al. (2013)	subspherical	36.6 × 33.4	ovoid	27.8 × 19.6
<i>C. megafalconis</i>	<i>F. biarmicus</i> ; <i>F. mexicanus</i> ; <i>F. peregrinus</i> ; <i>F. subbuteo</i>	Europe	Alfaleh et al. (2013)	subspherical or ovoid	43.6 × 35.8	spherical	23.8
<i>C. biarmicus</i>	<i>F. biarmicus</i>	Saudi Arabia	Alyousif et al. (2011); Alfaleh et al. (2013)	ovoid	40.2 × 34.7	spherical	20.1
<i>C. cherrughi</i>	<i>F. cherrug</i>	Saudi Arabia	Alfaleh et al. (2013)	ovoid	32.1 × 29.3	ellipsoidal	24.1 × 19.6
<i>C. peneireiroi</i>	<i>F. tinnunculus</i>	Portugal	current work	ellipsoidal	47.1 × 37.6	subspherical	25.1 × 24.3

These geographical ranges and the reports of *Caryospora* spp. from *F. tinnunculus* and other *Falco* spp. emphasize the potential transmission and dispersion of these coccidians in Europe, Asia and Africa. Moreover, this coccidian dispersion may be worldwide if we consider that some specimens of *F. tinnunculus* are vagrant in the Americas and that some *Falco* spp. are native in both the New and the Old World, for example: *Falco columbarius* Linnaeus, 1758, *Falco peregrinus* Tunstall, 1771, and *Falco rusticolus* Linnaeus, 1758. This assumption of coccidian dispersion across species of *Falco* and subsequently of Falconidae is grounded in the concept of intra-familial specificity, which was suggested by Duszynski & Wilber (1997) and has been reaffirmed in several subsequent studies on coccidians in birds (BERTO et al., 2011).

In this regard, the species described in the current study was primarily compared with the *Caryospora* spp. described from Falconidae, which is the single family of Falconiformes. However, since falconids are close to accipitrids, and because earlier studies on *Caryospora* spp. in raptors compared coccidians through descriptions from these two families (UPTON et al., 1986, 1990), *C. peneireiroi* was also compared in the current work with *Caryospora* spp. described from Accipitridae.

Therefore, based on the morphological features described above, *C. peneireiroi* is considered to be new to science and to be the fourth caryosporan species reported from *F. tinnunculus*.

Acknowledgements

We thank the staff at the LxCRAS facility at Monsanto Forest Park, Lisbon, Portugal, who enabled the collection of samples from birds held for rehabilitation and reintroduction into the wild. This study was supported by grants from the Fundação Carlos Chagas Filho de Amparo à Pesquisa do Estado do Rio de Janeiro (FAPERJ), Conselho Nacional de Desenvolvimento Científico e

Tecnológico (CNPq), Fundação Nacional de Desenvolvimento do Ensino Superior Particular (FUNADESP) and Ibero-American Scholarship Programme for Teachers and Researchers Santander Universities.

References

- Alfaleh FA, Alyousif MS, Al-Shawa YR, Al-Quraishy S. *Caryospora cherrughi* sp. n. (Apicomplexa: Eimeriidae) infecting *Falco cherrug* in Saudi Arabia. *Parasitol Res* 2013; 112(3): 971-974. <http://dx.doi.org/10.1007/s00436-012-3217-y>. PMid:23224640.
- Alyousif MS, Alfaleh FA, Al-Shawa YR. *Caryospora biarmiculus* sp. n. (Apicomplexa: Eimeriidae) infecting falcons from the genus *Falco* in Saudi Arabia. *J Egypt Soc Parasitol* 2011; 41(1): 1-6. PMid:21634236.
- Berto BP, Flausino W, McIntosh D, Teixeira-Filho WL, Lopes CWG. Coccidia of New World passerine birds (Aves: Passeriformes): a review of *Eimeria* Schneider, 1875 and *Isospora* Schneider, 1881 (Apicomplexa: Eimeriidae). *Syst Parasitol* 2011; 80(3): 159-204. <http://dx.doi.org/10.1007/s11230-011-9317-8>. PMid:22002022.
- Berto BP, McIntosh D, Lopes CWG. Studies on coccidian oocysts (Apicomplexa: Eucoccidioridae). *Rev Bras Parasitol Vet* 2014; 23(1): 1-15. <http://dx.doi.org/10.1590/S1984-29612014001>. PMid:24728354.
- Birdlife International. [online]. 2015 [cited 2016 Jan 30]. Available from: <http://www.birdlife.org>
- Duszynski DW, Wilber P. A guideline for the preparation of species descriptions in the Eimeriidae. *J Parasitol* 1997; 83(2): 333-336. <http://dx.doi.org/10.2307/3284470>. PMid:9105325.
- Lindsay DS, Blagburn BL. *Caryospora uptoni* and *Frenkelia* sp. -like coccidian infections in red-tailed hawks (*Buteo borealis*). *J Wildl Dis* 1989; 25(3): 407-409. <http://dx.doi.org/10.7589/0090-3558-25.3.407>. PMid:2503626.

- Lindsay DS, Blagburn BL. *Caryospora uptoni* n. sp. (Apicomplexa: Eimeriidae) from red-tailed hawks (*Buteo jamaicensis borealis*). *J Parasitol* 1986; 72(5): 762-765. <http://dx.doi.org/10.2307/3281470>. PMid:3806324.
- McAllister CT, Duszynski DW, McKown RD. A new species of *Caryospora* (Apicomplexa: Eimeriidae) from the bald eagle, *Haliaeetus leucocephalus* (Accipitriformes: Accipitridae), from Kansas. *J Parasitol* 2013a; 99(2): 287-289. <http://dx.doi.org/10.1645/GE-3236.1>. PMid:22992168.
- McAllister CT, Duszynski DW, McKown RD. A new species of *Caryospora* (Apicomplexa: Eimeriidae) from the sharp-shinned hawk, *Accipiter striatus* (Aves: Accipitriformes). *J Parasitol* 2013b; 99(3): 490-492. <http://dx.doi.org/10.1645/GE-3228.1>. PMid:23098064.
- Palma L. The present situation of birds of prey in Portugal. In: Newton I, Chancellor RD. *Conservation studies on raptors*. Cambridge: Technical Publication; 1985. p. 3-14.
- Papazahariadou MG, Georgiades GK, Komnenou AT, Ganoti M. *Caryospora* species in a snowy owl (*Nyctea scandiaca*). *Vet Rec* 2001; 148(2): 54-55. <http://dx.doi.org/10.1136/vr.148.2.54>. PMid:11202556.
- Santana-Sánchez G, Flores-Valle IT, González-Gómez M, Vega-Sánchez V, Salgado-Miranda C, Soriano-Vargas E. *Caryospora neofalconis* and other enteroparasites in raptors from Mexico. *Int J Parasitol Parasites Wildl* 2015; 4(3): 351-355. <http://dx.doi.org/10.1016/j.ijppaw.2015.08.004>. PMid:26543806.
- Upton SJ, Campbell TW, Weigel M, McKown RD. The Eimeriidae (Apicomplexa) of raptors: Review of the literature and description of new species of the genera *Caryospora* and *Eimeria*. *Can J Zool* 1990; 68(6): 1256-1265. <http://dx.doi.org/10.1139/z90-187>.
- Upton SJ, Current WL, Barnard SM. A review of the genus *Caryospora* Léger, 1904 (Apicomplexa: Eimeriidae). *Syst Parasitol* 1986; 8(1): 3-21. <http://dx.doi.org/10.1007/BF00010305>.
- Upton SJ, Odell DK, Walsh MT. Description of the oocysts of *Caryospora undata* (Apicomplexa) from tufted puffins, *Lunda cirrhata* (Charadriiformes). *Tex J Sci* 1992; 44(1): 79-82.
- Village A. *The Kestrel*. London: T. & A.D. Poyser; 1990.
- Volf J, Koudela B, Modrý D. Two new species of *Caryospora* Léger, 1904 (Apicomplexa, Eimeriidae) from accipitrid raptors. *Syst Parasitol* 2000; 46(1): 23-27. <http://dx.doi.org/10.1023/A:1006227911679>. PMid:10803432.
- Yang R, Brice B, Ryan U. A new *Caryospora* coccidian species (Apicomplexa: Eimeriidae) from the laughing kookaburra (*Dacelo novaeguineae*). *Exp Parasitol* 2014; 145: 68-73. <http://dx.doi.org/10.1016/j.exppara.2014.07.008>. PMid:25092438.