

Organization and variation of mitochondrial DNA control region in pleurodiran turtles

Ling Wang¹; Xuming Zhou² & Liuwang Nie^{1, 3}

¹ Life Science College, Anhui Normal University, The Provincial Key Lab of the Conservation and Exploitation Research of Biological Resources in Anhui, 1 East Beijing Road, Wuhu, Anhui, 241000, China.

² Key laboratory of Animal Ecology and Conservation Biology, Institute of Zoology, Chinese Academy of Sciences, Beijing 100101, China.

³ Corresponding author: E-mail: lwnie@mail.ahnu.edu.cn

ABSTRACT. Three complete mitochondrial DNA (mtDNA) control regions (CRs) of *Chelodina rugosa* (Ogilby, 1890), *Chelus fimbriata* (Schneider, 1783), and *Podocnemis unifilis* (Troschel, 1848) were firstly determined using Long-PCR method and the length were 1,016 bp, 1,149 bp, and 985bp, respectively. Together with CRs of *Pelomedusa subrufa* (Bonnaterre, 1789) and nearly complete CR of *Podocnemis expansa* (Schweigger, 1812) obtained from GenBank, the structural and evolutionary characteristics of mtDNA CRs in pleurodiran turtle were analyzed in this study. We identified three functional domains (TAS, CD, and CSB domains) as well as their conservation sequences (TAS, CSB-F, and CSB-1) according to their homology to those of other turtles. Within the TAS domain, an interrupted poly-C stretch was found in *C. rugosa*, *C. fimbriata*, and *P. subrufa*, which also exists in the published mt DNA CRs of *Chrysemys picta* (Schneider, 1783), *Trachemys scripta* (Thunberg in Schoepff, 1792), and *Trionyx triunguis* (Forskål, 1775). The analysis of the origin for the poly-C sequences in TAS domain from six turtles suggested that the poly-C sequences are more related to “goose hairpin” in birds rather than CSB2 in CSB domain. In the CSB domain, CSB2 and CSB3, which were determined in CRs of Cryptodira, were absent in Pleurodira CRs, indicating the regulative mechanisms of transcription may be varied in both two suborders and the lack of CSB2 and CSB3 could be proposed as one of diagnostic characters between Pleurodira and Cryptodira at molecular level. As for CR of other cryptodiran turtles, variable number of tandem repeats (VNTRs) in the 3' end of the CRs was found in the five pleurodiran turtles. Interestingly, the long repeated motifs from each species could form stable stem-loop secondary structures, suggesting that the repeated sequences may play an important role in regulating replication of the mitochondrial genome in Pleurodiran, and the secondary structures of VNTRs may provide some potential information in phylogenetic inference.

KEY WORDS. Control region; CSB; mitochondrial DNA; molecular evolution; Pleurodira.

The mitochondrial genome is highly conserved and compact, usually including genes that code for 13 proteins, 22 tRNAs and two rRNA, as well as an important noncoding sequence (control region, CR) in vertebrates. Particularly, the CR in vertebrate mitochondrial genomes contains two promoters (LSP and HSP) for the transcription, heavy-strand replication origin (O_H), and the displacement loop (D-loop) (CLAYTON 1982, CHANG & CLAYTON 1986). However, based on the distribution of the variable nucleotide positions and different nucleotide frequencies, the mitochondrial CR is divided into three domains (BROWN *et al.* 1986, SACCONI *et al.* 1991): termination associated sequence (TAS) domain, central conserved domain (CD), and conserved sequence block (CSB) domain. The TAS domain has been shown to contain sequences associated with termination of newly synthesized H-strands during replication (DODA *et al.* 1981, BROWN *et al.* 1986, SBISÀ *et al.* 1997). The CD, which contains several areas of highly

conserved sequences, is more conserved with respect to the TAS and CSB domains. However, the nature of these conserved sequences varies among vertebrate classes. For example, ANDERSON *et al.* (1981) identified several conserved sequence boxes (CSB), B, C, D, E, and F in the CD of the human mitochondrial CR. Of those, only three (CSB B, CSB D, and CSB F) can be found in the avian mitochondrial CD (RUKONEN & KVIST 2002). The CSB domain usually contains the origin of the H-strand transcription (WALBERG & CLAYTON 1981, BROWN *et al.* 1986, KING & LOW 1987, FORAN *et al.* 1988). In many taxa, the TAS and CSB domains, which are more variable than the CD, have variable numbers of tandem repeats (VNTRs) (BROUGHTON & DOWLING 1994, SBISÀ *et al.* 1997, ZARDOYA & MEYER 1998a, DELPORT *et al.* 2002, FU *et al.* 2006, ZHANG *et al.* 2009, XIONG *et al.* 2010).

Extant turtles have been divided into two monophyletic clades, Pleurodira and Cryptodira. However, virtually all stud-

ies on turtle mitochondrial genomes have focused on Cryptodira. Studies comparing the mitochondrial CRs of pleurodiran turtles have been limited because there is only one published complete CR from one species – *Pelomedusa subrufa* (Bonnaterre, 1789). In the present study, three complete CR of three pleurodiran turtles – *Chelodina rugosa* (Ogilby, 1890), *Chelus fimbriata* (Schneider, 1783), and *Podocnemis unifilis* (Troschel, 1848) – representing two families (Chelidae and Podocnemididae) are characterized. Additionally, together with the complete CR of *P. subrufa* and a nearly complete CR of *Podocnemis expansa* (Schweigger, 1812) obtained from GenBank, we have compared the CR sequences of pleurodiran turtles with the CRs of other vertebrates. Additionally, the features which are shared or not among the pleurodiran turtles and other vertebrates have been identified and are discussed in detail.

MATERIAL AND METHODS

Sample and sequencing

Specimens of *C. rugosa*, *C. fimbriata*, and *P. unifilis* were stored at the Anhui Normal University. Total genomic DNA was extracted from their muscles with the proteinase K method (SAMBROOK & RUSSELL 2001) and kept at -20°C for PCR amplification.

The mitochondrial CR was amplified by long-PCR. The entire CR and 3 tRNAs (tRNA^{Thr}, tRNA^{Pro}, and tRNA^{Phe}) as well as partial Cyt b and 12S rRNA gene sequences were amplified in one single step using a pair of long-PCR primers:

LCR-F: 5'-CTTCCTATTTGCTATGCTATC-3'

LCR-R: 5'-TATTTGGGCTCCTGGTGTA-3'

Long-PCR conditions were: one minute at 95°C, then 30 cycles of 10 seconds at 98°C, five minutes at 55°C, followed by a final extension for 10 minutes at 72°C. PCR products were isolated using a Gel Extract Purification Kit (TaKaRa Co., Ltd, Dalian, China) after 1% agarose gel electrophoresis. The purified Products were sequenced with an ABI3730 automated sequencer.

Sequence analysis

In order to determine the complete CR sequence of the three turtles, the sequences obtained for *C. rugosa*, *C. fimbriata*, and *P. unifilis* were compared with the complete mtDNA of *P. subrufa* (ZARDOYA & MEYER 1998b), and the tRNAs were identified using tRNAscan-SE 1.21 (LOWE & EDDY 1997). The sequence containing almost complete CRs of *P. expansa* was retrieved from GenBank. Subsequently, five pleurodiran turtle sequences were aligned using ClustalX 1.8 software (THOMPSON *et al.* 1997) and then checked manually in order to define the conserved sequence blocks.

After comparison with published data from other taxa (the sequences used are listed in the Appendix), the conserved box F (CSB F) was delimited in the aligned sequences. The conserved block 1 (CSB1) with the characteristic motif GACATA was also delimited. The boundary of the TAS domains and the Central conserved domain was the starting point of the CSB F.

The CSB domain was always set to start with the conserved sequence box 1 (CSB1) (ZHANG *et al.* 2009, XIONG *et al.* 2010). The program 'Tandem Repeats Finder' (BENSON 1999) was used in order to identify the VNTRs in the CRs. Furthermore, putative secondary structures in the CRs were determined using the software RNA structure (MATHEWS *et al.* 1999). Subsequently, the computer program RNAdraw (HOFACKER *et al.* 1995) was employed to prepare secondary structures for publication.

RESULTS AND DISCUSSION

The length and base composition of pleurodiran turtle CRs

The CRs of *C. rugosa*, *C. fimbriata*, and *P. unifilis* have 1,016 bp, 1,149 bp, and 985bp, respectively. They are rich in adenine and thymine and lack L strand guanines, which is evident in each domain, particularly in the CSB domain (Tab. I). However, the composition of the CR among the three domains is not uniform. The Central conserved domain is poor in adenine and rich in L strand guanines compared to the flanking TAS and CSB domains. Interestingly, among the five turtles compared, the TAS domain was found to be richer in adenine and thymine in three species: *C. rugosa*, *C. fimbriata*, and *P. subrufa*. This TAS domain composition is consistent with that of cryptodiran turtles (ZHANG *et al.* 2009, XIONG *et al.* 2010). However, in the other two pleurodiran turtles (*P. unifilis* and *P. expansa*), the TAS domain was rich in cytosine and thymine.

The organization of turtle CRs

Like in most cryptodiran turtles, the CRs in the mitochondrial genomes of pleurodiran turtles are located between the tRNA^{Pro} and tRNA^{Phe} genes. The CSB F and CSB-1 were easy to define in the five turtles (Fig. 1). In the CSB F, 12 of 29 (41.4%) nucleotide positions were fixed among the five turtles. In the CSB1, 9 of 20 (45%) nucleotide positions were fixed. Comparing with the cryptodiran turtles (ZHANG *et al.* 2009, XIONG *et al.* 2010), the conserved blocks of pleurodiran turtles varied greatly. This variation may due to the age of the group. The fossil record has shown that most pleurodiran turtles diverged about 100–150 mya (million years ago) (GAFFNEY 1990). During this long evolutionary history, the conserved blocks sequences of the mtDNA CRs might have changed independently, accumulating more variation than the CRs of cryptodiran turtles, in which most species have diverged after 100 mya (NEAR *et al.* 2005).

TAS domain

The TAS domain was located between the 5' end of the CR and the beginning of the CSB F. The length of the TAS domain varied from 243 bp to 357 bp, and the sequences were heterogeneous and could not be unambiguously aligned. However, the TAS sequences were easily determined and were identical among the pleurodiran and cryptodiran turtles (Fig. 1).

An interrupted poly-C stretch was found in the TAS domain of *C. rugosa* and *C. fimbriata*. This poly-C stretch also

Species	TAS	CSBF	CSB1
<i>Chelodina rugosa</i>	TACAT	AGC---TATCTTCGAGAAATCACCAATCCTTG	TTA-ATGTTTGTAAGACATAT
<i>Chelus fimbriata</i> ---.. G. C. T. A. C-. . . A. . AGT. . G.
<i>Pelomedusa subrufa</i> ---.. CTCA. . T. . . CATGA. . T. . . C. - . . . C. CAC. G. A
<i>Podocnemis unifilis</i> CCC. . TC. . AT. . . . C. . . . TC. . CAC	A. CC. . . CAC. C. G. A
<i>Podocnemis expansa</i> CCC. . . C. . A. C. . G. . TC. . CCT	... - . . . C. C. ACG. A
	*****	*** ** * *** ** *	* *** * ****

Figure 1. The alignment of TAS and conserved blocks from five pleurodiran turtles. Note: Dots indicate same nucleotides as top sequences, and asterisk shows the identical sites.

Table I. The base composition (%) and length of the five pleurodiran turtle CRs.

Species	Complete CR					TAS domain				
	T	C	A	G	Total	T	C	A	G	Total
<i>Chelodina rugosa</i>	32.00	19.9	35.00	13.10	1016.0	32.4	22.50	33.80	11.3	275.0
<i>Chelus fimbriata</i>	31.80	19.8	37.60	10.90	1149.0	28.6	24.90	32.70	13.9	245.0
<i>Pelomedusa subrufa</i>	34.80	18.7	37.90	8.60	1194.0	32.9	25.90	27.20	14.0	243.0
<i>Podocnemis unifilis</i>	32.40	26.0	28.90	12.70	985.0	31.4	32.50	24.60	11.5	357.0
<i>Podocnemis expansa</i>	32.30	25.1	29.80	12.80	1113.0	28.2	33.30	27.10	11.3	354.0
Average	32.66	21.9	33.84	11.62	1091.4	30.7	27.82	29.08	12.4	294.8

Species	Central conserved domain					CSB domain				
	T	C	A	G	Total	T	C	A	G	Total
<i>Chelodina rugosa</i>	37.00	19.20	26.90	16.90	349.0	27.30	18.60	43.10	11.0	392
<i>Chelus fimbriata</i>	37.70	19.00	27.50	15.90	353.0	29.40	18.00	46.30	6.4	551
<i>Pelomedusa subrufa</i>	34.70	24.80	25.30	15.20	363.0	35.70	11.90	50.00	2.4	588
<i>Podocnemis unifilis</i>	32.80	25.30	24.20	17.60	363.0	33.20	18.10	41.10	7.5	265
<i>Podocnemis expansa</i>	32.40	24.80	24.80	18.10	420.0	36.60	16.80	38.90	7.7	339
Average	34.92	22.62	25.74	16.74	369.6	32.44	16.68	43.88	7.0	427

Note: In the CR of *P. expansa*, the TAS and Central conserved domain are complete, and CSB domain is almost complete.

occurs in four other cryptodiran turtles: *Chrysemys picta* (Schneider, 1783), *Trachemys scripta* (Thunberg in Schoepff, 1792), *Trionyx triunguis* (Forskål, 1775), and *P. subrufa* (ZARDOYA & MEYER 1998a). It is one remarkable feature of the TAS domain sequence in these turtles. The interrupted poly-C stretch is repeated once in *C. picta* and twice in *T. scripta*. The similar poly-C stretch sequence in many birds (QUINN & WILSON 1993, RANDI & LUCCHINI 1998, RITCHIE & LAMBERT 2000) and crocodiles (RAY & DENSMORE 2002) is referred to as “goose hairpin”. In birds, it could potentially form a stable hairpin structure, which is usually characterized by several consecutive cytosine residues separated from the complementary guanine by a putative stop sequence (TCCC). The poly-C stretch in crocodiles (*Crocodylus*) was reported as having a region with high cytosine content that could form a similar stem-loop structure. However, in the present study, the only poly-C stretches that have the capacity to form a similar secondary structure are those of *C. fimbriata*

and *P. subrufa* (data not shown). ZARDOYA & MEYER (1998a) reported that the interrupted poly-C of the TAS domain in *P. subrufa* was remarkably similar to the CSB2. In order to discuss the origin of these similar poly-C sequences, we performed clustering analyses using the Neighbor-joining (NJ) method on the poly-C sequences of the TAS domain of the six turtles, the CSB2s from 31 turtles, and the “goose hairpin” sequence of *Gallus gallus* (Linnaeus, 1758). Interestingly, the sequences grouped into two major clades, A and B (Fig. 2). All CSB2s sequences grouped within Clade A. Though the relationships among these CSB2 may not represent the “species tree” of turtles, the fact that they clustered indicates that the CSB2 of these turtles is conserved. Clade B consists of the poly-C sequences in the TAS domain from the six turtles and the “goose hairpin” sequence from *G. gallus*. Clades A and B indicate that the origin of the poly-C sequences in the TAS domain were closer to “goose hairpin” in the TAS domain rather than the CSB2.




Figure 2. NJ tree based on the CSB2s and poly-C sequences. The tree was reconstructed according to SAITOU & NEI (1987) using the absolute number of differences in MEGA (KUMAR *et al.* 1993). Sequences used in the analysis are listed in the Appendix. A total of 1000 bootstrap replicates were performed and values greater than 50 are indicated at the respective nodes. Clade A consists of species in which CSB2 is located within the CSB domain, whereas clade B contains the species whose poly-C sequences are within the TAS domain.


Central conserved domain

The Central conserved domain (CD) is more conserved than the peripheral domains (TAS and CSB domains). It is 353–420bp long in the five turtles. Our resulting alignment had 102 conserved sites, 273 variable sites, and 127 parsimony-informative sites. Although the extreme conservation of this region in vertebrates may suggest that it has played an important role in the evolutionary history of the mitochondrial genome, its function remains completely unknown. On other hand, the high degree of similarity makes this domain suitable for phylogenetic inference (SACCONE *et al.* 1991). However, we failed to reconstruct the phylogenetic relationships based on it. We believe that the phylogenetic distances in this domain are subject to high statistical fluctuations among family-level taxa of turtles because of the reduced number of sites in this region (353–420bp). Thus, the phylogenetic relationships of the five pleurodiran turtles reconstructed based on this domain were not well resolved.

CSB domain

The CSB domain, ranging from 265bp (*P. unifilis*) to 588bp (*P. subrufa*), has several conserved blocks involved in the regulation of replication and transcription. These short conserved sequence blocks were firstly called CSB1, CSB2, and CSB3 by WALBERG & CLAYTON (1981). Researchers have proposed that they provide regulatory signals for the processing of the RNA primers for replication of the H-strand in human KB and mouse L cells. Since then, these three conserved sequence blocks have been identified in a number of vertebrate mtDNAs, such as fish (BUROKER *et al.* 1990, BROUGHTON & DOWLING 1994, LIU 2002), amphibians (DUNON-BLUTEAU *et al.* 1985), lizards (BREHM *et al.* 2003), turtles (SERB *et al.* 2001, ZHANG *et al.* 2009, XIONG *et al.* 2010), and mammals (SBISÀ *et al.* 1997). However, it is interesting that in turtles only the CSB-1 is found consistently, whereas in marsupials, all three CSBs are consistently present. In placental mammals, CSB-2 and CSB-3 are sometimes missing (NILSSON 2009). Thus, there seems to be a lot of plasticity across vertebrates concerning the CSB1, CSB-2 and CSB-3. Particularly, the CSB1 was found to be a basic element in the CR of the mtDNA of all vertebrates and duplications of the CSB1 have been found in sperm whales, shrews, hedgehogs, opossums (SBISÀ *et al.* 1997), and hornbills (SBISÀ *et al.* 1997, DELPORT *et al.* 2002). A sequence such as GACATA, is identical among several vertebrate mtDNA CSB-1 sequences, from fish to mammals, and birds (BUROKER *et al.* 1990, SBISÀ *et al.* 1997, RUOKONEN & KVIST 2002, NILSSON 2009). The conserved sequences and structure of CSB-1 may have important functions. For example, GHIVIZZANI *et al.* (1994) have reported that the origin of the heavy strand replication (OH) and transition from RNA transcription to DNA replication should occur in the proximity of the CSB-1.

The other two conserved sequence blocks, CSB2 and CSB3, are believed to play an important role in RNA processing during the transcription in mice (CHANG & CLAYTON 1986). In humans, the CSB2 and CSB3 may have another function: GHIVIZZANI *et al.* (1994) have proposed that together with C residues, they appear to function as domains which avoid interaction with the mitochondrial transcription factor A (mtTFA), thus reinforcing the organization of the mtTFA binding. Although they appear to have an important function in different taxa, both the CSB2 and CSB3 are not present in the CRs of the mtDNA of all vertebrates. In this study, the CSB2 and CSB3 were missing in all five pleurodiran turtles. The same condition has been reported in birds (RAMIREZ *et al.* 1993, BAKER & MARSHALL 1997, HÄRLID *et al.* 1997, RANDI & LUCCHINI 1998) and some mammals (SBISÀ *et al.* 1997). Interestingly, all mtDNA CRs of cryptodiran turtles published so far have the CSB2 and CSB3. Thus, there are two known types of mitochondrial CR in turtles at present, which correspond to variations in the CSB domain (Figs 3 and 4). In all cryptodiran turtles, the CSB domain contains three conserved blocks (CSB-1, CSB-2, and CSB-3) (SERB *et al.* 2001, JUNGT *et al.* 2006, AMER & KUMAZAWA 2009, ZHANG *et al.* 2009, XIONG *et al.* 2010) (Fig. 3) but only one (CSB-1) can be



Figures 3-4. Two typical mtDNA CR structures published in turtles: (3) the structure of mtDNA CR in Cryptodira. VNTRs a and VNTRs b are only identified in CRs of some trionychids; (4) The structure of mtDNA CR in Pleurodira. P: tRNA-Pro gene, F: tRNA-Phe gene.

identified in the CSB region of our five pleurodiran turtles (Fig. 4). This considerable variation shows that the regulative mechanisms of transcription may be different among vertebrates or may involve specific nuclear-mitochondrial co-evolutionary processes (SBISÀ *et al.* 1997), and in particular, the mechanisms of regulation may be different between the two suborders (Pleurodira and Cryptodira) in turtles. If the lack of the CSB2 and CSB3 in the CSB region turns out to be a pervading character state in Pleurodira, it could be used as a diagnostic character between Pleurodira and Cryptodira at the molecular level.

VNTRs

VNTRs have been identified in the mtDNA CRs of many animals, including invertebrates and vertebrates (HOELZEL 1993, FUMAGALLI *et al.* 1996, GORICKI & TRONTELJ 2006, JI *et al.* 2008). In cryptodiran turtles, VNTRs are inserted in the two peripheral domains, i.e. TAS and CSB domains (Fig. 3). VNTRs are only found within the TAS domain in some trionychids, such as *Apalone ferox* (Schneider, 1783), *Dogania subplana* (Geoffroy Saint-Hilaire, 1809), *Pelodiscus sinensis* (Wiegmann, 1835), and *Palea steindachneri* (Siebenrock, 1906) (XIONG *et al.* 2010). VNTRs

located between the CSB1 and the CSB2, identified in many mammalian species (GEMMELL *et al.* 1996, SBISÀ *et al.* 1997, LARIZZA 2002, PENG *et al.* 2007, SUN *et al.* 2009), are also presented in *P. sinensis* (XIONG *et al.* 2010) and *Indotestudo forstenii* (Schlegel and Müller, 1845). As for most turtles, VNTRs were commonly found at the 3' end of the CSB domain (ZARDOYA & MEYER 1998a, SERB *et al.* 2001, JUNGT *et al.* 2006, AMER & KUMAZAWA 2009, ZHANG *et al.* 2009, XIONG *et al.* 2010). Generally, VNTRs located between the CSB1 and CSB2, or within the TAS domain, or at the 3' end of the CRs identified in most turtles (ZARDOYA & MEYER 1998a, ZHANG *et al.* 2009, XIONG *et al.* 2010) support the idea that the CR can tolerate deletion/insertion events and accumulate repeats without loss of function (SBISÀ *et al.* 1997).

The VNTRs of the new determined mtDNA CRs from the three pleurodiran turtles, and two additional pleurodiran turtles (*P. expansa* and *P. subrufa*), which were only located at the 3' end of the mtDNA, were further analyzed in this study (Tab. II). As in pleurodiran turtles, the repeated motifs of each species were not homogenous at the end of the CR. The motifs of VNTRs in five pleurodiran turtles were similar in length. The lengths of the VNTRs are closer in closely related species.

Table II. Comparison of the repeat units in CR of the five pleurodiran turtles analyzed in the present study.

Species	Repeated motif	Sequences
<i>C. rugosa</i>	(74 bp)×3	5'-ATTATACACCAAGTGCAGTGTGCGAAAGATAATTATTTTTACACGCACCAAACACTGTGCCGAAATTATATTAAC-3'
<i>C. fimbriata</i>	(76 bp)×4	5'-GAAAAACAGACTACTTTTATATAAAAAATTAATTGTTTTTATTCAAAAATTTAAACAATACTATCTTTGTC-3'
	(36 bp)×5	5'-GAAAATTAACCCTAAAACACTACCAAGTCAAGTACC-3'
<i>P. unifilis</i>	(2 bp)×19	5'-AT-3'
	(66 bp)×2	5'-CTATAAATTCCTAGACTTACTAAGAAATTTACCCTAATCGCTAATATATATACATATTAATAGC-3'
<i>P. expansa</i>	(69 bp)×4	5'-CAATAATTTATAAATTCCTAGACTTACTAAGATATTTACCTCAATTGCTAATTGTATATATATATA-3'
<i>P. subrufa</i>	(65 bp)×6	5'-CAAACATATATACAAACAAAGTACTAATTACTAATCATATAAATATATATATATATATA-3'

Potential secondary structures of the repeated sequences in the five pleurodiran turtles' CR were constructed in this study, and the long repeated motif from each species formed a stable stem-loop structure (Figs 5-9). In hornbills, these VNTRs' stem-loop structures are believed to function as a termination sequence in mitochondrial genome replication (DELPORT *et al.* 2002). In turtles, whether these VNTRs have some function or not needs further study, but it is noteworthy that the potential secondary structures of the repeated sequences were very similar among *P. unifilis*, *P. expansa*, and *P. subrufa* (Figs 7-9).

Although the mechanism for generating the VNTRs in the mitochondrial genome is not well understood at present,

it is believed that the VNTRs can be used as molecular markers and provide sufficient phylogenetic information in molecular phylogenetics, population genetics, species identification, as well as genetic diversity and conservation, because they can be used to differentiate among genera, species, populations, and even individuals (ZARDOYA & MEYER 1998a, ZHANG *et al.* 2009). Pelomedusids and podocnemidids were placed into Pelomedusidae (sensu lato) based on morphological characteristic (GAFFNEY 1975, GAFFNEY & MEYLAN 1988). Although DE BROIN (1988) suggested that the extant Pelomedusinae and Podocneminae (GAFFNEY & MEYLAN 1988) should be recognized as two families, i.e. Pelomedusidae (consisting of the extant African *Pelusios*




Figure 5-9. Secondary structure of repeat units constructed from VNTRs of the five pleurodiran turtles: (5) the 74 bp repeat unit of *C. rugosa*; (6) the 76 bp repeat unit of *C. fimbriata*; (7) the 65 bp repeat unit of *P. subrufa*; (8) the 69 bp repeat unit of *P. expansa*; (9) the 66 bp repeat unit of *P. unifilis*.

and *Pelomedusa*) and Podocnemidae (consisting of extant Madagascan *Erymnochelys*, and the extant South American *Podocnemis* and *Peltocephalus*), morphological studies still support the monophyly of Pelomedusidae (sensu lato). Recently, molecular evidence has also suggested that the Pelomedusidae (sensu lato) may be monophyletic (GEORGES *et al.* 1998, NOONAN 2000, FUJITA *et al.* 2004, NOONAN & CHIPPINDALE 2006, THOMSON & SHAFFER 2010). In the present study, these secondary structures of VNTRs for *P. subrufa* (Pelomedusinae), *P. expansa*, and *P. unifilis* (Podocneminae) also supports the monophyly of Pelomedusidae (sensu lato). We suggest that comparative analyses of secondary structures of VNTRs may provide some potential information for phylogenetic inferences.

ACKNOWLEDGMENTS

This research was supported by the National Natural Science Foundation of China (NSFC, #30970351), the Research Fund for the Doctoral Program of Higher Education of China (#20080370001), and the Key Laboratory of Biotic Environment and Ecological Safety of Anhui province.

LITERATURE CITED

- AMER, S.A. & Y. KUMAZAWA. 2009. Complete sequence of the mitochondrial genome of the endangered Nile soft-shelled turtle *Trionyx triunguis*. *Egyptian Journal of Experimental Biology (Zoology)* 5: 43-50.
- ANDERSON, S.; A.T. BANKIER; B.G. BARRELL; M.H.L. DE BRUIJN; A.R. CNULSON; J. DROUIN; I.C. EPERON; D.P. NIERLICH; B.A. ROE; F. SANGER; P.H. SCHREIER; A.J.H. SMITH; R. STADEN & I.G. YOUNG. 1981. Sequence and organization of the human mitochondrial genome. *Nature* 290: 457-465.
- BAKER, A.J. & H.D. MARSHALL. 1997. Mitochondrial control region sequences as tools for understanding evolution, p. 51-82. *In*: D.P. MINDELL (Ed.). *Avian Molecular Evolution and Systematics*. San Diego, Academic Press.
- BENSON, G. 1999. Tandem repeats finder: a program to analyze DNA sequences. *Nucleic Acids Research* 27 (2): 573-580.
- BREHM, A.; A.D. JAMES HARRIS; C.D. ALVES; J.D. JESUS; F.D. THOMARAT & L.D. VICENTE. 2003. Structure and evolution of the mitochondrial DNA complete control region in the lizard *Lacerta dugesii* (Lacertidae, Sauria). *Journal of Molecular Evolution* 56 (1): 46-53.
- BROUGHTON, R.E. & T.E. DOWLING. 1994. Length variation in mitochondrial DNA of the minnow *Cyprinella spiloptera*. *Genetics* 138 (1): 179-190.
- BROWN, G.G.; G. GADALETA; G. PEPE; C. SACCONI & E. SBISA. 1986. Structural conservation and variation in the D-loop containing region of vertebrate mitochondrial DNA. *Journal of Molecular Biology* 192: 503-511.
- BUROKER, N. E.; J.R. BROWN; T.A. GILBERT; P.J. O'HARA; A.T. BECKENBACH; W.K. THOMAS & M.J. SMITH. 1990. Length Heteroplasmy of Sturgeon Mitochondrial DNA: An Illegitimate Elongation Model. *Genetics* 144: 157-163.
- CHANG, D. & D. CLAYTON. 1986. Identification of primary transcription start sites of mouse mitochondrial DNA: Accurate in vitro initiation of both heavy- and light-strand transcriptions. *Molecular Cellular Biology* 6: 1446-1453.
- CLAYTON, D.A. 1982. Replication of animal mitochondrial DNA. *Cell* 28: 693-705.
- DE BROIN, F. 1988. Les Tortues et le Gondwana. Examen des rapports entre le fractionnement du Gondwana et la dispersion géographique des tortues pleurodirées à partir du Crétacé. *Studia Geologica Salmantica* 2 (2): 103-142.
- DELPORT, W.; J.W. FERGUSON & P. BLOOMER. 2002. Characterization and evolution of the mitochondrial DNA control region in hornbills (Bucerotiformes). *Journal of Molecular Evolution* 54: 794-806.
- DODA, J.N.; C.T. WRIGHT & D.A. CLAYTON. 1981. Elongation of displacement-loop strands in human and mouse mitochondrial DNA is arrested near specific template sequences. *Proceeding of the National Academy of Sciences of the United States of America* 78: 6116-6120.
- DUNON-BLUTEAU, D.; M. VOLOVITCH & G. BRUN, 1985. Nucleotide sequence of a *Xenopus laevis* mitochondrial DNA fragment containing the D-loop, flanking tRNA genes and apocytobrome B gene. *Gene* 36: 65-78.
- FORAN, D.R.; J.E. HIXSON & W.M. BROWN. 1988. Comparisons of ape and human sequences that regulate mitochondrial DNA transcription and D-loop synthesis. *Nucleic Acids Research* 16: 5841-5861.
- FU, Y.Y.; W.W. GU; Y.Z. LIU; J.Y. PENG & H. CHANG. 2006. Genetic analysis of the polymorphism of mtDNA D loop and microsatellite loci in Tibet mini-pigs. *Acta Laboratorium Animalis Scientia Sinica* 14: 318-321.
- FUJITA, M.K.; T.N. ENGSTROM; D.E. STARKEY & H.B. SHAFFER. 2004. Turtle phylogeny: insights from a novel nuclear intron. *Molecular Phylogenetics Evolution* 31: 1031-1040.
- FUMAGALLI, L.; P. TABERLET; L. FAVRE & J. HAUSSER. 1996. Origin and evolution of homologous repeated sequences in the mitochondrial DNA control region of shrews. *Molecular Biology Evolution* 13: 31-46.
- GAFFNEY, E.S. 1975. A phylogeny and classification of the higher categories of turtles. *Bulletin of the American Museum of Natural History* 155: 391-436.
- GAFFNEY, E.S. 1990. The comparative osteology of the Triassic turtle *Proganochelys*. *Bulletin of the American Museum of Natural History* 194: 1-263.
- GAFFNEY, E.S. & P.A. MEYLAN. 1988. A phylogeny of turtles, p. 157-219. *In*: M.J. BENTON (Ed.). *The phylogeny and classification of Tetrapods. 1. Amphibians, Reptiles, Birds*. New York, Systematics Association Special, vol. 35A.
- GEMMELL, N.J.; P.S. WESTERN; J.M. WATSON & J.A. GRAVES. 1996. Evolution of the mammalian mitochondrial control region – Comparisons of control region sequences between mono-

- treme and therian mammals. **Molecular Biology Evolution** 13: 798-808.
- GEORGES, A.; J. BIRRELL; K.M. SAINT; W. MCCORD & S.C. DONNELLAN. 1998. A phylogeny for side-necked turtles (Chelonia: Pleurodira) based on mitochondrial and nuclear gene sequence variation. **Biological Journal of Linnnean Society** 67: 213-246.
- GHIVIZZIANI, S.C.; C.S. MADSEN; M.R. NELEN; C.V. AMMINI & W.W. HAUSWIRTH. 1994. In organelle foot print analysis of human mitochondrial DNA: Human mitochondrial transcription factor A interactions at the origin of replication. **Molecular Cellular Biology** 14: 7717-7730.
- Goricki, S. & P. TRONTELJ. 2006. Structure and evolution of the mitochondrial control region and flanking sequences in the European cave salamander *Proteus anguinus*. **Gene** 378: 31-41.
- HÄRLID, A.; A. JANKE & U. ARNASON. 1997. The mtDNA sequence of the ostrich and the divergence between paleognathous and neognathous birds. **Molecular Biology Evolution** 14: 754-761.
- HOELZEL, A.R. 1993. Evolution by DNA turnover in the control region of vertebrate mitochondrial DNA. **Current Opinion in Genetics & Development** 3: 891-895.
- HOFACKER, I.L.; P.F. STADLER; S. BONHOEFFER & W. FONTANA. 1995. **The Vienna RNA package: A comprehensive package for RNA secondary structure calculation and analysis for Windows95/NT3.51**. Available online at: <http://www.tbi.univie.ac.at/RNA> [Accessed: 04/VIII/2011].
- Ji, X.; X. WU; P. YAN & G. AMATO. 2008. Complete sequence and gene organization of the mitochondrial genome of Siamensis Crocodile (*Crocodylus siamensis*) **Molecular Biology Reports** 35: 133-138.
- JUNGT, S.O.; Y.M. LEE; Y. KARTAVTSEV; I.S. PARK; D.S. KIM & J.S. LEE. 2006. The complete mitochondrial genome of the Korean soft-shelled turtle *Pelodiscus sinensis* (Testudines, Trionychidae). **DNA Sequence** 17 (6): 471-483.
- KING, T.C. & R.L. LOW. 1987. Mapping of control elements in the displacement loop region of bovine mitochondrial DNA. **Journal of Biological Chemistry** 262: 6204-6213.
- KUMAR, S.; K. TAMURA & M. NEI. 1993. **MEGA: Molecular Evolutionary Genetics Analysis**. Pennsylvania State University, University Park.
- LARIZZA, A. 2002. Lineage Specificity of the Evolutionary Dynamics of the mtDNA D-Loop Region in Rodents. **Journal of Molecular Evolution** 54 (2): 145-155.
- LIU, H.Z. 2002. The structure and evolution of the mtDNA control region in fish: taking example for Acheilognathinae. **Progress in Natural Science** 12 (3): 266-270.
- LOWE, T.M. & S.R. EDDY. 1997. tRNAscan-SE: a program for improved detection of transfer RNA genes in genomic sequence. Available online at: <http://lowelab.ucsc.edu/tRNAscan-SE> [Accessed: 04/VIII/2011].
- MATHEWS, D.H.; M.E. BURKARD & D.H. TURNER. 1999. **RNAstructure: A computer program for Windows95/NT**. Available online at <http://rna.chem.rochester.edu> [Accessed: 04/VIII/2011].
- NEAR, T.J.; P.A. MEYLAN & H.B. SHAFFER. 2005. Assessing concordance of fossil calibration points in molecular clock studies: an example using turtles. **The American Naturalist** 165: 137-146.
- NILSSON, M.A. 2009. The structure of the Australian and South American marsupial mitochondrial control region. **Mitochondrial DNA** 20 (5-6): 126-38.
- NOONAN, B.P. 2000. Does phylogeny of pelomedusid turtles reflect vicariance due to continental drift? **Journal of Biogeography** 27: 1245-1249.
- NOONAN, B.P. & P.T. CHIPPINDALE. 2006. Vicariant origin of Malagasy reptiles supports Late Cretaceous Antarctic landbridge. **The American Naturalist** 168: 731-741.
- PENG, R.; B. ZENG; X. MENG; B. YUE; Z. ZHANG & F. ZOU. 2007. The complete mitochondrial genome and phylogenetic analysis of the giant panda (*Ailuropoda melanoleuca*). **Gene** 397: 76-83.
- QUINN, T.W. & A.C. WILSON. 1993. Sequence evolution in and around the mitochondrial control region in birds. **Journal of Molecular Evolution** 37: 417-425.
- RAMIREZ, V.; P. SAVOIE & R. MORAIS. 1993. Molecular characterization and evolution of a duck mitochondrial genome. **Journal of Molecular Evolution** 37: 296-310.
- RANDI, E. & V. LUCCHINI. 1998. Organization and Evolution of the Mitochondrial DNA Control Region in the Avian Genus *Alectoris*. **Journal of Molecular Evolution** 47: 449-462.
- RAY, D.A. & L.D. DENSMORE. 2002. The Crocodylian Control Region: General Structure, Conserved Sequences and Evolutionary Implications. **Journal of Experimental Zoology (Molecular Development and Evolution)** 294: 334-345.
- RHODIN, A.G.J.; P.P. VAN DIJK; J.B. IVERSON & H.B. SHAFFER. 2010. Turtles of the world, 2010 update: Annotated checklist of taxonomy and synonymy, Distribution, and Conservation Status, p. 000.85-000.164. *In*: A.G.J. RHODIN; P.C.H. PRITCHARD; P.P. VAN DIJK; R.A. SAUMURE; K.A. BUHLMANN; J.B. IVERSON & R.A. MITTERMEIER (Eds). **Conservation Biology of Freshwater Turtles and Tortoises: A Compilation Project of the IUCN/SSC Tortoise and Freshwater Turtle Specialist Group Chelonian Research Monographs** 5. Checklist v3.2010, available online at: <http://www.iucn-tftsg.org/cbftt>, doi:10.3854/crm.5.000.
- RITCHIE, P. & D. LAMBERT. 2000. A repeat complex in the mitochondrial control region of Adélie penguins from Antarctica. **Genome** 43: 613-618.
- RUOKONEN, M. & L. KVIST. 2002. Structure and evolution of the avian mitochondrial control region. **Molecular Phylogenetics and Evolution** 23: 422-432.
- SACCONE, C.; G. PESOLE & E. SBISA. 1991. The main regulatory region of mammalian mitochondrial DNA: structure-function model and evolutionary pattern. **Journal of Molecular Evolution** 33: 83-91.

- SAITOU, N. & M. NEI. 1987. The neighbor-joining method: A new method for reconstructing phylogenetic trees. **Molecular Biology and Evolution** 4: 406-425.
- SAMBROOK, J. & D. RUSSELL. 2001. **Molecular Cloning: A Laboratory Manual**. New York, Cold Spring Harbour Laboratory Press. 3rd ed.
- SBISÀ, E.; F. TANZARIELLO; A. REYES; G. PESOLE & C. SACCONI. 1997. Mammalian mitochondrial D-loop region structural analysis: identification of new conserved sequences and their functional and evolutionary implications. **Gene** 205: 125-140.
- SERB, J.M.; C.A. PHILLIPS & J.B. IVERSON. 2001. Molecular Phylogeny and Biogeography of *Kinosternon flavescens* Based on Complete Mitochondrial Control Region Sequences. **Molecular Phylogenetics and Evolution** 1 (18): 149-162.
- SUN, K.; J. FENG; L. JIN; Y. LIU; L. SHI & T. JIANG. 2009. Structure, DNA sequence variation and phylogenetic implications of the mitochondrial control region in horse shoe bats. **Mammalian biology** 74: 130-144.
- THOMPSON, J.D.; T.J. GIBSON; F. PLEWNIAK; F. JEANMOUGIN & D.G. HIGGINS. 1997. The ClustalX windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. **Nucleic Acids Research** 24: 4876-4882.
- THOMSON, R.C. & H.B. SHAFFER. 2010. Sparse Supermatrices for Phylogenetic Inference: Taxonomy, Alignment, Rogue Taxa, and the Phylogeny of Living Turtles. **Systematic Biology** 59 (1): 42-58.
- WALBERG, M.W. & D.A. CLAYTON. 1981. Sequence and properties of the human KB cell and mouse L cell D-loop regions of mitochondrial DNA. **Nucleic Acids Research** 9: 5411-5421.
- XIONG, L.; L. NIE; X. LI & X. LIU. 2010. Comparison research and phylogenetic implications of mitochondrial control regions in four soft shelled turtles of Trionychia (Reptilia, Testudinata). **Genes and Genomics** 32: 291-298.
- ZARDOYA, R. & A. MEYER. 1998a. Cloning and characterization of a microsatellite in the mitochondrial control region of the African side-necked turtle, *Pelomedusa subrufa*. **Gene** 216: 149-153.
- ZARDOYA, R. & A. MEYER. 1998b. Complete mitochondrial genome suggests diapsid affinities of turtles. **Proceeding of the National Academy of Sciences of the United States of AME** 95: 14226-14231.
- ZHANG, Y.; L. NIE; Y. HUANG; Y. PU & L. ZHANG. 2009. The Mitochondrial DNA Control Region Comparison Studies of Four Hinged Turtles and its Phylogenetic Significance of the Genus *Cuora Sensu Lato* (Testudinata: Geoemydidae). **Genes & Genomics** 31: 349-359.

Submitted: 05.IV.2011; Accepted: 24.VII.2011.
 Editorial responsibility: Marcio R. Pie

Appendix. Species and associated GenBank accession numbers used in the present study.

Family name	Species name	GenBank accession number
Trionychidae	<i>Trionyx triunguis</i>	NC_012833
	<i>Apalone ferox</i>	FJ890514
	<i>Palea steindachneri</i>	FJ541030
	<i>Dogania subplana</i>	AF366350
	<i>Lissemys punctata</i>	NC_012414
	<i>Pelodiscus sinensis</i>	AY687385
	<i>Carettochelys insculpta</i>	FJ862792
Carettochelyidae	<i>Carettochelys insculpta</i>	FJ862792
Pelomedousidae	<i>Pelomedusa subrufa</i>	AF039066
Podocnemididae	<i>Podocnemis unifilis</i>	this study
	<i>Podocnemis expansa</i>	AY572983
	<i>Chelus fimbriata</i>	this study
Chelidae	<i>Chelodina rugosa</i>	this study
	<i>Mauremys reevesi</i>	AY676201
Geoemydidae	<i>Cuora aurocapitata</i>	AY874540
	<i>Cuora flavomarginata</i>	EU708434
	<i>Mauremys mutica</i>	DQ453753
	<i>Cuora mouhotii</i>	DQ659152
	<i>Sacalia quadriocellata</i>	EF088646
	<i>Cuora galbinifrons</i>	NC_014102

Continue

Appendix. Continued.

Family name	Species name	GenBank accession number
Chelydridae	<i>Chelydra serpentine</i>	NC_011198
	<i>Macrochelys temmincki</i>	EF071948
Emydidae	<i>Chrysemys picta</i>	AF069423
	<i>Trachemys scripta</i>	FJ392294
Cheloniidae	<i>Chelonia mydas</i>	AB012104
	<i>Eretmochelys imbricata</i>	NC_012398
	<i>Lepidochelys olivacea</i>	AM258984
Testudinidae	<i>Testudo kleinmanni</i>	DQ080048
	<i>Stigmochelys pardalis</i>	DQ080041
	<i>Malacochersus tornieri</i>	DQ080042
	<i>Manouria emys</i>	DQ080040
	<i>Indotestudo elongata</i>	DQ080043
	<i>Testudo horsfieldii</i>	DQ080045
	<i>Testudo graeca</i>	DQ080049
	<i>Indotestudo forstenii</i>	DQ080044
	<i>Testudo marginata</i>	DQ080047
	Platysternidae	<i>Platysternon megacephalum</i>
Phasianidae	<i>Gallus gallus</i>	AY235571

Note: The taxonomical classification provided by Genbank entries was used for the species and the classification of RHODIN *et al.* (2010) was used for the following four species: *Macrolemys temminckii* (Troost in Harlan, 1835) was replaced by *Macrochelys temminckii* (Troost, 1835); *Chinemys reevesii* (Gray, 1831) should be *Mauremys reevesii* (Gray, 1831); *Geochelone pardalis* (Bell, 1828) should be *Stigmochelys pardalis*; and *Pyxidea mouhotii* (Gray, 1864) should be *Cuora mouhotii* (Gray, 1862).