

Vitamin K: content in food consumed in São Paulo, Brazil

Simone Aparecida dos Santos Conceição Faria*, Vanilda Aparecida Soares de Arruda, Elias da Silva Araújo, Marilene De Vuono Camargo Penteado

Department of Food and Experimental Nutrition, Faculty of Pharmaceutical Sciences, University of São Paulo

Recent research on Vitamin K has shown its importance in maintaining vascular and bone health. Brazilian food composition tables do not show phylloquinone content in national foods. These data are needed to obtain more reliable results in nutritional status assessment studies of individuals in relation to this vitamin as studies have shown a geographical influence in food phylloquinone content. This study aims to determine phylloquinone (Vitamin K₁) levels in its most important source: dark green leaved vegetables. Several varieties of vegetables were purchased directly from CEAGESP (General Warehouse Company of São Paulo) at different times. Phylloquinone was extracted using organic solvents and quantified by High Performance Liquid Chromatography - HPLC. Results show the concentrations of phylloquinone in commonly consumed foodstuffs. In general, results showed variations with data from literature on the amount of Vitamin K in the plants analysed.

Keywords: Phylloquinone/determination/leaf vegetables. Food composition. High Performance Liquid Chromatography. Vitamin K/study/Brazil. Cooking/Loss.

INTRODUCTION

Vitamin K is a fat-soluble vitamin that was originally identified as an essential factor in blood synthesis of coagulation proteins (factors II, VII, IX, and X) (Booth, Davidson, Sadowski, 1994). Several recent studies have demonstrated an important role in maintaining vascular and bone health through the Vitamin K-dependent proteins (Claussen *et al.*, 2015; Viegas *et al.*, 2014; Yamaguchi, 2014; Beulens *et al.*, 2013; Booth, 2012). The natural forms of Vitamin K are: phylloquinone and menaquinone, the latter being synthesized by bacteria in the intestinal flora and known as “menaquinone-n” or Vitamin K₂ (Booth, 2012; Penteado, 2003). Vitamin K₁ (phylloquinone) is found in foods of both animal and vegetable origin, but is the only analogue of the vitamin present in plants, being predominantly found in vegetables and vegetable oils (Booth, 2012; Damon *et al.*, 2005). Although it is generally assumed that Vitamin K deficiency is probably more common than previously believed, there are several questions about recommended adequate intake (AI) levels and the true dietary requirement of

Vitamin K is unknown (Booth, 2012; Booth *et al.*, 2003). The precise physiological functions of newly discovered Vitamin K proteins are not known except for the function of osteocalcin in bone growth regulation (Vermeer *et al.*, 1996). It is also unclear which markers are best for evaluating Vitamin K status (Booth *et al.*, 2003). With the discovery of new roles for different forms of Vitamin K, it is important to quantify their content in a variety of foods (Elder *et al.*, 2006). In place of biological and chemical methods, the introduction of High Performance Liquid Chromatography (HPLC) has facilitated routine analysis of Vitamin K in foods (Ogles, Cagindi, 2007; Pérez-Ruiz *et al.*, 2007; Jakob, Elmadfa, 2000; 1996; Ferland, Sadowski, 1992). Recently, methods have been developed for determining Vitamin K levels in foods using High Performance Liquid Chromatography-Tandem Mass Spectrometry HPLC-MS/MS (Claussen *et al.*, 2015) and Gas Chromatography-Mass Spectrometry GC-MS (Jang, Moon, Shibamoto, 2015). Although, these methods offer high sensitivity and accuracy, they are very expensive for routine assays. However, separation and detection by HPLC with electrochemical and fluorescence detectors after post-column reduction is relatively convenient and stable, and provides enough sensitivity and selectivity for analysing phylloquinone in plant products (Pérez-Ruiz *et al.*, 2007; Jakob, Elmadfa, 2000). Brazil still has no tables

*Correspondence: S.A.S.C. Faria. Departamento de alimentos de nutrição Experimental. Faculdade de Ciências Farmacêuticas. Universidade de São Paulo. Av. Prof. Lineu Prestes, 580, Bl. 14 - 05508-900 - São Paulo, SP - Brasil. E-mail: sinett@usp.br

of phylloquinone levels for nationally produced foods (Dôres, Paiva, Campana, 2001). There is therefore a need to analyse Vitamin K levels in foods grown in Brazilian soil to assist researchers in the nutrition area, as studies have shown that geography can influence phylloquinone levels in analysed foods (Otles, Cagindi, 2007; Ferland, Sadawski, 1992). This study therefore aims to determine phylloquinone levels in its most important source: leafy green vegetables by HPLC fluorescence using post-column chemical reduction for Vitamin K derivatives.

MATERIAL AND METHODS

Samples

Samples were acquired directly from CEAGESP (São Paulo General Warehouse Company), the largest supply centre in Latin America (Brasil, 2010). Twenty different leafy vegetables were collected between February 2011 and October 2012 and a “pool” of samples was set up with approximately 10 kg per collection for each vegetable type (Faria, 2013). The samples were analysed at the Food Analysis Laboratory of the Department of Food and Experimental Nutrition, Pharmaceutical Sciences Faculty, University of São Paulo. The day after collection sample pre-analysis preparation began following procedures described by Kawashima and Soares (2003). For vegetables that underwent a cooking process, the procedures were as established by UNICAMP (2011) according to vegetable characteristics: for broadleaf samples, baking time was 10 minutes and for floral samples, 20 minutes. About 300 g of each sample was weighed on an analytical scale. They were then placed in a 4-L stainless steel pan and 1.5 L of water added. Cooking was performed on a conventional stove, always using the same stove with a medium flame. Cooking time began when the water started boiling.

Analytical reference standards

Phylloquinone analytical standard and Vitamin K₂, used as internal standards (IS), were purchased from the Sigma Chemical Co. (St. Louis, MO, USA). A stock solution containing 20 µg/mL total phylloquinone in hexane was prepared. A similar stock solution containing 20 µg/mL total IS in hexane was prepared. All standard solutions were stored refrigerated (-20 °C) in amber Eppendorf tubes. Working standard solutions and subsequent dilutions for construction of the standard curve were prepared according to Gao and Ackman (1995). The concentration range was 0.052–2.08 µg mL⁻¹ (n=10 points)

to determine the limit of detection (LOD) and limit of quantitation (LOQ) for the method. The whole procedure was protected from light.

The reagents used to prepare mobile phases and solutions for the standard curves were HPLC grades from Merck (Merck Millipore Corp., Germany). Ultra-pure water was obtained through a Milli-Q Plus Direct – Q3 system from Millipore. All standard and sample solutions were filtered through a 13 mm diameter 0.45 µm pore size Millipore filter unit prior to injection into the chromatographic system. The mobile phases were vacuum-filtered through 0.45 µm pore size nylon Millipore filter (FHLC04700) and degassed by ultrasonic bath for the required period before injection into the chromatographic system.

HPLC analyses

The extraction and concentration of phylloquinone from food matrices used organic solvents with subsequent quantification by HPLC. Analysis was based on the methodology by Jakob and Elmadfa (2000; 1996) with an adjustment in the extract purification stage; after 5 minutes centrifuging at 3000 rpm and 4 °C, the supernatant was collected, filtered through a membrane, and transferred to a 50 mL evaporation flask and evaporated at 40 °C for 15 minutes using a rotary evaporator and vacuum system. The residue was dissolved in 200 µL of mobile phase and 50 µL was injected. A Shimadzu HPLC included a CBM-20A system (SCL-10AVP), LC-Solution Software, a SIL-20A auto sampler, an LC-20AT isocratic pump, and an RF-10AXL fluorescence detector. The separation was in reverse phase with a LiChrospher RP-18 5 µm endcapped LiChroCART 250-4.6 column, with a pre-column from Merck and a mobile phase consisting of dichloromethane/methanol (10:90 v/v) with the addition of 5ml of methanol solution with zinc chloride (1.37 g), sodium acetate (0.41 g) and acetic acid (0.30 g) per litre of mobile phase and was pumped at a flow rate 1.00 mL min⁻¹ with isocratic elution. The post-column reduction (20 x 4.0 mm id) was filled manually with zinc dust p.a. grade from Merck with particles <45 µ, kept in a furnace (Shimadzu-CTO-6A) at 40 °C, with fluorescence detector excitation 243 nm and emission 430 nm.

Validation method

A sequence of chromatographic optimization and validation runs were performed using a set of calibration samples assayed in triplicate and quality control samples at three levels in triplicate; these were carried out on

seven separate occasions. Procedures were performed to validate the method for: Linearity, Limit of Detection and Quantification, Precision, Accuracy, Selectivity and Robustness (Lanças, 2004; Ribani *et al.*, 2004; Youden, Steiner, 1975).

Statistical analysis

The experiments were conducted in a completely randomized way and all data were tested for normal distribution (Shapiro-Wilk's test) and homogeneity of variance (Levene and Brown-Forsythe's tests). For fresh vegetables analysed *in nature*, analyses of variance (ANOVA) were performed with a random factor followed by Bonferroni multiple comparisons (Neter, *et al.*, 1996) to compare Vitamin K levels at two different times per sample. Vitamin K greenery from CEAGESP and American values were compared by using the Student t test against a fixed value (Kirkwood, Sterne, 2006). The results were expressed as means of results \pm standard deviation. All statistical analyses were performed using the STATISTICA 8.0 program with a significance level of 5% ($p < 0.05$).

RESULTS AND DISCUSSION

Vitamin K can be separated by normal and reverse phase chromatography. In this work, the reversed-phase C₁₈ system was used as the chromatography column showed good resolution on simultaneous separation of phylloquinone peaks. These conditions were also used by Otles and Cangindi (2007). The use of post-column reduction proved important for detection as no natural Vitamin K homologues emit fluorescence. Sample fluorescence was induced using a chemical reduction method employed by some authors to quantify Vitamin K in green vegetables (Pérez-Ruiz *et al.*, 2007; Pentead, 2003; Kamao *et al.*, 2005; Maccreehan, Schönberger, 1995). The method was stated to be specific and linear in the range of 0.025-2.8 $\mu\text{g/mL}$ ($r = 0.9997$). Precision was demonstrated by calibration curve standard deviation ($SD = 0.0088$). The method is robust relative to changes in flow rate, column, and temperature. Detection and quantitation limits were 0.0051 and 0.0157 $\mu\text{g/mL}$ respectively.

Selected method precision was tested and the accuracy of the selected methods was tested and phylloquinone recovery calculated by adding three standard levels (25, 50 and 100%). Standard recovery corresponded to 80% based on sample concentration.

Leafy vegetables occupy an important place in well

balanced diets with the most common leafy vegetables consumed raw in Southeast Brazil being lettuce, arugula, and watercress, with cabbage, cauliflower, endive, and spinach used as cooked vegetables (Kawashima, Soares, 2003). The lack of information on the composition of Brazilian foods highlights the need for more data on the composition of leafy vegetables as these are present in the main meals of all social classes (Kawashima, Soares, 2003; Dôres, Paiva, Campana, 2001). Brazilian researchers explain that the national bibliography is quite sparse regarding Vitamin K; for this reason, it is necessary to carry out further studies and improve composition tables, to increase knowledge of Vitamin K levels in Brazilian foods (Klack, Carvalho, 2006; Dôres, Paiva, Campana, 2001).

Table I shows the means of triplicate data for both *in nature* and processed samples collected at CEAGESP at different times of the year in 2011 and 2012. Results show that higher levels of phylloquinone were found in raw leafy vegetables, which was present in those with darker green leaves and/or darker flowers, such as parsley, spinach, broccoli, kale, chicory catalogna, and watercress. These results agree with literature (Beulens *et al.*, 2013; Booth, 2012) which mentions the group of dark green vegetables as foods rich in phylloquinone and state that this compound should be associated with tissues that perform photosynthesis, causing dark green vegetables with high chlorophyll concentrations to contain the highest levels of the vitamin. In contrast, lower levels were obtained in samples of cauliflower, lettuce, and chard; these vegetables are lighter green. Data from *in nature* vegetables were compared for different seasons of the year. These results generally showed statistical differences in Vitamin K levels for all samples. There was significant temporal variation in maximum and minimum Vitamin K levels except for samples of chicory catalogna 279-280 $\mu\text{g}/100\text{ g}$ and chard 150-153 $\mu\text{g}/100\text{ g}$. This variation factor was particularly accentuated in broccoli which presented high and low mean vitamin values of 374 μg and 192 $\mu\text{g}/100\text{ g}$, in broccoli raab of 369 and 165 $\mu\text{g}/100\text{ g}$, and in spinach of 383 and 113 $\mu\text{g}/100\text{ g}$. These results showed that for most *in nature* samples, the spring/summer seasons were favourable for achieving higher Vitamin K levels whereas the autumn/winter periods produced the lowest concentrations phylloquinone (Faria, 2013). Booth (2012) likewise reports that the amount of phylloquinone increases more in summer than winter months. Some possible causes for the variations in phylloquinone levels of samples grown in Southeast Brazil are cultivating, planting location, agricultural practices, such as irrigation and possibly temperature.

For the latter, drastic temperature changes during the year cause prolonged periods of drought and irregular rainfall – conditions which have also been cited by Kawashima and Soares (2003) to explain the variations in mineral content found in samples of fresh leafy vegetables such as lettuce, kale, spinach, cabbage and chicory purchased from markets in Southeast Brazil.

Regarding processed samples, the aim was to reproduce the domestic practice of heat treatment for products such as leafy and floral vegetables commonly used in Brazilian cuisine (UNICAMP, 2011). Table I shows the influence of domestic processing on total phyloquinone content for the evaluated vegetables. In most cases the cooking process produced losses in

TABLE I - Concentration of phyloquinone in fresh and processed vegetables

Sample Identification	Mean ($\mu\text{g}/100\text{ g}$)	Range
Arugula (<i>Eruca sativa</i> L.)	289	228 - 319
Basil (<i>Ocimum basilicum</i> L.)	335	283 - 386
Raw Broccoli (<i>Brassica oleracea</i> L. var. <i>italica</i>)	279	192 - 374
Cooked Broccoli	267	131 - 404
Raw Broccoli Raab (<i>Brassica oleracea</i> L. var. <i>italica</i> Plenck)	260	165 - 369
Cooked Broccoli Raab	228	205 - 250
Raw Cabbage (<i>Brassica oleracea</i> L. var. <i>capitata</i>)	328	319 - 353
Cooked Cabbage	244	197 - 290
Raw Chicory Catalogna (<i>Cichorium Intybus</i> L.)	280	279 - 280
Cooked Chicory Catalogna	226	171 - 281
Raw Cauliflower (<i>Brassica oleracea</i> var. <i>botrytis</i> L.)	37	23 - 46
Cooked Cauliflower	56	47- 66
Raw Celery (<i>Apium graveolens</i> L. var. <i>rapaceum</i>)	295	268 - 322
Cooked Celery	309	293 - 324
Raw Swiss chard (<i>Beta orientalis</i> L.)	152	150 - 153
Cooked Swiss chard	122	118 - 127
Raw Endive (<i>Cichorium endivia</i> L.)	176	172 - 205
Cooked Endive	209	175 - 243
Chives (<i>Allium fistulosum</i>)	160	105 - 192
Raw Kale (<i>Brassica oleracea</i> var. <i>acephala</i>)	280	212 - 362
Cooked Kale	269	221 - 326
Coriander (<i>Coriandrum sativum</i> L.)	258	215 - 300
Raw Chinese Cabbage (<i>Brassica rapa pekinensis</i> (Lour.) Hanelt)	153	118 - 208
Cooked Chinese Cabbage	189	172 - 243
Raw Escarole (<i>Cichorium endivia</i>)	174	110 - 230
Cooked Escarole	155	119 - 234
Iceberg Lettuce (<i>Lactuca sativa capitata</i>)	113	86 - 147
Leafy Lettuce (<i>Lactuca sativa</i> L.)	135	113 - 158
Mint (<i>Mentha longifolia</i>)	364	380 - 348
Parsley (<i>Petroselinum hortense</i>)	491	357 - 558
Raw Spinach (<i>Spinacea oleracea</i> L.)	375	316 - 423
Cooked Spinach	262	113 - 383
Watercress (<i>Nasturtium officinale</i>)	301	264 - 347

Analyses carried out in triplicate and results expressed as mean \pm SD = Standard Deviation. Samples that have undergone the cooking process followed the methodology of TACO – Brazilian Table of Food Composition - UNICAMP.

phylloquinone content compared to the raw values with some samples such as spinach, broccoli and cabbage showing losses of up to 30% indicating consumption in the raw form would provide better utilization of the vitamin. Kale samples lost an average of 1.15% of phylloquinone, indicating practically no loss was suffered from the effects of cooking. Chicory, cauliflower and celery samples showed higher levels of the vitamin in processed samples than their raw counterparts; we can therefore conclude that the cooking process favoured higher vitamin extraction due to disrupting the cell membranes of the food matrices by heating leaving the compound of interest more exposed thus facilitating its extraction. This factor was also reported by Gliszczynska-Swiglo *et al.* (2006) in a study evaluating cooking by boiling vegetable samples. The differentiated behaviour observed in relation to vitamin content of samples which have experienced the cooking process may have been caused by the loss or amount of water incorporated by the samples through the process. The effects of cooking by steam and microwave on phytochemical substances (polyphenols, carotenoids and ascorbic acid) in foods such as fresh fruits and vegetables have been extensively studied with both positive and negative effects reported depending on differences in process conditions, morphology, and nutritional characteristics of different plant species (Mazzeo *et al.*, 2011; Pellegrini *et al.*, 2010; Miglio *et al.*, 2008; Gliszczynska-Swiglo *et al.*, 2006; Turkmen, Sari, Velioglu, 2005). Mazzeo *et al.* (2011) evaluated the effects of steaming and boiling spinach and cauliflower samples and reported that both treatments showed significant changes in the nutritional composition both vegetables. Pellegrini *et al.* (2010) demonstrated that different thermal household processes increased the bioavailability of polyphenols and carotenoids, highlighting the positive role of cooking on the nutritional qualities of vegetables. These improvements offered by heat treatment in vegetables were also reported by Damon *et al.* (2005) and Miglio *et al.* (2008). Pellegrini *et al.* (2010) reported that the absorption efficiency of phylloquinone can vary substantially and be less efficient when the food is eaten raw because the vitamin is intimately linked to the thylakoid and chloroplast membranes in the cells of these plants. Absorption is most efficient when foods are ingested processed because breaking the food matrix promotes increased bioavailability of many phytochemicals thereby improving the nutritional quality of the plants. The authors also emphasize that existing data on different methods of thermal vegetable processing are fragmented and incomplete making it difficult to make comparisons between studies.

Evaluation of phylloquinone levels in samples collected at CEAGESP compared to those of USDA. Table II presents the mean of triplicate samples collected at CEAGESP during different seasons of 2011 and 2012 compared to levels for the same vegetables in the USDA database.

TABLE II – Comparison between Brazilian and American phylloquinone concentration data in fresh vegetables ($\mu\text{g}/100\text{ g}$)

Samples <i>in nature</i>	CEAGESP* Mean	USDA**	p
Iceberg lettuce	113	24	<0.001
Leaf lettuce	135	126	0.315
Parsley	491	1640	<0.001
Arugula	289	109	<0.001
Chard	152	231	<0.001
Cabbage	328	76	<0.001
Spinach	375	483	<0.001
Kale	280	437	<0.001
Cauliflower	37	15	<0.001
Broccoli	279	102	<0.001
Broccoli Raab	260	225	<0.001
Chives	160	213	<0.001
Coriander	258	310	<0.001
Celery	295	29	<0.001
chicory	176	298	<0.001
Watercress	301	250	<0.001

* Triplicate results from GEACESP samples analysed at different times of the year in 2011 and 2012. **USDA - United States Department of Agriculture. p = Descriptive level of statistical test.

Table II shows that only raw leafy lettuce has statistically the same amount of phylloquinone ($p>0.05$) as online data from the United States Department of Agriculture - USDA. Parsley, chard, spinach, kale, chives, coriander, and chicory samples sourced from CEAGESP have lower vitamin content, and iceberg lettuce, arugula, cabbage, cauliflower, broccoli, celery and watercress have higher vitamin content than online USDA data ($p<0.05$). Table III shows data from other studies and highlights significant variations in phylloquinone content for the same vegetables. As described here and by other authors, differences in phylloquinone levels for certain vegetables according to geographic location suggest that levels of this vitamin may be influenced by growing conditions, soil, climate, and agricultural tracts (Booth, 2012; Klack, Carvalho, 2006).

TABLE III – Comparison between different studies for phylloquinone content in vegetables ($\mu\text{g}/100\text{ g}$)

	1	2	3	4	5	This study
Broccoli	147 - 205	178	110	113 - 180	179	308
Cauliflower	27 - 39	-	20	20	31	37
Leafy lettuce	120 - 140	519 - 1180	160	122	129	147
Iceberg lettuce	-	-	40	31	-	126
Spinach	294 - 433	1001- 1439	270	300 - 380	380	404
Cabbage	174 - 204	449 - 719	60	145	339	336
Parsley	530 - 560	-	360	-	548	558
Watercress	-	-	-	-	315	320
Kale	-	621 - 1657	-	440	-	313

1) Bolton-Smith *et al.*, 2000 2) Ferland, Sadowski, 1992; 3) Koivu *et al.*, 1997; 4) Booth, 2012 and Booth *et al.*, 1994; 5) Shearer *et al.*, 1996

Table III compares phylloquinone content in vegetables from this study with other data from literature.

Knowledge on Vitamin K in food is very important in studies which evaluate the relationship between dietary vitamin intake and nutritional status of the individual. Significant variations can be seen in consumption levels for this vitamin in different studies: Booth *et al.* (2004) analysing Vitamin K intake and bone mineral density in men and women, observed a consumption of 171 $\mu\text{g}/\text{day}$ for women and 153 $\mu\text{g}/\text{day}$ for men. However, average phylloquinone consumption in The Rotterdam Study was 244 $\mu\text{g}/\text{day}$ for women and 257 $\mu\text{g}/\text{day}$ for men (Geleijnse *et al.*, 2004). The discrepancies between Vitamin K levels in different food composition tables may reflect the results of these studies on the intake of this vitamin. Research has shown that Vitamin K has been inadequately studied; one of the biggest issues is related to the food composition tables presenting large variations in Vitamin K levels for the same food. Researchers emphasize that geographical differences in food composition data should be considered when applying database composition values for regional surveys (Booth, 2012; Souza, Rodrigues, Penteadó, 2012; Klack, Carvalho, 2006; Dôres, Paiva, Campana, 2001). For this reason, further studies are needed to improve composition tables and increase knowledge on Vitamin K levels in Brazilian foods. In summary, the results of this study show variations in phylloquinone content between analysed samples and data described in international literature for the same vegetables; this can be explained by the influence of geographical differences. The validation procedure for the analysis method used in this study was effective and can be used in determining phylloquinone levels in leafy vegetable samples. The processing of leafy vegetables by cooking caused changes in phylloquinone levels compared to fresh samples. Most

vegetables obtained in spring showed the highest levels of phylloquinone.

ACKNOWLEDGEMENTS

The authors would like to thank São Paulo Teaching and Research Foundation Support - CNPq (the Brazilian National Council Scientific and Technological Development) for their financial support.

REFERENCES

- BEULENS, J.W.J.; BOOTH, S.L.; VAN DEN HEUVEL, E.G.H.M.; STOECKLIN, E.; BAKA, A.; VERMEER, C. The role of menaquinones (Vitamin K₂) in human health. *Brit. J. Nutr.*, v.110, n.8, p.1-12, 2013.
- BOLTON-SMITH, C.; PRICE, R.J.G.; FENTON, S.T.; HARRINGTON, D.J.; SHEARER, M.J. Compilation of provisional UK database for the phylloquinone (Vitamin K₁) content of foods. *Brit. J. Nutr.*, v.83, n.4, p.389-399, 2000.
- BOOTH, S.L. Vitamin K: food composition and dietary intakes. *J. Food Nutr.*, v.56, p. 5505, 2012.
- BOOTH, S.L.; GOLLY, I.; SACHECK, J.M.; ROUBENOFF, R.; DALLAL, E.G.; HAMADA, K.; BLUMBERG, J.B. Effect of vitamin E supplementation on Vitamin K status in adults with normal coagulation status. *Am. J. Clin. Nutr.*, v.80, n.1, p.143-148. 2004.
- BOOTH, S.L.; MARTINI, L.; PETERSON, J.W.; SALTZMAN, E.; DALLAL, G.E.; WOOD, R.J. Dietary phylloquinone depletion and repletion in older women. *J. Food Nutr.*, v.133, n.8, p.2565-2569, 2003.

- BOOTH, S.L.; DAVIDSON, K.W.; SADOWSKI, J.A. Evaluation of an HPLC method for the determination of phyloquinone (Vitamin K₁) in various food matrices. *J. Agric. Food Chem.*, v.42, n.2, p.295-300, 1994.
- BRASIL. Ministério da Agricultura, Pecuária e Abastecimento. PROHORT - *Programa Brasileiro de Modernização do Mercado Hortigranjeiro*. Disponível em: <<http://www.ceasa.gov.br/index.php?pag=10>>. Access: May 19, 2010.
- CLAUSSEN, F.A.; TAYLOR, M.L.; BREEZE, M.L.; LIU, K. Measurement of Vitamin K₁ in commercial canola cultivars from growing locations in north and south America using High-Performance Liquid Chromatography-Tandem Mass Spectrometry. *J. Agric. Food Chem.*, v.63, p.1076-1081, 2015.
- DAMON, M.; ZHANG, N.Z.; HAYTOWITZ, D.B.; BOOTH, S.L. Phyloquinone (Vitamin K₁) content of vegetables. *J. Food Compos. Anal.*, v.18, n.8, p.751-758, 2005.
- DÔRES, S.M.C.; PAIVA, S.A.R.; CAMPANA, A.O. Vitamina K: metabolismo e nutrição. *Rev. Nutr.*, v.14, n.3, p.207-218, 2001.
- ELDER, S.J.; HAYTOWITZ, D.B.; HOWE, J.; PETERSON, J.W.; BOOTH, S.L. Vitamin K contents of meat, dairy, and fast food in the U.S. diet. *J. Agric. Food Chem.*, v.54, n.2, p.463-467, 2006.
- FARIA, S.A.S.C. *Teores de vitamina K em hortaliças consumidas na cidade de São Paulo*. 2013. 113f. Tese (Doutorado em Ciências dos Alimentos) - Faculdade de Ciências Farmacêuticas, Universidade de São Paulo, São Paulo, 2013.
- FERLAND, G.; SADAWSKI, J.A. Vitamin K₁ (phyloquinone) content of green vegetables: Effects of plant Maturation and geographical growth location. *J. Agric. Food Chem.*, v.40, n.10, p.1874-1877, 1992.
- GAO, Z.H.; ACKMAN, R.G. Determination of vitamin K₁ in canola oils by high performance liquid chromatography with menaquinone-4 as an internal standard. *Food Res. Int.*, v.28, n.1, p.61-69, 1995.
- GELEIJNSE, J.M.; VERMEER, C.; GROBBEE, D.E.; SCHURGERS, L.J.; KNAPEN, M.H.; VAN DER MEER, I.M.; HOFMAN, A.; WITTEMAN, J.C. Dietary intake of menaquinone is associated with a reduced risk of coronary heart disease: the Rotterdam Study. *J. Nutr.*, v.134, n.11, p.3100-3105, 2004.
- GLISZCZYNSKA-SWIGLO, A.; CISKA, E.; PAULAK-LEMANSKA, K.; CHMIELEWSKW, J.; BORKOWSKW, T.; TYRAKOWSKA, B. Changes in the content of health-promoting compounds and antioxidant activity of broccoli after domestic processing. *Food Addit. Contam.*, v.23, n.11, p.1088-1098, 2006.
- JAKOB, E.; ELMADFA, I. Rapid and simple HPLC analysis of Vitamin K in food, tissues and blood. *Food Chem.*, v.68, n.2, p.219-221, 2000.
- JAKOB, E.; ELMADFA, I. Application of a simplified HPLC assay for the determination of phyloquinone (Vitamin K₁) in animal and plant food items. *Food Chem.*, v.56, n.1, p.87-91, 1996.
- JANG, H.S.; MOON, J-K.; SHIBAMOTO, T. Analysis and antioxidant activity of extracts from broccoli (*Brassica oleracea* L.) Sprouts. *J. Agric. Food Chem.*, v.63, n.4, p.1169-1174, 2015.
- KAWASHIMA, L.M.; SOARES, L.M.V. Mineral profile of raw and cooked leafy vegetables consumed in southern Brazil. *J. Food Compos. Anal.*, v.16, n.5, p.605-611, 2003.
- KAMAO, M.; SUHARA, Y.; TSUGAWA, N.; OKANO, T. Determination of plasma Vitamin K by high-performance liquid chromatography with fluorescence detection using Vitamin K analogs as internal standards. *J. Chromatogr. B.*, v.816, n.1-2, p.41-48, 2005.
- KIRKWOOD, B.R.; STERNE, J.A.C. *Essential medical statistics*. 2.ed. Malden: Blackwell Science, 2006.
- KLACK, K.; CARVALHO, J.F. Vitamina K: metabolismo, fontes e interação com o anticoagulante varfarina. *Rev. Bras. Reumatol.*, v.46, n.6, p.398-406, 2006.
- KOIVU, T.J.; PIIRONEN, V.I.; HENTTONEN, S.K.; MATTILA, P.H. Determination of phyloquinone in vegetables, fruits, and berries by High-Performance Liquid Chromatography with electrochemical detection. *J. Agric. Food Chem.*, v. 45, n.12, p.4644-4649, 1997.
- LANÇAS, F.M. *Validação de métodos cromatográficos de análise*. 6. ed. São Carlos: Rima, 2004. v.1, 62p.
- MACCREHAN, W.A.; SCHÖNBERGER, E. Determination of Vitamin K₁ in serum using catalytic-reduction liquid chromatography with fluorescence detection. *J. Chromatogr. B.*, v.670, n.2, p.209-217, 1995.

- MAZZEO, T.; N'DRI, D.; CHIVARO, E.; VISCONTI, A.; FOGLIANO, V.; PELLEGRINI, N. Effect of two cooking procedures on phytochemical compounds, total antioxidant capacity and colour of selected frozen vegetables. *Food Chem.*, v.128, n.3, p.627-233, 2011.
- MIGLIO, C.; CHIAVARO, E.; VISCONTI, A.; FOGLIANO, V.; PELLEGRINI, N. Effects of different cooking methods on nutritional and physicochemical characteristics of selected vegetables. *J. Agric. Food Chem.* v.56, n.1, p.139-147, 2008.
- NETER, J.; KUTNER, M.H.; NACHTSHEIM, C.J.; WASSERMAN, W. *Applied linear statistical models*. Boston: WCB, McGraw-Hill, 1996. 1408p. (Irwin Series in Statistics).
- OTLES, S.; CAGINDI, O. Determination of vitamin K₁ content in olive oil, chard and human plasma by RP-HPLC method with UV-Vis detection. *Food Chem.*, v.100, n.3, p.1220-1222, 2007.
- PELLEGRINI, N.; CHIAVARO, E.; GARDANA, C.; MAZZEO, T.; CONTINO, D.; GALLO, M.; RISO, P.; FOGLIANO, V.; PORRINI, M. Effect of different cooking methods on color, phytochemical concentration, and antioxidant capacity of raw and frozen brassica vegetables. *J. Agric. Food Chem.*, v.58, n.7, p.4310-4312, 2010.
- PENTEADO, M.D.V.C. *Vitaminas: aspectos nutricionais, bioquímicos, clínicos e analíticos*. Barueri: Manole, 2003. v.1, 612p.
- PÉREZ-RUIZ, T.; MARTÍNEZ-LOZANO, C.; GARCÍA, M.D.; MATÍN, J. High-performance liquid chromatography-photochemical reduction in aerobic conditions for determination of K vitamins using fluorescence detection. *J. Chromatogr. A.*, v.1141, n.1, p.67-72, 2007.
- RIBANI, M.; BOTTOLI, C.B.G.; COLLINS, C.H.; JARDIM, I.C.S.F.; MELO, L.F.C. Validação de métodos cromatográficos e eletroforéticos. *Quim. Nova*, v.27, n.5, p.771-780, 2004.
- SHEARER, M.J.; BACH, A.; KOHLMEIER, M. Chemistry, nutritional sources, tissue distribution and metabolism of Vitamin K with special reference to bone health. *J. Nutr.*, v.126, n.4, p.1181S-1186S, 1996.
- SOUZA, W.N.; RODRIGUES, M.L.; PENTEADO, M.D.V.C. Ingestão habitual de vitamina K em adultos e idosos. *Rev. Nutr.*, v.25, n.4, p.507-515, 2012.
- TURKMEN, N.; SARI, F.; VELIOGLU, Y.S. The effect of cooking methods on total phenolics and antioxidant activity of selected green vegetables. *Food Chem.* v.93, n.4, p.713-718, 2005.
- UNIVERSIDADE ESTADUAL DE CAMPINAS. UNICAMP. *TACO – Tabela de composição de alimentos*. Núcleo de Estudos e Pesquisas em Alimentação. Campinas: NEPA, 2011. Disponível em: <https://www.unicamp.br/nepa/taco/contar/taco_4_edicao_ampliada_e_revisada>. Acesso em: 05 Fev. 2011.
- VERMEER, C.; GLJSBERS, B.L.MLG.; CRACIUN, A.M.; DOOREN, M.M.C.L.G.; KNAPEN, M.H.J. Effects of Vitamin K on bone mass and bone metabolism. *J. Nutr.*, v.126, Suppl. 4, p.1187S-1191S, 1996.
- VIEGAS, C.S.B., HERFS, M.; RAFAEL, M.S.; ENRIQUEZ, J.L.; TEIXEIRA, A.; LUÍS, I.M.; HOOFD, C.M.R.; JOÃO, A.; MARIA, V.L.; CAVACO, S.; FERREIRA, A.; SERRA, M., THEUWISSEN, E.; VERMEER, C.; SIMES, D.C. Glarich protein is a potential new Vitamin K target in cancer: Evidences for a direct GRP-mineral interaction. *Biomed Res. Int.*, v.14, ID340216, 14p., 2014.
- YAMAGUCHI, M. Osteoporosis treatment with functional food factor: Vitamin K₂. *J. Osteopor. Phys. Act.*, v.2, n.1, p.1-3, 2014.
- YOUNDEN, W. J.; STEINER, E. H. *Statistical manual of Association of Official Analytical Chemists: official methods of analysis of AOAC*. Washington: AOAC, 1975. v.1, p.88.

Received for publication on 06th October 2015

Accepted for publication on 19th December 2016