



An Overview About Ultrastructure of Kranz Anatomy in Cyperaceae (Poales)

Shirley Martins^{1,4}, Silvia Rodrigues Machado² and Vera Lucia Scatena³

Abstract

An overview about ultrastructure of Kranz anatomy in Cyperaceae (Poales) - The ultrastructure of the Kranz tissues in Cyperaceae species with the four types of Kranz anatomy (chlorocyperoid, eleocharoid, fimbristyloid and rhynchosporoid) was studied and compared with species already described in the literature, with the goal verifying the occurrence of patterns for the Kranz types. In addition, chloroplasts and mitochondria were quantified for the first time for all Kranz types. The chloroplast's structure is similar in the mesophyll cells (PCA) in the four Kranz types, but differs in some features in the bundle sheath cells (PCR). In the chlorocyperoid and rhynchosporoid types the bundle sheath cells (PCR) present centrifugal or scattered chloroplasts with convoluted or parallel thylakoids. In the eleocharoid types, the chloroplasts are scattered with parallel thylakoids. The fimbristyloid type presents centrifugal chloroplasts with convoluted and parallel thylakoids. To eleocharoid and fimbristyloid types and to *Cyperus* and *Pycreus* (chlorocyperoid) was observed pattern in the chloroplasts position. In most of the studied species the number of mitochondria in the bundle sheath cells (PCR) is significantly higher than in the mesophyll cells (PCA) and this result diverges from the results found in previous studies.

Key words: C₄ photosynthesis, chloroplast, mitochondria, Cyperaceae, Kranz.

Introduction

Species with C₄ photosynthesis are in general anatomically characterized by the presence of two types of specialized photosynthetic tissues: the photosynthetic carbon assimilation (PCA) tissue, constituted by the mesophyll cells (chlorophyllian parenchyma) arranged radially, and the photosynthetic carbon reduction (PCR) tissue, constituted by the vascular bundle sheath with chloroplasts (Soros & Bruhl 2000; Sage 2004; Sage *et al.* 2011; Sage *et al.* 2012). However, in some species of Amaranthaceae that presents C₄ photosynthesis without Kranz anatomy, in which all photosynthetic process occur within a single chlorenchyma cell (Edwards *et al.* 2004; Edwards & Voznesenskaya 2011)

From an ultrastructural point of view, C₄ plants differ from C₃ ones by the great number

of chloroplasts in the bundle sheath cells (PCR), which are absent or fewer in C₃ plants (Brown *et al.* 1983; Sage *et al.* 2011). The structure of the chloroplasts varies between the two photosynthetic tissues (PCA and PCR) mainly in relation to the thylakoids organization (Brown 1958; Black & Mollenhauer 1971; Besnard *et al.* 2009). In addition, the chloroplasts of the PCR tissue can differ in location and organization of the thylakoids system among species from different families or from the same family (Gutierrez *et al.* 1974; Bruhl & Perry 1995). In general, these variations are related to the biochemical subtype of the decarboxylation enzymes: NADP-ME; NAD-ME and PCK (Gutierrez *et al.* 1974; Hatch *et al.* 1975; Yoshimura *et al.* 2004).

Cyperaceae is one of the families with the highest numbers of species with C₄ photosynthesis

¹ Universidade Estadual do Oeste do Paraná (UNIOESTE), Centro de Ciências Biológicas e da Saúde, R. Universitária, Jardim Universitário, 85819-110, C.P. 000711, Cascavel, PR, Brazil.

² Universidade Estadual Paulista (UNESP), Inst. Biociências, Depto. Botânica, Botucatu, 18618-000, São Paulo, Brazil.

³ Universidade Estadual Paulista (UNESP), Inst. Biociências, Depto. Botânica, Rio Claro, 13506-900, São Paulo, Brazil.

⁴ Author for correspondence: shirley_botany@yahoo.com.br

(Sage *et al.* 2011) and includes four Kranz anatomical types (chlorocyperoid, eleocharoid, fimbristylloid and rhynchosporoid) (Soros & Bruhl 2000). In the family each anatomical type are usually associate with a taxonomic group (chlorocyperoid - tribe Cypereae; eleocharoid – *Eleocharis*; fimbristylloid - tribe Abildgaardieae; rhynchosporoid - *Rhynchospora*), but in *Eleocharis* also occurs species with fimbristylloid type and in *Rhynchospora* the chlorocyperoid type (Soros & Bruhl 2000; Martins & Scatena 2011).

The Kranz types in Cyperaceae differ anatomically in the number and continuity of the vascular bundle sheath and in the presence or not of chloroplasts in the bundle sheath (Soros & Bruhl 2000; Martins & Scatena 2011). The fimbristylloid type possesses three bundle sheaths: the outer with thin-walled cells, reduced lumen, and chloroplasts; the middle one has slightly thick-walled cells, reduced lumen, and no chloroplasts; and the inner one (PCR) present thin-walled cells, ample lumen and chloroplasts, being discontinuous in the major veins. In the chlorocyperoid and eleocharoid types the bundles are surrounded by two sheaths: the outer one with slightly thick-walled cells, reduced lumen and no chloroplasts, and the inner one (PCR) with thin-walled cells, large lumen and chloroplast, differing by the continuity of the inner bundle sheath in the major veins only in the eleocharoid type. The rhynchosporoid type differs from the others by the presence of only one sheath (PCR), constituted by thick-walled cells with chloroplasts (Martins & Scatena 2011).

The origin of the PCA and PCR tissues, besides of the others bundle sheaths were demonstrated by Soros and Dengler (2001) and Martins and Scatena (2011). These studies showed that in Cyperaceae Kranz species the PCA tissues originates from ground meristem and the PCR tissues, in all Kranz types, develops from procambium.

The biochemical subtype of the decarboxylation enzymes (NADP-ME or NAD-ME) also varies in the C₄ species of the family (Ueno *et al.* 1989; Bruhl *et al.* 1987; Bruhl & Perry 1995; Voznesenskaya *et al.* 2005), in which the species of the chlorocyperoid, fimbristylloid (tribe Abildgaardieae) and rhynchosporoid types possess the NADP-ME subtype, and the eleocharoid and fimbristylloid (*Eleocharis*) types contain the NAD-ME subtype (Bruhl & Perry 1995; Murphy *et al.* 2007).

The chloroplast characteristics in groups with C₄ photosynthesis might be of great relevance for taxonomy and systematic purposes (Bruhl & Perry

1995; Jacobs 2001). However, in Cyperaceae it is difficult to establish patterns related to the chloroplast structure because of the small number of studied species, mainly in species C₄ of *Rhynchospora* and *Eleocharis*. The ultrastructural studies with C₄ Cyperaceae conducted in species from the four Kranz types have mainly examined species of tribe Cypereae (chlorocyperoid) and tribe Abildgaardieae (fimbristylloid) (Carolin *et al.* 1977; Ueno *et al.* 1988; Estelita 1992; Bruhl & Perry 1995; Rodrigues & Estelita 2003; Martins *et al.* 2008). To *Rhynchospora*, presenting rhynchosporoid and chlorocyperoid Kranz types, some works described the ultrastructural features to *Rhynchospora rubra* (Lour.) Makino (rhynchosporoid) (Carolin *et al.* 1977; Bruhl *et al.* 1987; Bruhl & Perry 1995) and recently, other Kranz *Rhynchospora* species with both Kranz types were described by Ueno (2013), however, were used dry material and therefore some characteristic were unclear, leading the author detach the importance of new studies using fresh material. To eleocharoid type the ultrastructural analysis were conducted mainly with *Eleocharis vivipara* Link and *Eleocharis baldwinii* (Torr.) Chapm. (Bruhl & Perry 1995; Ueno 1996; Uchino *et al.* 1998; Ueno 2004; Murphy *et al.* 2007).

Quantifying organelles is important for characterizing the cell structure of the different Kranz types, and this process has been performed with several Poaceae species (Brown *et al.* 1983; Yoshimura *et al.* 2004). However, in Cyperaceae this approach has only been performed with individuals of *Eleocharis vivipara* (fimbristylloid) growing in distinct environmental conditions (Ueno 1996). The information about quantifying organelles are considerate important for indicating the biochemical subtype of the decarboxylation enzymes (Bruhl *et al.* 1987; Ueno 1996) by the proportion of mitochondria present in the mesophyll cells (PCA - Photosynthetic Carbon Assimilative tissue) relative to the bundle sheath cells (PCR - Photosynthetic Carbon Reductive tissue). The majority studied in Cyperaceae only indicated this proportion by visual observation (Carolin *et al.* 1977; Bruhl *et al.* 1987; Bruhl & Perry 1995).

The aim of this study was to characterize and to quantify the chloroplasts and mitochondria of Cyperaceae species with different types of Kranz anatomy, using fresh material, and compared with species already described in the literature to identify if there is pattern to the types and if the similarities can reflect the taxonomical proximity.

Material and methods

Eight species of Cyperaceae with different types of Kranz anatomy were studied: *Cyperus ligularis* L. (S. Martins 280, 330), *Kyllinga brevifolia* Rottb. (S. Martins 281, 288) and *Rhynchospora barbata* (Vahl) Kunth (S. Martins 313; V.L. Scatena 321) - chlorocyperoid Kranz anatomy; *Eleocharis minima* Kunth (S. Martins 405, 406) - eleocharoid Kranz anatomy; *Bulbostylis scabra* (J. Presl. & C. Presl.) C.B. Clarke (S. Martins 248, 408) and *Fimbristylis autumnalis* L. (V.L. Scatena 337, 343) - fimbristylid Kranz anatomy; *Rhynchospora globosa* Lindl. (S. Martins 261, 305) and *R. terminalis* Kunth (S. Martins 250, 302) - rhynchosporoid Kranz anatomy. The species were collected in their natural habitats and the voucher materials were deposited at the Herbarium of the Department of Botany, Universidade Estadual Paulista (HRCB).

Samples of the mid-region of at least three fully expanded leaf blade or scapes (*Eleocharis minima*) fully developed were fixed in 2.5% glutaraldehyde solution in 0.1 M phosphate buffer at a pH of 7.3 for 24 h at 5°C, post-fixed with 1% osmium tetroxide in the same buffer for 1 h at 25°C, dehydrated with an acetone series and embedded in Araldite resin. Ultrathin transverse sections were obtained with a Diatome diamond knife and were stained with uranyl acetate and lead citrate (Reynolds 1963). The material was examined with a Philips EM 301 transmission electron microscope (MET). The numbers of chloroplasts and mitochondria per cell were counted for 30–40 cells of the mesophyll cells (PCA) and of the bundle sheath (PCR) in eight vascular bundles of at least three leaves under an electron microscope.

Statistical analyses were performed to verify the significant differences among the number of organelles in the PCA and PCR tissues. A nonparametric test (Wilcoxon/Kruskal-Wallis) was applied using R (R Development Core Team 2010).

Results

Mesophyll cells (PCA)

In all studied species, the mesophyll cells (Fig. 1a - asterisk), in transverse sections, have generally lenticular chloroplasts located near the cell walls (Fig. 1b,c) and with well-developed grana (Gr) (Fig. 1d). Plasmodesmata (Pl) occur between adjacent PCA cells (Fig. 1e) and also between the PCA cells and adjacent bundle sheath cells. The PCA cells in the studied species differ in the number of chloroplasts (Tab. 2) and in to the development of the peripheral reticulum (Pr), which is well-developed in *Eleocharis*

minima (eleocharoid), *Rhynchospora terminalis* (rhynchosporoid) and *R. barbata* (chlorocyperoid) (Fig. 1f) and which is reduced or absent in the other species. The mitochondria (Mi) in the PCA cells are oval with developed cristae (Fig. 1c).

Bundle sheaths (Chlorocyperoid)

The Kranz chlorocyperoid species *Cyperus ligularis* (Fig. 1g), *Kyllinga brevifolia* (Fig. 1h) and *Rhynchospora barbata* (Fig. 1i), possess two sheaths around the vascular bundles: the outer one (Os) lacks chloroplasts, whereas the inner one (Is) has larger chloroplasts (PCR) (Fig. 1g, i). In the inner sheath cells (Is), the chloroplasts are located centrifugally in *C. ligularis* (Fig. 1g), but in *K. brevifolia* (Fig. 1h) and *R. barbata* (Fig. 1i) they are centrifugal or do not present a pattern of localization (Tab. 1). The lamellar system of the chloroplasts is composed of thylakoids parallel or convoluted in *C. ligularis* (Fig. 1j), parallel in *K. brevifolia* (Fig. 1k) and parallel or convoluted, sometimes forming concentric circles, in *R. barbata* (Fig. 1l). The peripheral reticulum is reduced or absent (Fig. 1j, k, l and Tab. 1). Starch grains were observed only in *Kyllinga brevifolia*, which can be numerous, large and generally ellipsoids (Fig. 1k). The mitochondria in the PCR cells are usually oval in transverse view (Fig. 1m) and in *K. brevifolia* and *R. barbata*, they were more numerous in PCR cells than in the PCA cells, while in *C. ligularis*, the PCR and PCA cells had similar numbers of mitochondria (Tab. 2). Plasmodesmata (Pl) are frequent between the anticlinal walls of the inner sheath cells (Fig. 1m), connecting them, and between the inner sheath cells and the outer bundle sheath cells (Fig. 1i).

Bundle sheaths (Eleocharoid)

Eleocharis minima possess Kranz eleocharoid anatomy and have two vascular bundle sheaths: the outer bundle sheath (Os) lacks chloroplasts, whereas the inner bundle sheath (Is) present chloroplasts (PCR) (Fig. 2a). In the inner sheath cells, the chloroplasts do not show a constant location and occupy most of the cell lumen (Fig. 2a). These organelles show parallel thylakoids (Fig. 2b), reduced peripheral reticulum and few starch grains. Additionally, in these cells, the mitochondria (Mi) are oval or elongate in transverse view (Fig. 2b) and were more numerous compared to those in the mesophyll cells (PCA) (Tab. 2). Plasmodesmata (Pl) occur between the anticlinal walls of the inner sheath cells and between the inner sheath cells and the outer bundle sheath cells.

Bundle sheaths (Fimbristylloid)

Bulbostylis scabra and *Fimbristylis autumnalis* have Kranz fimbristylloid anatomy (Fig. 2c), with vascular bundles surrounded by three sheaths: the outer sheath (Os) and the inner sheath (Is) with chloroplasts, while the middle sheath (Ms) lacks chloroplasts (Fig. 2c). In the cells of the outer bundle sheath, the chloroplasts are smaller than those of the PCA cells, but are similar in lenticular shape, well-developed grana (Fig. 2d) and reduced peripheral reticulum. In the inner sheath (PCR), the chloroplasts are mainly centrifugal (Fig. 2c, e), with convoluted thylakoids in *Bulbostylis scabra* and convoluted or parallel thylakoids, sometimes forming concentric circles, in *Fimbristylis autumnalis* (Fig. 2e) and a reduced peripheral reticulum. The mitochondria of the inner sheath cells are oval or circular in transverse view with well developed cristae (Fig. 2e). In these species the mitochondria were more numerous in inner sheath cells (PCR) than in the mesophyll cells (PCA) (Tab. 2). Plasmodesmata (Pl) are frequent among the cells of the different sheaths and between the anticlinal walls of the inner sheath cells. Starch grains were not observed in the PCR cells.

Bundle sheaths (Rhynchosporoid)

The rhynchosporoid species, *Rhynchospora globosa* (Fig. 2f) and *R. terminalis* (Fig. 2g), have

vascular bundles surrounded by a single bundle sheath containing chloroplasts (PCR) (Fig. 2f). In the cells of this sheath, the chloroplasts located in general centrifugally (Fig. 2g) and possess usually parallel thylakoids (Fig. 2h, i), a reduced peripheral reticulum and few starch grains. The mitochondria are oval in transverse view and they were observed in similar numbers as those in the mesophyll cells (PCA) (Tab. 2). Plasmodesmata are frequent between the anticlinal walls of the inner sheath cells (Fig. 2i) and between the inner sheath cells and those of the mesophyll cells.

The characteristics of the organelles in the studied species (Tab. 1) were compared with those species of the literature (Tab. 3) to further the discussion.

Discussion

The Kranz Cyperaceae species of the four anatomical types studied here and described in the literature (Carolin *et al.* 1977; Ueno *et al.* 1988; Bruhl & Perry 1995) have chloroplasts in the mesophyll cells (PCA) with well-developed grana. These same characteristics was observed in the chloroplasts of the outer bundle sheath cells in the Kranz fimbristylloid species study here and in the literature (Bruhl & Perry 1995). Therefore, the mesophyll and outer bundle sheath cells in the

Table 1 – Chloroplast features of the vascular bundle sheath cells (PCR) in the Cyperaceae studied species (A = absent; Fd = few developed; Md = moderately developed)

Taxa studied	Chloroplast (PCR)		
	Thylakoid system	Peripheral Reticulum	Position
Chlorocyperoid			
Cypereae			
<i>Cyperus ligularis</i>	convoluted	A	centrifugal
<i>Kyllinga brevifolia</i>	convoluted/parallel	Fd	centrifugal/scattered
Rhynchosporae			
<i>Rhynchospora barbata</i>	contorted/parallel	Fd	centrifugal/scattered
Eleocharoid			
Scirpeae			
<i>Eleocharis minima</i>	parallel	Md	scattered
Fimbristylloid			
Abildgaardieae			
<i>Bulbostylis scabra</i>	convoluted/parallel	A	centrifugal
<i>Fimbristylis autumnalis</i>	convoluted/parallel	A	centrifugal
Rhynchosporoid			
Rhynchosporae			
<i>Rhynchospora globosa</i>	parallel (contorted)	A	centrifugal/scattered
<i>Rhynchospora terminalis</i>	parallel (contorted)	A	centrifugal/scattered




Figure 1 – a-f. transmission electron micrographs of leaves of Cyperaceae species with Kranz anatomy in transverse sections showing the radiate parenchyma cells (PCA). g-m. vascular bundle sheath cells (PVC). a, d, g, j. *Cyperus ligularis* (chlorocyperoid) – a. general view of the vascular bundle showing the arrangement of radiate parenchyma cells (PCA); d. detail of the grana in the chloroplast; g. general view of the bundle sheaths; j. detail of the chloroplasts in the inner bundle sheath. b, e, h, k, m. *Kyllinga brevifolia* (chlorocyperoid) – b. PCA cells with chloroplasts; e. plasmodesmata between the PCA cells; h. general view of the bundle sheaths; k. detail of the chloroplasts in the inner bundle sheath; m. detail of the inner bundle sheath cells showing the mitochondria and plasmodesmata. c. *Fimbristylis autumnalis* (fimbristylid) – Chloroplasts and mitochondria of the PCA cells. f, i, l. *Rhynchospora barbata* (chlorocyperoid) – f. detail of the peripheral reticulum of the chloroplast; i. general view of the bundle sheaths; l. detail of the chloroplasts in the inner bundle sheath. Symbols: Asterisk, radiate parenchyma cells; Gr, grana; Is, inner bundle sheath; Mi, mitochondria; Os, outer bundle sheath; Pa, radiate parenchyma cells; Pl, plasmodesmata; Rp, peripheral reticulum. Scale bars: (a) = 5 μm; (b), (g-i) = 2 μm; (c) and (e) = 1 μm; (d), (f), (j), (l-m) = 0.5 μm; (k) = 0.2 μm.

fimbristylloid type, that present the same origin (Martins & Scatena 2011), can be interpreted as having similar functions as indicated by Carolin *et al.* (1977).

The variations in location and structure of chloroplasts in the bundle sheath cells, that constitute the PCR tissue, observed in the species studied here (Tab. 1) also was observed in others studied species (Tab. 3), differing in some aspects (Tabs. 1, 3).

Pattern to chloroplast location only can be indicated to eleocharoid and fimbristylloid (tribe Abildgaardieae) types (Tabs. 1, 3) in the Cyperaceae Kranz species. To the others types this character varies (Tabs. 1, 3) and then it cannot be applied in taxonomical context to taxa with these Kranz types. According to Hattersley & Browning (1981), the chloroplast location can be related with the photosynthetic activities and the centrifugal position facilitate the transport of metabolites. Therefore, the centrifugal chloroplasts position in all fimbristylloid species from the Abildgaardieae described can be influenced by the distance between the PCR and PCA, because this Kranz type is the unique type with three bundle sheaths. In the others Kranz types with two (chlorocyperoid and eleocharoid) or one sheaths (rhynchosporoid) is more common the scattered distribution.

Yet as to the fimbristylloid type (tribe Abildgaardieae), thylakoid system organization was described as convoluted or contorted (Tab. 3), but in the species studied here also occur the parallel organization (Tab. 1). The occurrence of convoluted and contorted thylakoids has been related to the increase to the stromal area (Carolin *et al.* 1977; Rodrigues & Estelita 2003).

In the Kranz chlorocyperoid species, that included members of tribe Cypereae (*Cyperus ligularis* and *Kyllinga brevifolia*) and tribe Rhynchosporae (*Rhynchospora barbata*) from present study and literature, it was not possible establish a pattern because of variation in the chloroplast position, development of the peripheral reticulum and the organization of the thylakoid system (Tabs. 1, 3). Nevertheless, within the tribe Cypereae, the Kranz *Cyperus* and *Pycnus* species had chloroplasts in the centrifugal position (Tab. 3), representing a potential taxonomic character and reflecting similarity between the genera, already indicated by Tucker (1994), who grouped all species of these genera within *Cyperus*.

The similarity observed about the chloroplast location and structure in *Eleocharis* species with eleocharoid (Tabs. 1, 3) or fimbristylloid (Tab. 3) types can be interpreted as a pattern to the Kranz species of the genus and can be useful to the taxonomy of this group.

The characteristics of the chloroplasts in *Rhynchospora* species from both Kranz types (chlorocyperoid and rhynchosporoid) are generally similar in position, thylakoid system organization and the development of the peripheral reticulum (Tabs. 1, 3). So, these features can be considerate common to Kranz species of the genus.

The peripheral reticulum constitutes a series of anastomosed tubules located in the stroma periphery of the chloroplast (Laestch 1974) and present taxonomic values to Kranz Cyperaceae species (Ueno *et al.* 1988; Bruhl & Perry 1995). However, in the studied species here and from the literature, the degree of peripheral reticulum development is varied, even in close groups (Tabs. 1, 3), remaining constant only in Kranz *Rhynchospora*.

Some authors indicated that the variations in the chloroplast location (centripetal or centrifugal), organization of thylakoids system (with or without grana developed) and ratios of mitochondria quantity between the PCA and PCR tissues in Poaceae and Cyperaceae species can be related with the biochemical subtypes (NADP-ME or NAD-ME) (Hatch *et al.* 1975; Hattersley & Browning 1981; Bruhl & Perry 1995). In Poaceae, the species with NADP-ME subtype present centrifugal chloroplasts with reduced or absent grana and similar ratios of mitochondria between the PCA and PCR tissues (Hatch *et al.* 1975; Hattersley & Browning 1981). Whereas in NAD-ME species, the chloroplasts are centripetal, with grana developed and major number of mitochondria in the PCR tissues (Hatch *et al.* 1975; Hattersley & Browning 1981).

About the chloroplast location, in the Kranz species of Cyperaceae studied here (Tab. 1) and described in the literature (Tab. 2), they present centrifugal distribution only in the fimbristylloid (tribe Abildgaardieae) and chlorocyperoid (*Cyperus*, *Pycnus*) types. These Kranz types were indicated to present NADP-ME subtype (Ueno *et al.*, 1989; Bruhl & Perry 1995), corroborating the same chloroplast location observed in NADP-ME Poaceae species (Hattersley & Browning 1981). However, other Kranz Cyperaceae species,




Figure 2 – Transmission electron micrographs of leaves of Cyperaceae species with Kranz anatomy in transverse sections showing the vascular bundle sheaths cells. a-b. *Eleocharis minima* (eleocharoid) – a. general view of the bundle sheaths; b. detail of the chloroplasts and mitochondria in the inner bundle sheath. c. *Bulbostylis scabra* (fimbristylloid) – general view of the bundle sheaths. d-e. *Fimbristylis autumnalis* (fimbristylloid) – detail of the chloroplasts in the outer bundle sheath and in the inner bundle sheath, respectively. f. *Rhynchospora globosa* (rhynchosporoid) – general view of the bundle sheaths. g-i. *Rhynchospora terminalis* (rhynchosporoid) – g-h. general view and detail of the chloroplasts in the inner bundle sheath, respectively; i. plasmodesmata between the inner bundle sheath cells. Symbols: Mi, mitochondria; Ms, middle bundle sheath; Os, outer bundle sheath; Pa, radiate parenchyma cells; Ss, single bundle sheath; Is, inner bundle sheath; Pl, plasmodesmata. Scale bars: (f) = 5 μm ; (a), (c), (g) = 2 μm ; (d-e), (h) = 1 μm ; (b) and (i) = 0.5 μm .

Table 2 – Number of chloroplasts and mitochondria in the PCA and PCR tissues of Cyperaceae studied species. Values are given as means \pm SD. Asterisks represent significant difference at $P < 0.05$ between the PCA and PCR cells and ns represent not significant difference. (PCA = Primary Carbon Assimilation tissue; PCR = Photosynthetic Carbon Redution tissue)

Taxa studied	Rariate Parenchyma (PCA)		Bundle sheath (PCR)		Mitochondria (PCR:PCA)
	Chloroplast	Mitochondria	Chloroplast	Mitochondria	
Chlorocyperoid					
Cypereae					
<i>Cyperus ligularis</i>	2.7 \pm 0.9	2.2 \pm 1.6	1.5 \pm 0.6*	2 \pm 0.7 ^{ns}	~1
<i>Kyllinga brevifolia</i>	2.9 \pm 1.1	2.5 \pm 0.8	2.2 \pm 0.8	5.6 \pm 1.7*	>1
Rhynchosporae					
<i>Rhynchospora barbata</i>	3.1 \pm 1.7	1.9 \pm 2.2	2.9 \pm 0.8 ^{ns}	5.5 \pm 3.3*	>1
Eleocharoid					
Scirpeae					
<i>Eleocharis minima</i>	2.5 \pm 1.1	3 \pm 1	2.3 \pm 1.2 ^{ns}	7.7 \pm 5.1*	>1
Fimbristylid					
Abildgaardieae					
<i>Bulbostylis scabra</i>	3.8 \pm 1.2	2.9 \pm 0.8	3.3 \pm 0.9 ^{ns}	5.8 \pm 1.4*	>1
<i>Fimbristylis autumnalis</i>	2.5 \pm 0.9	1.8 \pm 1.2	2.2 \pm 0.5 ^{ns}	2.2 \pm 1.7*	>1
Rhynchosporoid					
Rhynchosporae					
<i>Rhynchospora globosa</i>	3.3 \pm 0.8	1.8 \pm 0.9	1.9 \pm 0.7*	1.7 \pm 0.9 ^{ns}	~1
<i>R. terminalis</i>	4.6 \pm 2.0	3.6 \pm 1.9	3.9 \pm 2.0 ^{ns}	3.7 \pm 2.5 ^{ns}	~1

Table 3 – Chloroplast features of the vascular bundle sheath cells (PCR) and ratio of mitochondria between the PCA and PCR tissues of Cyperaceae species described in previous studies (A = absente; Fd = few developed; Md = moderately developed; PCA = Primary Carbon Assimilation tissue; PCR = Photosynthetic Carbon Redution tissue; Wd = well developed)

Genera Kranz	Chloroplast (PCR)			Mitochondria (PCR:PCA)
	Thylakoid system	Peripheral Reticulum	Position	
Chlorocyperoid				
Cypereae				
<i>Cyperus</i> ^{2,3,5,6,7}	contorted/convoluted	Fd to Wd	centrifugal	~1
<i>Pycreus</i> ^{1,7}	convoluted	Md	centrifugal	~1
<i>Remirea</i> ⁴	parallel	Wd	centripetal	not seen
<i>Rhynchospora</i> ¹⁰	convoluted	Fd	unclear	not seen
Eleocharoid				
Scirpeae				
<i>Eleocharis</i> ^{1,8,9}	parallel	Fd to Md	scattered	>1
Fimbristylid				
Abildgaardieae				
<i>Bulbostylis</i> ⁷	convoluted	Fd	centrifugal	~1
<i>Fimbristylis</i> ^{1,2,3,7}	contorted/convoluted	Fd to Md	centrifugal	~1
Rhynchosporoid				
Rhynchosporae				
<i>Rhynchospora</i> ^{1,7,10}	convoluted/parallel	A to Fd	centrifugal/scattered	~1

¹Bruhl & Perry (1995); ²Carolin *et al.* (1977); ³Estelita-Teixeira & Handro (1987); ⁴Estelita (1993); ⁵Kim *et al.* (1999); ⁶Rodrigues & Estelita (2003); ⁷Ueno *et al.* (1988); ⁸Ueno (1996); ⁹Ueno (2004); ¹⁰Ueno (2013)

such as rhynchosporoid (*Rhynchospora*) and chlorocyperoid (*Kyllinga*) types, also indicated to have NADP-ME (Ueno *et al.* 1989; Bruhl *et al.* 1987), the distribution of the chloroplast in PCR cells can be scattered or without location pattern (Tabs. 1, 3). Only Kranz *Eleocharis* species were indicated to have NAD-ME biochemical subtypes (Bruhl *et al.* 1987; Bruhl & Perry 1995), but different from those Poaceae species with this subtype, the chloroplasts do not present location pattern (Tabs. 1, 3).

In relation to the ratios of mitochondria between the PCA and PCR tissues (Hatch *et al.* 1975; Hattersley & Browning 1981), in *Kyllinga brevifolia* and *Rhynchospora barbata* (chlorocyperoid) and in *Bulbostylis scabra* and *Fimbristylis autumnalis* (fimbristylloid), the number of mitochondria was significantly greater in the PCR cells relative to the PCA cells, even these Kranz types being indicated to have NADP-ME. To *Rhynchospora* species from the both Kranz types (chlorocyperoid and rhynchosporoid) was indicated no significant increase in the number of mitochondria in the PCR cells in relation to the PCA cells (Ueno *et al.* 1988; Bruhl & Perry 1995; Ueno 2013). This indication is in accord with our observations only to rhynchosporoid species. Thus, the centrifugal localization of the chloroplasts in the PCR cells and the similar ratio of mitochondria between PCA and PCR do not represent consistent characteristics for indicating the biochemical subtype NADP-ME in the Cyperaceae different from that described to NADP-ME Poaceae species.

The species of the *Eleocharis* genus studied here (Tab. 1) and described in the literature (Tab. 3) present chloroplast without location pattern (Tab. 1) or scattered (Tab. 3) and have been indicated to possess the NAD-ME biochemical subtype (Bruhl & Perry 1995), differing from the Poaceae species with this subtype that possess centripetal chloroplast (Hattersley & Browning 1981; Yoshimura *et al.* 2004). Thus, the data observed here show that the characteristics of location, thylakoids organization and mitochondria ratios between PCA and PCR tissues that were indicated as useful traits to distinguish NADP-ME and NAD-ME biochemical subtypes to Poaceae, in general, cannot be applied to Cyperaceae.

The dimorphism between PCA and PCR tissues was corroborated in the present study, in which the PCA cells possess chloroplast with well-developed grana and PCR cells show chloroplasts with

grana absent or reduced (parallel thylakoids). The characterization and quantification of chloroplasts and mitochondria amplify the knowledge about the different types of Kranz anatomy in Cyperaceae, showing that Kranz species of the eleocharoid, rhynchosporoid and fimbristylloid types, have patterns in relation to chloroplast position and/or thylakoids organization. In addition, to the genera *Cyperus*, *Eleocharis*, *Pycneus* and *Rhynchospora* some characters can be useful for future studies of their systematics. To *Cyperus* and *Pycneus* the similarities in the chloroplast structure reinforce the proximity of these genera. To *Eleocharis* and *Rhynchospora* species with different Kranz types show similarities in chloroplasts location and organization of thylakoids.

Acknowledgements

We thank the Fundação de Amparo à Pesquisa do Estado de São Paulo - FAPESP for a PhD grant (2008/09380-2) to S. Martins and financial support (2011/18275-0) and the Conselho Nacional de Desenvolvimento Científico e Tecnológico - CNPq for financial support to V. L. Scatena (301692/2010-6 and 471837/2011-3) and to S. R. Machado (2008/301464-1) and the Centre of Microscopy of the Universidade Estadual Paulista (Botucatu).

References

- Besnard, G.A.; Muasya, M.; Russier, F.; Roalson, E.H.; Salamin, N. & Christin, P.A. 2009 Phylogenomics of C₄ photosynthesis in sedge (Cyperaceae): multiple appearances and genetic convergence. *Molecular Biology and Evolution* 26:1909-1919.
- Black, C.C. & Mollenhauer, H.H. 1971. Structure and distribution and other organelles in leaves with various rates of photosynthesis. *Plant Physiology* 47: 15-23.
- Brown, W.V. 1958. Leaf anatomy in grass systematics. *Botanical Gazette* 119: 170-178.
- Brown, R.H.; Bouton, J.H.; Rigsby, L. & Rigler, M. 1983. Photosynthesis of grass species differing in carbon dioxide fixation pathways. VIII. Ultrastructural characteristics of *Panicum* species in the *Laxa* group. *Plant Physiology* 71: 425-431.
- Bruhl, J.J. & Perry, S. 1995. Photosynthetic pathway-related ultrastructure of C₃, C₄ and C₃-C₄ intermediate sedge (Cyperaceae), with special reference to *Eleocharis*. *Australian Journal of Plant Physiology* 22: 521-530.
- Bruhl, J.J.; Stone, N.E. & Hattersley, P.W. 1987. C₄ acid decarboxylation enzymes and anatomy in sedge

- (Cyperaceae): first record of NAD-Malic enzyme species. *Australian Journal of Plant Physiology* 14: 719-728.
- Carolin, R.C.; Jacobs, S.W.L. & Vesk, M. 1977. The ultrastructure of Kranz cells in the family Cyperaceae. *Botanical Gazette* 138: 413-419.
- Edwards, G.E.; Franceschi, V.R. & Voznesenskaya, E.V. 2004. Single-cell C₄ photosynthesis versus the dual-cell (Kranz) paradigm. *Annual Review of Plant Biology* 55:173-96.
- Edwards, G.E. & Voznesenskaya, E.V. 2011. C₄ photosynthesis: Kranz forms and single-cell C₄ in terrestrial plants. *In: Raghavendra, A.S & Sage, R.F. (eds.). C₄ photosynthesis and related CO₂ concentrating mechanisms. Advances in Photosynthesis and Respiration. Vol. 32. Springer Science + Business Media B.V., Dordrecht. Pp 29-61*
- Estelita, M.E.M. 1992. Origin and structure of the Kranz tissues in Cyperaceae. *Boletim de Botânica da Universidade de São Paulo* 13: 41-48.
- Estelita, M.E.M. 1993. Anatomia dos órgãos vegetativos de *Remirea maritima* Aubl. (Cyperaceae). *Naturalia* 18: 123-134.
- Estelita-Teixeira, M.E. & Handro, W. 1987. Kranz pattern in leaf, scape and bract of *Cyperus* and *Fimbristylis* species. *Revista Brasileira de Botânica* 10: 105-111.
- Gutierrez M; Gracen, V.E. & Edwards, G.E. 1974. Biochemical and cytological relationships in C₄ plants. *Planta* 119:279-300.
- Hatch, M.D.; Kagawa, T. & Craig, S. 1975. Subdivision of C₄-pathway species based on differing C₄ acid decarboxylating systems and ultrastructural features. *Australian Journal of Plant Physiology* 2: 111-128.
- Hattersley, P.W. & Browning, A.J. 1981. Occurrence of the suberized lamella in leaves of grasses of different photosynthetic types. In parenchymatous bundle sheat and PCR (Kranz) sheaths. *Protoplasma* 109: 371-401.
- Kim, I.S.; Pak, J.H.; Seo, B.B. & Song, S.D. 1999. Ultrastructure of leaves in C₄ *Cyperus iria* and C₃ *Carex siderosticta*. *Journal of Plant Biology* 42: 213-221.
- Laetsch, W.M. 1974. The C₄ syndrome: a structural analysis. *Annual Review of Plant Physiology* 25: 27-52.
- Martins, S.; Machado, S.R. & Alves, M. 2008. Anatomia e ultra-estrutura de *Cyperus maritimus* Poir. (Cyperaceae): estratégias adaptativas ao ambiente de dunas litorâneas. *Acta Botanica Brasilica* 22: 289-299.
- Martins, S. & Scatena V.L. 2011. Bundle sheath ontogeny in Kranz and non-Kranz species of Cyperaceae (Poales). *Australian Journal of Botany* 59: 554-562.
- Murphy, L.R.; Barroca, J.; Franceschi, V.R.; Lee, R.; Roalson, E.H.; Edwards, G.E. & Ku, M.S.B. 2007. Diversity and plasticity of C₄ photosynthesis in *Eleocharis* (Cyperaceae). *Functional Plant Biology* 34: 571-580.
- R Development Core Team. 2010. R: A language and environment for statistical computing. Vienna: R Foundation for Statistical Computing. Available at <<http://www.R-project.org>>. Access on 27 April 2011.
- Reynolds, E.S. 1963. The use of lead citrate at high pH as an electron-opaque stain in electron microscopy. *The Journal of Cell Biology* 17: 208-212.
- Rodrigues, A.C. & Estelita M.E.M. 2003. Origin and structure of the Kranz tissue in bracts of *Cyperus giganteus* Vahl (Cyperaceae). *Revista Brasileira de Botânica* 26: 445-452.
- Sage, R.F. 2002. C₄ photosynthesis in terrestrial plants does not require Kranz anatomy. *Trends in Plant Science* 7: 283-285.
- Sage, R.F. 2004. The evolution of C₄ photosynthesis. *New Phytologist* 161: 341-370.
- Sage, R.F.; Christin, P.A. & Edwards, E.J. 2011. The C₄ plant lineage of planet, earth. *Journal Experimental of Botany* 62: 3155-3169.
- Sage, R.F.; Sage, T.L. & Kocacinar, F. 2012. Photorespiration and the evolution of C₄ photosynthesis. *Annual Review of Plant Biology* 63:19-47.
- Soros, C.L. & Bruhl, J.J. 2000. Multiple evolutionary origins of C₄ photosynthesis in the Cyperaceae. *In: Wilson, K.L. & Morrison, D.A. (eds.). Monocots: systematics and evolution. CSIRO Publishing, Melbourne. Pp. 629-636.*
- Tucker, G.C. 1994. Revision of the Mexican species of *Cyperus* L. (Cyperaceae). *Systematic Botany Monographs* 43: 1-213.
- Ueno, O. 1996. Structural characterization of photosynthetic cells in an amphibious sedge, *Eleocharis vivipara*, in relation to C₃ and C₄ metabolism. *Planta* 199: 382-393.
- Ueno, O. 2004. Environmental regulation of photosynthetic metabolism in the amphibious sedge *Eleocharis baldwinii* and comparisons with related species. *Plant Cell Environment* 27: 627-639.
- Ueno, O. 2013. Ultrastructure and carbon isotope ratios of leaves in C₄ species of *Rhynchospora* (Cyperaceae) that differ in the location of Kranz cells. *International Journal of Plant Science* 174: 702-709.
- Ueno, O.; Samejima, M. & Koyama, T. 1989. Distribution and evolution of the C₄ syndrome in *Eleocharis*, a sedge group inhabiting wet and aquatic environments, based on culm anatomy and carbon isotope ratios. *Annals of Botany* 64: 425-438.
- Ueno, O.; Takeda, T. & Maeda, E. 1988. Leaf ultrastructure of C₄ species possessing different Kranz anatomical

- types in the Cyperaceae. *The Botanical Magazine of Tokyo* 101: 141-152.
- Voznesenskaya, E.V.; Chuong, S.D.X.; Kiirots, O.; Franceschi, V.R. & Edwards, G.E. 2005. Evidence that C_4 species in genus *Stipagrostis*, family Poaceae, are NADP-malic enzyme subtype with nonclassical type of Kranz anatomy (Stipagrostoid). *Plant Science* 168: 731-739.
- Yoshimura, Y.; Kubota, F. & Ueno, O. 2004. Structural and biochemical bases of photorespiration in C_4 plants: quantification of organelles and glycine decarboxylase. *Planta* 220: 307-317.