

Parity Indices and Two-Line Matrix Representation for Partitions

I.M. CRAVEIRO^{1*}, A. WAGNER² and D. DOMINGUES³

Received on November 6, 2014 / Accepted on October 28, 2015

ABSTRACT. In this work we present a solution for Andrews’s Problem 5 [1] by establishing a bijection between the sets D_n^0 and D_n^1 defined in Fine’s Theorem [8] and the sets of partitions indexed by their lower parity index [1]. We also solve Andrews’s Problem 6, conjectured in [1].

Keywords: partition, parity index, mock theta function.

1 INTRODUCTION

In this section we present one of Fine’s renowned theorems. Its combinatorial proof can be found in [8]. That proof does not use complicated analytic formulae and its popularization occurred because of its elegance and simplicity, being similar to the famous involutive proof by Franklin. Franklin’s involution was used in many occasions to prove several refinements of Euler’s Pentagonal Number Theorem [7].

Theorem 1.1 (Fine). *Let D_n^0 and D_n^1 be the sets of partitions of n into distinct parts, such that the largest part $a(\lambda) = \lambda_1$ is even and odd, respectively. Then*

$$|D_n^0| - |D_n^1| = \begin{cases} 1, & \text{if } n = \frac{k(3k+1)}{2}; \\ -1, & \text{if } n = \frac{k(3k-1)}{2}; \\ 0, & \text{otherwise.} \end{cases}$$

Problems that involve parity are related to classic partition identities such as Euler’s, Rogers’s, Ramanujan’s and Gordon’s identities. The theorem we just stated is an example of a problem that involves the concept of parity in the study of identities related to Partition Theory. Several authors

*Corresponding author: Irene Magalhães Craveiro.

¹Faculdade de Ciências Exatas e Tecnologia – FACET, Universidade Federal de Grande Dourados – UFGD, Caixa Postal 364, 79084-970 Dourados, MS, Brasil. E-mail: irenecraveiro@ufgd.edu.br

²DMA-IMECC, Universidade de Campinas – Unicamp, 13083-859 Campinas, SP, Brasil.
 E-mail: dricawagner@gmail.com

³Centro de Ensino Monteiro de Carvalho, 20240-090 Santa Teresa, Rio de Janeiro, RJ, Brasil.
 E-mail: deborarodrigues@gmail.com

studied questions that involve parity in partitions, [4], [5], [6]. In [1], Andrews described several results on this subject. At the end of the same work he enumerated fifteen problems. Problems 1, 2 and 3 were solved in [10]. Problems 9 and 10 were solved in [12], with a purely analytic solution. In [11] other solutions for problems 9 and 10 are presented, and in the same paper a proof for problem 5 is given, which is based on some results related to generating functions for partitions and q -series. In [9] there is also a solution to problem 5, based on the involutive proof by Franklin, 1881.

In section 2 we present a solution to Problem 5 establishing a bijection between the sets D_n^0 and D_n^1 defined on Fine's Theorem and the sets of partitions indexed by their lower odd parity index [1]. In section 4 we present a solution for Problem 6, conjectured in [1].

We adopt the standard q -series notation: for each integer $n > 0$ define

$$(a)_n = (a; q)_n = \prod_{j=0}^{n-1} (1 - aq^j) \text{ and } (a; q)_0 = 1.$$

2 LOWER INDEX AND FINE'S THEOREM

Definition 2.1. Let λ be a partition $\lambda = \lambda_1 + \lambda_2 + \dots + \lambda_j$, where

$$\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_j > 0.$$

The lower odd parity index of λ , denoted by I_{lo} , is defined as the maximum length of the subsequences of $\{\lambda_1, \lambda_2, \dots, \lambda_j\}$ whose terms alternate in parity beginning with an odd λ_i .

Example. Consider $\lambda = 8 + 7 + 7 + 6 + 5 + 4 + 4 + 2 + 2 + 1$ a partition of 46. Thus, $I_{lo}(\lambda) = 6$.

We will denote by $p_o(r, m, n)$ the number of partitions of n in m distinct parts with $I_{lo}(\lambda)$ equals to r . Besides, $P_o(y, x; q)$ is the generating function for partitions λ of n into m distinct parts with $I_{lo}(\lambda) = r$, that is,

$$P_o(y, x; q) = \sum_{r, m, n \geq 0} p_o(r, m, n) y^r x^m q^n \tag{2.1}$$

The next result, given in [1], provides an explicit formula for $P_o(y, x; q)$.

Theorem 2.1. The generating function for partitions enumerated by $p_o(r, m, n)$ is given by:

$$P_o(y, x; q) = \sum_{n \geq 0} \frac{x^n y^n q^{\frac{n(n+1)}{2}} \left(-\frac{q}{y}\right)_n}{(q^2; q^2)_n}. \tag{2.2}$$

Evaluating (2.1) and (2.2) in $y = -1$ and $x = 1$ we have:

$$P_o(-1, 1; q) = \sum_{r, N \geq 0} p_o(r, m, N) (-1)^r q^N = \sum_{n \geq 0} \frac{(-1)^n q^{\frac{n(n+1)}{2}} (q; q)_n}{(q^2; q^2)_n}. \tag{2.3}$$

Problem 5 in [1] asks for a combinatorial proof for:

$$\sum_{r,m \geq 0} p_o(r, m, N)(-1)^r = \begin{cases} 1, & \text{if } N = n(3n + 1)/2 \\ -1, & \text{if } N = n(3n + 5)/2 + 1 \\ 0, & \text{otherwise} \end{cases}$$

The following lemma is very useful to solve Problem 5.

Lemma 2.1. Consider $\lambda = \lambda_1 + \lambda_2 + \dots + \lambda_m$ a partition of the positive integer N in m distinct parts, with $I_{lo}(\lambda) = r$ and $\lambda_1 > \lambda_2 > \dots > \lambda_m$. Then r is even (odd) if, and only if, λ_1 is even (odd).

Proof. Let $\lambda = \lambda_1 + \lambda_2 + \dots + \lambda_m$ be a partition of the positive integer N in m distinct parts, with $I_{lo}(\lambda) = r$, $\lambda_1 > \lambda_2 > \dots > \lambda_m$ and r even. Hence there exists an increasing subsequence of $\{\lambda_1, \lambda_2, \dots, \lambda_m\}$, whose length is equal to r , $(\lambda_{i+r-1}, \dots, \lambda_{i+1}, \lambda_i)$ with an odd λ_{i+r-1} . Because λ_{i+r-1} is odd and r is even, then λ_i is even. In the same way, we can conclude that if λ_i is even, then r is even, because otherwise λ_{i+r-1} would be even, a contradiction.

If $\lambda_1 = \lambda_i$ then the proof is over. Suppose, then, that $\lambda_1 > \lambda_i$.

For all $i > j$, we have that λ_j is even. If λ_j is odd we have one more term of the sequence that alternates parity with λ_i , which is even, and so the subsequence has length $r + 1$. However $I_{lo}(\lambda) = r$. Therefore λ_j is even. It follows from taking $j = 1$ that λ_1 is even.

Conversely, suppose that λ_1 is even. Suppose by absurd that $I_{lo}(\lambda) = r$ with an odd r , then there exists a subsequence of λ , $(\lambda_{i+r-1}, \dots, \lambda_{i+1}, \lambda_i)$, with an odd λ_{i+r-1} , and r being the maximum with this property. Hence λ_i is odd. We have that $\lambda_1 \neq \lambda_i$, because λ_1 is even. Since λ_i must alternate parity with λ_1 we have that $I_{lo}(\lambda) \geq r + 1$. Therefore r if even. It is similar in the case r is an odd number. □

We will denote the set of partitions of a positive integer N , with distinct parts, whose lower odd parity index is $I_{lo}(\lambda) = r$ by: \mathcal{P}_N^r . When r is even we use the notation \mathcal{P}_N^0 , and when r is odd, the notation \mathcal{P}_N^1 .

It follows from Lemma 2.1 that given a partition $\lambda = \lambda_1 + \lambda_2 + \dots + \lambda_m$ of the positive integer N in m distinct parts with $I_{lo}(\lambda) = r$ and $\lambda_1 > \lambda_2 > \dots > \lambda_m$, then:

$$\lambda \in \mathcal{P}_N^0 \text{ if, and only if, } \lambda \in D_N^0.$$

In the same way, it follows from Lemma 2.1 that

$$\lambda \in \mathcal{P}_N^1 \text{ if, and only if, } \lambda \in D_N^1.$$

This result establishes a 1-1 correspondence between the sets defined on Fine’s Theorem and the set of partitions λ indexed by $I_{lo}(\lambda) = r$. So we can conclude that

$$|D_N^0| = \sum_{r,m \geq 0; r \text{ even}} p_o(r, m, N)(-1)^r \tag{2.4}$$

and

$$-|D_N^1| = \sum_{r,m \geq 0; r \text{ odd}} p_o(r, m, N)(-1)^r. \tag{2.5}$$

It follows from (2.4) and (2.5) that

$$\sum_{r,m \geq 0} p_o(r, m, N)(-1)^r = |D_N^0| - |D_N^1|. \tag{2.6}$$

Therefore it follows from Theorem 1.1 that

$$\begin{aligned} \sum_{r,m \geq 0} p_o(r, m, N)(-1)^r &= |D_N^0| - |D_N^1| \\ &= \begin{cases} 1, & \text{if } N = n(3n + 1)/2 \\ -1, & \text{if } N = n(3n + 5)/2 + 1 \\ 0, & \text{otherwise.} \end{cases} \end{aligned} \tag{2.7}$$

Therefore we have another solution for Andrews’s Problem 5, [1].

Example. Let $N = 11, 12$, m denote the number of distinct parts of a partition λ of N and r be its lower odd parity index.

Table 1: Distinct partitions of 11 and 12.

11	12
10 + 1	11 + 1
9 + 2	10 + 2
8 + 3	9 + 3
7 + 4	8 + 4
6 + 5	7 + 5
8 + 2 + 1	9 + 2 + 1
7 + 3 + 1	8 + 3 + 1
6 + 4 + 1	7 + 4 + 1
6 + 3 + 2	7 + 3 + 2
5 + 4 + 2	6 + 5 + 1
5 + 3 + 2 + 1	6 + 4 + 2
	5 + 4 + 3
	6 + 3 + 2 + 1
	5 + 4 + 2 + 1

It follows from equation (2.7)

$$\sum_{r,m \geq 0} p_o(r, m, 11)(-1)^r = 0$$

and

$$\sum_{r,m \geq 0} p_o(r, m, 12)(-1)^r = -1.$$

In particular see cases $n = 11$ and $n = 12$, respectively in Table 2.

Table 2: The values of $p_o(r, m, N)$, where $N = 11$ and $N = 12$, respectively.

m	r	$p_o(r, m, 11)$	$p_o(r, m, 11)(-1)^r$	m	r	$p_o(r, m, 12)$	$p_o(r, m, 12)(-1)^r$
1	0	0	0	1	0	1	1
1	1	1	-1	1	1	0	0
2	0	0	0	2	0	2	2
2	1	2	-2	2	1	3	-3
2	2	3	3	2	2	0	0
3	0	0	0	3	0	1	1
3	1	2	-2	3	1	1	-1
3	2	3	3	3	2	2	2
3	3	0	0	3	3	3	-3
4	0	0	0	4	0	0	0
4	1	0	0	4	1	0	0
4	2	0	0	4	2	0	0
4	3	1	-1	4	3	1	-1
4	4	0	0	4	4	1	1

3 PROBLEM 6 GIVEN IN [1]

The main goal of this section is to present a solution to Andrews’s Problem 6, [1].

Definition 3.1. Given a partition λ of a positive integer $n = \lambda_1 + \lambda_2 + \dots + \lambda_m$ with $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_m$, whose lower odd parity index equals to $I_{lo}(\lambda) = r$, the weight of λ , denoted by $w(\lambda)$ is the real number given by $(-1)^{r+m}$.

Lemma 3.1. Consider $\lambda = \lambda_1 + \lambda_2 + \dots + \lambda_m$ a partition of the positive integer N , with $I_{lo}(\lambda) = r$ and $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_m$. Then: r is even (odd) if, and only if, λ_1 is even (odd).

Proof. The proof of this lemma is similar to the proof of Lemma 2.1. □

It follows from Lemma 3.1 that given any partition λ of a positive integer N ,

$$w(\lambda) = (-1)^{m+r} = (-1)^{m+\lambda_1},$$

where m is the number of parts and λ_1 is its greatest part.

In the following table we describe partitions λ of 7 with its respectives $I_{lo}(\lambda) = r$, number of parts m and weights $w(\lambda)$.

It follows from [1] that

$$U_o(y, x; q) = \sum_{r,m,n \geq 0} u_o(r, m, n) y^r x^m q^n \tag{3.1}$$

is the generating function for the partitions λ of the positive integer n in exactly m parts, with $I_{lo}(\lambda) = r$. Evaluating (3.1) in $x = y = -1$ we have:

$$U_o(-1, -1; q) = \sum_{r,m,n \geq 0} u_o(r, m, n) (-1)^{m+r} q^n. \tag{3.2}$$

Observe that the coefficient of q^7 in (3.2) is given by $\sum_{r,m \geq 0} u_o(r, m, 7) (-1)^{m+r}$. This is equivalent to sum the elements in the fourth column of Table 3, which in this case equals to 7.

Table 3: Partitions of 7 with its respectives $I_{lo}(\lambda) = r$.

partitions λ of 7	$I_{lo}(\lambda) = r$	m	$w(\lambda) = (-1)^{m+r}$
7	1	1	$(-1)^{1+1} = 1$
6 + 1	2	2	$(-1)^{2+2} = 1$
5 + 2	1	2	$(-1)^{1+2} = -1$
5 + 1 + 1	1	3	$(-1)^{1+3} = 1$
4 + 3	2	2	$(-1)^{2+2} = 1$
4 + 2 + 1	2	3	$(-1)^{3+2} = -1$
4 + 1 + 1 + 1	2	4	$(-1)^{4+2} = 1$
3 + 3 + 1	1	3	$(-1)^{3+1} = 1$
3 + 2 + 2	1	3	$(-1)^{3+1} = 1$
3 + 2 + 1 + 1	3	4	$(-1)^{4+3} = -1$
3 + 1 + 1 + 1 + 1	1	5	$(-1)^{5+1} = 1$
2 + 2 + 2 + 1	2	4	$(-1)^{4+2} = 1$
2 + 2 + 1 + 1 + 1	2	5	$(-1)^{5+2} = -1$
2 + 1 + 1 + 1 + 1 + 1	2	6	$(-1)^{6+2} = 1$
1 + 1 + 1 + 1 + 1 + 1 + 1	1	7	$(-1)^{7+1} = 1$

In [2] it is given a bijection between the set of unrestricted partitions and two-line matrices A defined as:

$$A = \begin{pmatrix} c_1 & c_2 & \dots & c_l \\ d_1 & d_2 & \dots & d_l \end{pmatrix}, \tag{3.3}$$

whose entries satisfy

$$\begin{cases} c_l > 0 \\ c_t \geq 2 + c_{t+1} + d_{t+1} & \text{with } t < l \end{cases},$$

with $c_t, d_t \geq 0$ and $n = \sum_{i=1}^l (c_i + d_i)$. Matrices defined in this way are presented as:

$$A = \begin{pmatrix} (2l - 1) + j_1 + \dots + j_l + d_2 + \dots + d_l & \dots & 3 + j_{l-1} + j_l + d_l & 1 + j_l \\ & d_1 & \dots & d_{l-1} & d_l \end{pmatrix},$$

$l \geq 1, s, d_1, \dots, d_l, j_1, j_2, \dots, j_l \in \mathcal{N}$.

Given a partition λ of a integer $n, n = \lambda_1 + \lambda_2 + \dots + \lambda_m$, the procedure to obtain the matrix A from λ [2] is given below:

- the side l of the Durfee square is the number of columns of the matrix;
- the parameters d_1, d_2, \dots, d_l indicate the amount of parts $1's, 2's, \dots, l's$, respectively, that appear below the Durfee square of side l .

$$\begin{cases} \lambda_1 = l + j_1 + j_2 + \dots + j_l \\ \vdots \\ \lambda_{l-1} = l + j_{l-1} + j_l \\ \lambda_l = l + j_l \end{cases},$$

where j_1, j_2, \dots, j_l are obtained in the solution of the system:

$$\begin{cases} c_1 = (2l - 1) + j_1 + j_2 + \dots + j_l + d_2 + \dots + d_l \\ \vdots \\ c_{l-1} = 3 + j_{l-1} + j_l + d_l \\ c_l = l + j_l \end{cases}.$$

In [3], the weight of the matrix A , given in (3.3) is defined by $w(A) = (-1)^{c_1+d_1+1}$. We will check that the weight of the matrix A has the same weight of the partition λ , associated to A by means of the bijection given in [3]. Indeed, $c_1+d_1+1 = (2l-1)+j_1+\dots+j_l+d_2+\dots+d_l+d_1+1 = 2l+j_1+\dots+j_l+d_1+d_2+\dots+d_l = (l+j_1+\dots+j_l) + (l+d_1+d_2+\dots+d_l)$. Observe that the greatest part of the partition is $\lambda_1 = l + j_1 + \dots + j_l$ and the number of parts of the partition is: $m = l + d_1 + d_2 + \dots + d_l$. Therefore, $w(A) = (-1)^{c_1+d_1+1} = (-1)^{\lambda_1+m} = (-1)^{r+m} = w(\lambda)$, where $r = I_{l_0}(\lambda)$. It follows from [3] that

Theorem 3.1. *The third order mock theta function $f(q) = \sum_{n=0}^{\infty} \frac{q^{n^2}}{(-q; q)_n^2}$, is the generating function for the weighted number of matrices A of the form (3.3). Where matrix is to be counted with the weight $w(A) = (-1)^{c_1+d_1+1}$.*

With these observations and together with Theorem 3.1 we have:

Theorem 3.2. The third order mock theta function $f(q) = \sum_{n=0}^{\infty} \frac{q^{n^2}}{(-q; q)_n^2}$, is the generating function for the weighted number of unrestricted partitions $\lambda = \lambda_1 + \lambda_2 + \dots + \lambda_m$ with $I_{l_0}(\lambda) = r$. Where each partition is to be counted with the weight $w(\lambda) = (-1)^{m+r}$ and m is the number of parts. From this from Theorem we have then, that

$$U_o(-1, -1; q) = \sum_{r, m, n \geq 0} u_o(r, m, n) (-1)^{m+r} q^n = \sum_{n=0}^{\infty} \frac{q^{n^2}}{(-q; q)_n^2}$$

RESUMO. Neste trabalho apresentamos uma solução para o Problema 5 de Andrews [1] por meio de uma bijeção entre os conjuntos D_n^0 and D_n^1 definidos no Teorema de Fine [8] e os conjuntos das partições indexadas pelo índice de paridade inferior [1]. Também resolvemos o Problema 6 de Andrews, conjecturado em [1].

Palavras-chave: partição, índice de paridade, mock theta function.

REFERENCES

- [1] G.E. Andrews. Parity in partition identities. *Ramanujan Journal*, **23** (2010), 45–90.
- [2] E.H.M. Brietzke, J.P.O. Santos & R. Silva. Bijective proofs using two-line matrix representation for partition. *Ramanujan Journal*, **23** (2010), 265–295.
- [3] E.H.M. Brietzke, J.P.O. Santos & R. Silva. Combinatorial interpretations as two-line array for the mock theta functions. *Bulletin Brazilian Mathematical*, **44** (2013), 233–253.
- [4] H. Göllnitz. “Einfache Partionen”. Thesis, Göttingen, (1960).
- [5] H. Göllnitz. Partitionen mit Differenzenbedingungen. *J. Reine Angew. Math.*, **225** (1967), 154–190.
- [6] B. Gordon. Some continued fractions of the Rogers-Ramanujan type. *Duke Math J.*, **31** (1965), 741–748.
- [7] I. Martinjak. Refinements and Extensions of the Euler Partition theorem. <http://imartinjak.files.wordpress.com/2013/06/refinementslecturenotes1.pdf>.
- [8] I. Pak. On Fine’s partition theorems, Dyson, Andrews and missed opportunities. *Mathematical Intelligencer*, **25**(1) (2003), p. 10.
- [9] R. Silva, J.C. Filho & J.P.O. Santos. Proving Two Partition Identities. *TEMA*, **13**(2) (2012), 133–142.
- [10] A.J. Yee & S. Kim. Göllnitz-Gordon Identities and Parity Questions in Partitions. *European Journal of Combinatorics*, **32**(2) (2011), 288–293.
- [11] A.J. Yee. Ramanujan’s partial theta series and parity in partitions. *The Ramanujan Journal*, **23** (2010), 215–225.
- [12] C. Wenchang. Two Problems of George Andrews on Generating Functions for partitions. *Miskolc Mathematical Notes*, **13**(2) (2012), 293–302.