



Propolis from Southwest of Parana produced by selected bees: Influence of seasonality and food supplementation on antioxidant activity and phenolic profile

MATHEUS A. CALEGARI¹, ANACLARA PRASNIEWSKI¹, CLEIDIANE DA SILVA¹, RICARDO Y. SADO², FABIANA M.C. MAIA², LARISSA M.S. TONIAL¹ and TATIANE L.C. OLDONI¹

¹ Departamento de Química, Universidade Tecnológica Federal do Paraná/UTFPR, Campus Pato Branco, Via do Conhecimento, Km 1, 85503-390 Pato Branco, PR, Brazil

² Departamento de Ciência Animal, Universidade Tecnológica Federal do Paraná/UTFPR, Campus Dois Vizinhos, Estrada para Boa Esperança, Km 4, 85660-000 Dois Vizinhos, PR, Brazil

Manuscript received on July 28, 2016; accepted for publication on September 29, 2016

ABSTRACT

Propolis produced by selected bees *Apis mellifera* were collected from March to June of 2013 and in March of 2015 and analyzed in order to evaluate the influence of climate, colony of origin, and food supplementation of colonies on the content of total phenolic and flavonoid by chromatographic analysis and antioxidant activity by radical scavenging of 2,2-diphenyl-1-picrylhydrazyl hydrate (DPPH) and 2,2'-azino-bis(3-ethylbenzothiazoline-6-sulphonic acid) (ABTS) and ferric reducing antioxidant power (FRAP) methods. The Principal Component Analysis (PCA) was carried out with propolis collected in 2013 and two clusters were formed. Propolis produced in the months of March and April showed a higher content of total phenolic compounds (TPC) and antioxidant capacity than those produced in May and June. The results of PCA obtained from samples collected in March of 2013 and 2015 showed two clusters, and propolis collected in 2015 were more bioactive and presented a higher content of TPC. The chromatographic analysis of extracts allowed the identification of phenolic acids *p*-coumaric, ferulic and caffeic with similar chemical profiles that could be closely related to the botanical origin of propolis. It can be concluded that the season and food supplementation of colonies influenced the chemical composition and the biological activity of samples analysed.

Key words: *Apis mellifera*, ethanolic extract, polyphenols, principal component analysis.

INTRODUCTION

Propolis is a natural resinous material collected by bees from different vegetable sources, and used to protect the hive against microorganisms, enclose small spaces and embalm foreign intruders (Daneshmand et al. 2015). It is widely known for presenting diverse biological properties such as anti-inflammatory (Luis-Villaroya et al. 2015,

Valenzuela-Barra et al. 2015, Wang et al. 2014), antibacterial (Gülçin et al. 2010, Luis-Villaroya et al. 2015, Oldoni et al. 2011), antioxidant (Cottica et al. 2015, Luis-Villaroya et al. 2015, Mahmoud et al. 2015, Oldoni et al. 2015), antifungal (Luis-Villaroya et al. 2015, Da Silva et al. 2015) and anticariogenic (Gülçin et al. 2010, Oldoni et al. 2011).

Phenolic compounds being their main components, and they have a high redox potential, which allows them to work as reducing agents

Correspondence to: Tatiane Luiza Cadorin Oldoni
E-mail: tatianeoldoni@utfpr.edu.br

(Miguel et al. 2010). The antioxidants act as defense agents against free radicals that are naturally present in organisms. Studies indicate that these compounds protect the organism from chronic diseases caused by oxidative stress, such as cancer and metabolic disorders (Wang et al. 2014, De-Melo et al. 2014, Kumazawa et al. 2004, Mohammadzadeh et al. 2007).

Studies described in the literature show that the chemical composition of propolis is varied and complex and it is intimately related to the biodiversity of flora, geographical region and climate where it is collected (Buratti et al. 2007, Gong et al. 2012, López et al. 2014). It is generally composed of resin, vegetable balm, wax, essential oils, pollen, and several other substances (Kalogeropoulos et al. 2009, Luis-Villaroya et al. 2015, Valenzuela-Barra et al. 2015).

Analysis of TPC and total flavonoids content (TFC) and the determination of antioxidant activity, contribute to the determination of both chemical and biological composition of propolis, allowing a relationship to be established with its geographical origin (Falcão et al. 2013). Nevertheless, a detailed discussion could be promoted if chemometric tools were used. Among them, PCA receives attention because it allows the evaluation of dataset, establishing relationships between variables, reducing its dimensions, and maintaining most of the statistical information (Randelovic et al. 2015).

Therefore, the goals of this study were to determine the influence of climate, origin of colony, and food supplementation of colonies in the phenolic profile by High-Performance Liquid Chromatography with Diode-Array Detection (HPLC-DAD), TPC and TFC and the antioxidant activity in Brazilian propolis produced by selected bees (by means of the animal model approach for weight at the emergence of Africanized queens) collected from March to June of 2013 and March of 2015, in addition to classifying these samples using the parameters evaluated, and the PCA tool.

MATERIALS AND METHODS

REAGENTS

The reagents DPPH, ABTS, (6-hydroxy-2,5,7,8-tetramethylchroman-2-carboxylic acid) (Trolox), 2,4,6-Tripyridyl-s-triazine (TPTZ) and authentic standards (grade HPLC) were acquired from Sigma Aldrich Co. (St. Louis, MO) with $\geq 99\%$ purity. All chemical products were of analytical grade and its aqueous solutions were prepared with ultrapure water (Milli-Q system).

SAMPLE COLLECTION

The samples of propolis were acquired from selected colonies of Africanized honey bees (Hymenoptera: Apidae) collected at an apiary within the Unidade de Ensino e Pesquisa de Apicultura (UNEPE) in Federal Technological University of Parana, Dois Vizinhos, Parana, Brazil (latitude: $-25,699063^\circ$, longitude: $-53,095273^\circ$, altitude: 546m). The collections were performed between March and June of 2013 and March of 2015. The selection of Africanized bees was done through generations and was based on the weight of queen at emergence. In 2013 the colonies were under systematic selection and in 2015 under relaxed selection. In 2015 the colonies received food supplementation every three days throughout the year. Samples were collected of propolis produced by fifth generation bees and obtained from three distinct colonies. The selection of the colonies used in this study was based on phenotypic reference to the potential of the bees.

The identification used in this study is described in Table I relates to the variables: month of collection, year, color, and colony.

SAMPLE PREPARATION

After collection, the samples were cleaned, crushed with liquid nitrogen, homogenized, weighed, and stored at -6°C .

TABLE I
Identification's table of samples of propolis.

Code of samples		Month of collection (2013)	Color	Colony
A	A1	03	Greenish brown	20
	A2	04	Greenish brown	
	A3	05	Red	
	A4	06	Red	
B	B1	03	Brown	02
	B2	04	Chesty	
	B3	05	Red	
	B4	06	Red	
C	C1	03	Greenish brown	44
	C2	04	Chesty	
	C3	05	Brown	
	C4	06	Brown	
Code of samples		Month of collection (2015*)	Color	Colony
D	D1	03	Red	20
	D2		Red	02
	D3		Red	44

*In the year of 2015, samples were collected only in March because was observed that propolis collected in March and April of 2013 showed better results and no significant difference.

PREPARATION OF ETHANOLIC EXTRACT OF PROPOLIS (EEP)

The EEP was prepared as described by Oldoni et al. (2015). Samples (2 g) of propolis were adding 25 mL of ethanol:water 80:20 (v v⁻¹) was added. The total volume was transferred to a thermostatically controlled water bath at 70 °C and held for 45 min. The samples were cooled and filtered using qualitative filter paper. All extracts and analyses were performed in triplicate.

ANTIOXIDANT ACTIVITY USING THE ABTS METHOD

The antioxidant activity was performed through the ABTS method according to Re et al. (1999) where the ABTS^{•+} radical was formed by the reaction of 7 mmol L⁻¹ of ABTS with potassium persulfate 140 mmol L⁻¹, incubated at 25 °C in the dark for 16 hours. The radical was diluted with ethanol and resulted in an absorbance value of 0.700 ± 0.200 at 734 nm. Under dark conditions, 3.0 mL of the ABTS^{•+} solution was added to 30 µL of EEP (16

g L⁻¹) and the absorbance was read at 734 nm at a spectrophotometer after 6 min. Ethanol was used as a blank and Trolox was used as reference in concentrations that varied from 100 to 2000 µmol L⁻¹. The results of antioxidant activity were equivalent to the antioxidant capacity of Trolox per gram of sample (µmol Trolox g⁻¹).

ANTIOXIDANT ACTIVITY USING THE DPPH FREE RADICAL SCAVENGING METHOD

The antioxidant activity by the free radical scavenging method DPPH was performed as described by Brand-Williams et al. (1995) where the reaction medium consisted in 0.5 mL of EEP (1.6 g L⁻¹), 3 mL of ethanol and 0.3 mL of radical DPPH solution 0.5 mmol L⁻¹ in ethanol. The mixture was incubated without light at room temperature for 45 min. and, later, the absorbance was measured using a spectrophotometer (model UV-VIS Lambda 25, Perkin Elmer) at 517 nm. The ethanol was used as blank and the quantification was performed based on an analytical curve using Trolox as standard in

concentrations that varied from 15 to 200 $\mu\text{mol L}^{-1}$. The results were expressed in μmol of Trolox per gram of propolis ($\mu\text{mol Trolox g}^{-1}$).

ANTIOXIDANT ACTIVITY USING THE FRAP METHOD

The antioxidant activity by the FRAP was proposed by Benzie and Strain (1996). The FRAP reagent was obtained from a mixture of 25 mL of acetate buffer 0.3 mol L^{-1} , 2.5 mL of a solution of TPTZ, 10 mmol L^{-1} and 2.5 mL of iron chloride in an aqueous solution at 20 mmol L^{-1} , used after preparation. The method consisted of a mixture of 100 μL of EEP (1.6 g L^{-1}) with 3 mL of FRAP reagent. The mixture was homogenized and kept in a water bath at 37 °C for 30 min. and, next, the absorbance in 595 nm was determined by a spectrophotometer (model UV-VIS Lambda 25, Perkin Elmer). The FRAP reagent was used as a blank and an aqueous solution of ferrous sulphate (100 - 2000 $\mu\text{mol L}^{-1}$) was used for the calibration curve. The results were expressed as μmol of Iron²⁺ ion per gram of propolis ($\mu\text{mol Fe}^{2+} \text{g}^{-1}$).

TOTAL PHENOLIC CONTENT

The TPC was determined through colorimetric analysis using the Folin-Ciocalteu reagent (Singleton et al. 1998). In a test tube, 0.5 mL of EEP (3.2 g L^{-1}) was added, 2.5 mL of Folin-Ciocalteu reagent diluted in 10 g L^{-1} and 2.0 mL of Na_2CO_3 at 40 g L^{-1} . After incubation without light for a period of 2 hours at room temperature, the absorbance was measured in a spectrophotometer (model UV-VIS Lambda 25, Perkin Elmer) at 740 nm. The blank was performed using ethanol:water 80:20 (v v^{-1}). The TPC of the extracts was determined by comparison with a standard calibration curve of gallic acid and represented in equivalent mg of gallic acid (EAG) per gram of propolis (mg EAG g^{-1}).

TOTAL FLAVONOID CONTENT

The TFC present in EEP was determined by applying the colorimetric method described by

Jurd and Geissman (1956). An aliquot of 0.5 mL of EEP (16 g L^{-1}) was added to a series of tubes identified as with or without nitrate. In the tubes that received nitrate, 4.3 mL of ethanol:water 80:20 (v v^{-1}) were added, and in those tubes that had not received nitrate, 4.4 mL of ethanol:water 80:20 (v v^{-1}) was added. In all tubes, 0.1 mL of potassium acetate 1 mol L^{-1} was added. In the series of tubes identified as “with nitrate”, 0.1 mL of aluminum nitrate at 10% was added. A control (blank) was also prepared from 4.9 mL of ethanol at 80% and 0.1 mL of potassium acetate. After 40 min., the absorbance was measured in a spectrophotometer (model UV-VIS Lambda 25, Perkin Elmer) at 415 nm. The TFC was determined by comparison with the calibration curve using quercetin as standard, and the results were expressed in mg of equivalent to quercetin (EQ) per gram of propolis (mg EQ g^{-1}).

HIGH-PERFORMANCE LIQUID CHROMATOGRAPHY WITH PHOTODIODE ARRAY DETECTION (HPLC-PDA)

For the analysis by HPLC-PDA a volume of 10 μL of EEP was injected at a concentration of 30 g L^{-1} in a liquid chromatography instrument coupled to a detector of photodiode array (PDA) (Varian 920-LC). An analytical column MICROSORB-MV 100 C18 was used; the composition of the mobile phase was constituted of (A) $\text{H}_2\text{O}:\text{CH}_3\text{COOH}$ 98:2 (v v^{-1}) and (B) $\text{H}_2\text{O}:\text{CH}_3\text{COOH}:\text{CH}_3\text{CN}$ 58:2:40 (v v^{-1}), and elution was made in the gradient mode starting with 5% solvent B to 7% of B in 7 minutes, 20% B in 15 minutes, 50% B in 30 minutes, 90% B in 50 minutes, and 95% B in 55 minutes with flow of 1 mL min^{-1} and the column was maintained at a constant temperature of 30°C. The identification was performed through comparison of the retention time and maximum absorbance wavelength with authentic standards.

MULTIVARIATE ANALYSIS

Multivariate analysis was performed with Pirouette software. The PCA charts were developed based

on the propolis characterization (ABTS, DPPH, FRAP, TFC and TPC). The pre-processing approach of the data used in this study was autoscaling, where each variable is centered in the average and divided by its standard deviation. Samples of propolis produced in the months from March to June 2013 were submitted to multivariate analysis, through the PCA to check the influence of seasonality in quality of material. After that, the results obtained from samples produced in March of 2013 and 2015 were compared with the main objective of verifying the influence of supplementation and year of production in the chemical composition of propolis.

RESULTS AND DISCUSSION

Propolis produced in March and April of 2013 showed a difference in color (Table I) and higher

results for antioxidant activity and content of TPC when compared with those produced in May and June (Table II) of the same year, indicating that there is a variation in chemical composition of the material and this variation can be related to the month of production. Among the samples of propolis collected in March of 2013 and 2015 the best results for TPC and TFC content as well as antioxidant activity (Table II) were obtained for propolis collected in March of 2015 suggesting that food supplementation of colonies can improve the quality of material. The antioxidant activity through the ABTS radical scavenging varied from 25.5 to 439.2 $\mu\text{mol Trolox g}^{-1}$, corresponding to samples C4 and D3 respectively.

Samples C4 and D3 also presented the lowest and highest capacity to scavenge the radical DPPH,

TABLE II
Total content of phenolic and flavonoid compounds and antioxidant activity determined by the FRAP, ABTS and DPPH methods.

Code	ABTS ($\mu\text{mol Trolox g}^{-1}$)	DPPH	FRAP ($\mu\text{mol Fe}^{2+} \text{g}^{-1}$)	Phenolic (mg EAG g^{-1})*	Flavonoid (mg EQ g^{-1})**
2013					
A1	85.5 \pm 1.33 ^c	39.2 \pm 5.39 ^b	243.9 \pm 29.18 ^b	17.9 \pm 0.73 ^a	1.23 \pm 0.09 ^b
A2	84.9 \pm 2.88 ^c	40.6 \pm 0.55 ^b	250.2 \pm 31.58 ^b	16.9 \pm 0.46 ^b	0.92 \pm 0.07 ^c
A3	89.1 \pm 2.90 ^b	41.9 \pm 0.94 ^b	229.4 \pm 7.85 ^c	18.3 \pm 0.40 ^a	1.00 \pm 0.06 ^c
A4	57.2 \pm 5.90 ^f	27.5 \pm 4.80 ^c	90.5 \pm 8.03 ^f	10.0 \pm 0.25 ^d	0.34 \pm 0.06 ^f
B1	78.2 \pm 0.81 ^d	25.3 \pm 0.46 ^d	121.0 \pm 6.13 ^c	10.0 \pm 0.46 ^d	0.62 \pm 0.03 ^c
B2	92.2 \pm 3.10 ^b	29.0 \pm 3.07 ^c	223.6 \pm 5.81 ^c	13.4 \pm 0.59 ^c	0.56 \pm 0.06 ^c
B3	49.1 \pm 4.16 ^h	23.2 \pm 1.94 ^d	112.5 \pm 6.19 ^c	10.2 \pm 0.47 ^d	0.48 \pm 0.03 ^c
B4	55.2 \pm 3.82 ^g	21.0 \pm 1.07 ^d	124.0 \pm 6.82 ^c	8.54 \pm 0.31 ^c	0.24 \pm 0.08 ^g
C1	109.2 \pm 1.19 ^a	49.4 \pm 2.03 ^a	286.7 \pm 7.88 ^a	17.0 \pm 0.79 ^b	2.19 \pm 0.06 ^a
C2	53.4 \pm 0.91 ^e	30.0 \pm 0.58 ^c	267.1 \pm 4.99 ^a	13.5 \pm 0.27 ^c	0.79 \pm 0.13 ^d
C3	61.5 \pm 1.78 ^c	21.0 \pm 1.05 ^d	152.7 \pm 5.56 ^d	8.67 \pm 0.37 ^c	0.65 \pm 0.16 ^c
C4	25.5 \pm 1.39 ⁱ	11.8 \pm 0.34 ^c	89.7 \pm 3.00 ^f	5.36 \pm 0.23 ^f	0.16 \pm 0.02 ^g
RSD (%)	4.18	8.26	7.51	3.87	10.90
2015					
D1	370.7 \pm 5.57 ^b	145.2 \pm 12.80 ^b	1041.0 \pm 95.44 ^a	39.4 \pm 3.92 ^b	3.03 \pm 0.19 ^b
D2	298.1 \pm 25.1 ^c	159.7 \pm 24.63 ^b	690.7 \pm 49.80 ^b	36.6 \pm 0.94 ^b	5.56 \pm 0.92 ^a
D3	439.2 \pm 3.51 ^a	235.6 \pm 8.14 ^a	638.1 \pm 17.64 ^b	48.2 \pm 1.29 ^a	1.02 \pm 0.13 ^c
RSD (%)	4.54	10.1	6.85	6.09	20.6

The same letter in the column, are not significantly different at the level of 0.05 according to ANOVA. The value is the average \pm standard deviation (n = 4). RSD: Relative Standard Deviation.

*equivalents to gallic acid.

**equivalent to quercetin.

11.8 and 235.6 μmol of Trolox g^{-1} , respectively (Table II). Kalogeropoulos et al. (2009) evaluated the antioxidant activity of ethanolic propolis extracts originating from Greece and Cyprus through the DPPH radical scavenging method and obtained as response values between 0.33 and 1.11 mmol Trolox g^{-1} , respectively.

Sample D1 produced in 2015 was the most successful in reducing the complex Fe^{3+} to Fe^{2+} , 1041.0 μmol Fe^{2+} g^{-1} , and presented high TFC (3.03 mg EQ g^{-1}) (Table II). However, sample C4, produced in 2013, showed the lowest antioxidant activity through the FRAP method and the lowest TFC, 89.7 μmol of Fe^{2+} g^{-1} and 0.16 mg EQ g^{-1} , respectively.

A high variability of results was also observed in propolis originating from Maringá – PR, which showed variation from 528 to 2068 μmol Fe^{2+} g^{-1} of propolis for antioxidant activity through the FRAP method, and from 10 to 26 mg EQ g^{-1} for the TFC (Cottica et al. 2011). In propolis produced in Transylvania, values between 740 and 2540 μmol Fe^{2+} g^{-1} were determined (Mihai et al. 2011).

The lowest and highest contents of TPC were observed for C4 and D3 samples, with 5.36 and 48.2 mg EAG g^{-1} , respectively. Propolis collected in Anhui, China, showed values for TPC that

varied from 174.7 to 235.6 μg EAG mg^{-1} (Yang et al. 2011). Matos et al. (2014) evaluated the content of TPC in propolis collected in the Northeast of Brazil and obtained values between 7.68 to 36.78 ± 1.52 mg EAG g^{-1} , corroborating the results of this study.

It can be concluded that there is a significant difference among propolis produced in different months and years by fifth generation bees. It is evident that propolis produced in 2015 is rich in phenolic compounds which should be directly related to higher antioxidant activity.

PCA is a technique used in multivariate analysis, allowing eased visualization of all information contained in a data set and classification of samples according to similarity, determination of objects showing different properties from others (outliers), and definition of important variables for classification (Morlock et al. 2014). In the current study, the first PCA (Figure 1) was performed using results obtained from samples collected in the summer and fall of 2013. Two principal components (PCs) described 92% of the total data variability. PC1 and PC2 described 86.2 and 5.8% of the variability respectively.

There were two general clusters on the 2D PC score (Figure 1a), one group corresponding




Figure 1 - Clustering on a 2D PC-score (a) and loading plot (b) based on the variables ABTS, DPPH, FRAP, total contents of phenolic and flavonoid compounds for propolis collected in 2013.

to samples collected in March and April, with the exception of sample A3, and the other group was composed of samples collected in May and June, with the exception of sample B1. All parameters evaluated had a positive correlation with PC1 and their concentration in sample collected in March and April was significantly higher. According to Figure 1b, PC2 showed a negative correlation with TPC and FRAP while DPPH, ABTS and TFC showed a positive correlation.

Nevertheless, results showed that is impossible to group the samples in relation to the colony, which indicates that the physical-chemical characteristics of the propolis cannot be associated with these classes.

The variations in the results for samples collected from March to June 2013 can be explained by variations in temperature observed in this period (Table III). Depending on the yield in the production of propolis in UTFPR - campus Dois Vizinhos, samples were collected between summer and fall, as well as in the transitional period between seasons. In southern Brazil there is a gradual decrease in temperature during this period and the change in average temperature directly influences the activity of bees, that is, the lower the temperature, the lower the activity of the bees will be. According to the data obtained by Jiang et al. (2016) activity of bees becomes more energetic when the temperature exceeds 25 °C and relative

humidity is between 60% and 70%. It was also observed that the bees leave their hives at sunrise and cease activities at sunset.

In 2013 changes were observed in the average temperatures of 5 °C and 6 °C for the minimum and maximum respectively between the months of March to June, with an average maximum above 25 °C in March and April. The propolis samples collected in the months with the lowest average temperature showed the lowest levels for TPC and TFC, as well as for the antioxidant activity. It is possible that due to the lower activity presented by bees and lower availability of material collected, product quality was affected, and as a result, different biological activities were observed.

In the study by McCall and Primack (1992), the shape of the flower, light, and temperature of the year are the most important variables that influence the rate of visitation of insects to flowers. Malerbo-Souza and Silva (2011) observed in their results that *Apis mellifera* bees have a different foraging behavior (insects searching for food) during the year. The bees collected less nectar in July 2007 (winter) and more in February 2008 (summer), which correlates with our results.

A second PCA was performed with samples collected in March of 2013 and 2015. The main objective was to compare the influence of year of production and food supplementation in the quality of propolis. Two PCs described 96.3% of the

TABLE III
Temperature and precipitation data in 2013 and 2015.

2013	Temperature (°C)		Precipitation (mm)
	Minimum mean	Maximum mean	Mean
March	17.629	27.384	28.446
April	14.947	27.023	10.760
May	13.300	23.410	20.333
June	12.927	21.287	39.488
2015			
March	18.722	30.084	109.832

Temperature and precipitation information obtained from the Meteorological Parana System (SIMEPAR), Hydroelectric Power Station Salto Osorio, the closest the UTFPR – Dois Vizinhos/PR.

total data variability. PC1 and PC2 described 75.8 and 20.5% of the variability respectively. Two general clusters on the 2D PC score (Figure 2a) were formed related to the samples of 2013 and 2015. All parameters evaluated had a positive correlation with PC1 and their concentration in samples collected in 2015 was significantly higher. The parameter flavonoid is related to the D2 sample, while results obtained by the FRAP method were higher for D1 sample. PC2 showed a negative correlation with TPC, ABTS and DPPH while FRAP and TFC showed a positive correlation (Figure 2b).

These results indicate that, despite different methods being used to evaluate the antioxidant activity, there is a strong positive correlation between content of TPC and antioxidant activity by DPPH and ABTS methods (Figure 2b), indicating that the antioxidant activity by these methods could be related to TPC content.

The phenolic profiles of EEP were identified by using High Performance Liquid Chromatography and the samples A2, A4 and D1 (Figure 3) are representative of summer and fall of 2013 and 2015 respectively and presented a chromatographic profile with differences in intensities of absorbance. The phenolic acids *p*-coumaric, ferulic and caffeic were identified and quantified (Table IV). The chromatographic method used to identify the TPC

was effective, as it showed a high selectivity and resolution for the majority of signals analyzed.

Samples showed a complex chemical composition with multiple signals mainly eluting with retention times of between 20 and 30 minutes, which indicates compounds with chemical structures of medium polarity (Figure 3). The phenolic acids caffeic, *p*-coumaric and ferulic were identified in the most extracts and from the identified compounds, which showed the highest levels. These results correlate with those obtained by Oldoni et al. (2015).

It is important to note the similarity of chemical profiles that samples produced in 2013 and 2015 showed, indicating that the botanical origin of the material analyzed is the same. The main difference between the propolis extracts was the content of phenolic acids identified in extracts (Table IV and Figure 3). The extracts D1, D2 and D3, produced in 2015 showed a higher content of phenolic acids caffeic, ferulic and *p*-coumaric. The D3 extract showed the highest levels of *p*-coumaric acid ($617.69 \mu\text{g g}^{-1}$) and ferulic acid ($305.56 \mu\text{g g}^{-1}$) and sample D1 showed higher content of caffeic acid ($150.84 \mu\text{g g}^{-1}$) indicating a positive correlation of these compounds with antioxidant activities determined by radical scavenger methods DPPH and ABTS (Table II).




Figure 2 - Clustering on a 2D PC-score (a) and loading plot (b) based on the variables ABTS, DPPH, FRAP, total contents of phenolic and flavonoid compounds for samples collected in March of 2013 and 2015.




Figure 3 - HPLC chromatograms obtained at 280 nm. **a)** sample A2; **b)** sample A4; **c)** sample D1; 1: caffeic acid, 2: *p*-coumaric acid, 3: ferulic acid. Spectral similarity comparison of the signals obtained from samples with the standard library is shown. Caffeic acid $R^2 = 99.25$; cumaric acid $R^2 = 99.77$; ferrulic acid $R^2 = 99.75$.

TABLE IV
Content of phenolic acids present in the extracts of propolis determined by HPLC-DAD.

Samples	Caffeic acid ($\mu\text{g g}^{-1}$)	<i>p</i> -coumaric acid ($\mu\text{g g}^{-1}$)	Ferulic acid ($\mu\text{g g}^{-1}$)
A1	126.7	n.d.	85.76
A2	121.0	n.d.	101.1
A3	124.3	n.d.	84.48
A4	128.3	53.55	64.58
B1	n.d.	43.22	51.32
B2	124.3	29.21	n.d.
B3	n.d.	38.55	62.37
B4	127.6	n.d.	56.85
C1	127.5	379.5	67.93
C2	124.3	47.89	n.d.
C3	117.7	57.23	44.25
C4	n.d.	66.6	45.79
D1	150.8	n.d.	211.6
D2	n.d.	206.7	145.3
D3	137.6	617.7	305.6

n.d.: not detected.

In 2013 and 2015 changes were observed in average temperatures of 1 to 3 °C for the minimum and maximum respectively. This variation may have had an impact on bee activity, as well as the daily supplementation provided in 2015. In addition, the temperature can influence the production of secondary metabolites in plants as a function of environmental variables, such as attack by microorganisms, temperature and UV radiation. Resins, flower buds and resinous exudates are sources of material for the production of propolis and a variation in the chemical composition of these materials means changes in the composition of propolis.

De Figueiredo et al. (2015) evaluated the chemical composition of propolis produced in Minas Gerais, Brazil by RP-HPLC and identified *p*-coumaric acid Li et al. (2016) obtained caffeic and ferulic acids in large quantities by HPLC in aqueous extract of propolis from China.

CONCLUSIONS

The PCA results provided two clusters for propolis produced in 2013 and it was possible to separate samples produced in warmer months from those produced in colder months. Among the samples collected in March of 2013 and 2015, it was possible to verify more bioactivity in those collected in 2015 from colonies that received food supplementation.

In addition, a positive correlation was verified between the TPC and the antioxidant activity through the DPPH and ABTS radical scavenging. The chromatographic analysis showed a similar chemical profile for propolis produced in different months and years, with identification of phenolic acids *p*-coumaric, caffeic and ferulic.

ACKNOWLEDGMENTS

The authors acknowledge fellowship from Conselho Nacional de Desenvolvimento Científico e Tecnológico – (CNPq) and Central de Análises

da Universidade Tecnológica Federal do Paraná (UTFPR).

REFERENCES

- BENZIE IF AND STRAIN JJ. 1996. The ferric reducing ability of plasma (FRAP) as a measure of “antioxidant power”: the FRAP assay. *Anal Biochem* 239: 70-76.
- BRAND-WILLIAMS W, CUVELIER ME AND BERSET C. 1995. Use of a free radical method to evaluate antioxidant activity. *LWT - Food Sci Technol* 28: 25-30.
- BURATTI S, BENEDETTI S AND COSIO MS. 2007. Evaluation of the antioxidant power of honey, propolis and royal jelly by amperometric flow injection analysis. *Talanta* 71: 1387-1392.
- COTTICA SM, SABIK H, ANTOINE C, FORTIN J, GRAVELINE N, VISENTAINER JV AND BRITTEN M. 2015. Characterization of Canadian propolis fractions obtained from two-step sequential extraction. *LWT - Food Sci Technol* 60: 609-614.
- COTTICA SM, SAWAYA ACHF, EBERLIN MN, FRANCO SL, ZEOULA LM AND VISENTAINER JV. 2011. Antioxidant Activity and Composition of Propolis Obtained by Different Methods of Extraction. *J Braz Chem Soc* 22: 929-935.
- DANESHMAND A, SADEGHI GH, KARIMI A, VAZIRY A AND IBRAHIM SA. 2015. Evaluating complementary effects of ethanol extract of propolis with the probiotic on growth performance, immune response and serum metabolites in male broiler chickens. *Livest Sci* 178: 195-201.
- DA SILVA RO, ANDRADE VM, RÊGO ESB, DÓRIA GAA, LIMA BS, DA-SILVA FA, ARAÚJO AAS, JÚNIOR RLCA, CARDOSO JC AND GOMES MZ. 2015. Acute and sub-acute oral toxicity of Brazilian red propolis in rats. *J Ethnopharmacol* 170: 66-71.
- DE FIGUEIREDO SM ET AL. 2015. Green Propolis: Thirteen Constituents of Polar Extract and Total Flavonoids Evaluated During Six Years through RP-HPLC. *Curr Drug Discov Technol* 12: 229-239.
- DE-MELO AAM, MATSUDA AH, FREITAS ADSD, BARTH OM AND ALMEIDA-MURADIAN LBD. 2014. Capacidade antioxidante da própolis. *Pes Agrop Trop* 44: 341-348.
- FALCÃO SI, TOMÁS A, VALE N, GOMES P, FREIRE C AND VILAS-BOAS M. 2013. Phenolic quantification and botanical origin of Portuguese propolis. *Ind Crops Prod* 49: 805-812.
- GONG S, LUO L, GONG W, GAO Y AND XIE M. 2012. Multivariate analyses of element concentrations revealed the groupings of propolis from different regions in China. *Food Chem* 134: 583-588.
- GÜLÇİN I, BURSAL E, ŞEHİTOĞLU MH, BILSEL M AND GÖREN AC. 2010. Polyphenol contents and antioxidant

- activity of lyophilized aqueous extract of propolis from Erzurum, Turkey. *Food Chem Toxicol* 48: 2227-2238.
- JIANG JA, WANG CH, CHEN CH, LIAO MS, SU YL, CHEN WS, HUANG CP, YANG EC AND CHUANG CL. 2016. A WSN-based automatic monitoring system for the foraging behavior of honey bees and environmental factors of beehives. *Comput Electron Agric* 123: 304-318.
- JURD L AND GEISSMAN TA. 1956. Absorption Spectra of Metal Complexes of Flavonoid Compounds. *J Org Chem* 21: 1395-1401.
- KALOGEROPOULOS N, KONTELES SJ, TROULLIDOU E, MOURTZINOS I AND KARATHANOS VT. 2009. Chemical composition, antioxidant activity and antimicrobial properties of propolis extracts from Greece and Cyprus. *Food Chem* 116: 452-461.
- KUMAZAWAS, HAMASAKA T AND NAKAYAMAT. 2004. Antioxidant activity of propolis of various geographic origins. *Food Chem* 84: 329-339.
- LIA, XUAN H, SUN A, LIU R AND CUI J. 2016. Preparative separation of polyphenols from water-soluble fraction of Chinese propolis using macroporous absorptive resin coupled with preparative high performance liquid chromatography. *J Chromatogr B Anal Technol Biomed Life Sci* 1012-1013: 42-49.
- LÓPEZ BGC, SCHMIDT EM, EBERLIN MN AND SAWAYA ACHF. 2014. Phytochemical markers of different types of red propolis. *Food Chem* 146: 174-180.
- LUIS-VILLAROYA A, ESPINA L, GARCÍA-GONZALO D, BAYARRI S, PÉREZ C AND PAGÁN R. 2015. Bioactive properties of a propolis-based dietary supplement and its use in combination with mild heat for apple juice preservation. *Int J Food Microbiol* 205: 90-97.
- MAHMOUD UT, ABDEL-RAHMAN MAM, DARWISH MHA, APPLGATE TJ AND CHENG HW. 2015. Behavioral changes and feathering score in heat stressed broiler chickens fed diets containing different levels of propolis. *Appl Anim Behav Sci* 166: 98-105.
- MALERBO-SOUZA DT AND SILVA FAS. 2011. Comportamento forrageiro da abelha Africanizada *Apis mellifera* L. no decorrer do ano. *Acta Sci Anim Sci* 33: 183-190.
- MATOS VR, ALENCAR SM AND SANTOS FAR. 2014. Pollen types and levels of total phenolic compounds in propolis produced by *Apis mellifera* L. (Apidae) in an area of the Semiarid Region of Bahia, Brazil. *An Acad Bras Cienc* 86: 407-418.
- MCCALL C AND PRIMACK RB. 1992. Influence of Flower Characteristics, Weather, Time of Day, and Season on Insect Visitation Rates in Three Plant Communities. *Am J Bot* 79: 434.
- MIGUEL MG, NUNES S, DANDLEN SA, CAVACO AM AND ANTUNES MD. 2010. Phenols and antioxidant activity of hydro-alcoholic extracts of propolis from Algarve, South of Portugal. *Food Chem Toxicol* 48: 3418-3423.
- MIHAI CM, MARGHITAS LA, DEZMIREAN DS AND BARNUTIU L. 2011. Correlation between Polyphenolic Profile and Antioxidant Activity of Propolis from Transylvania. *Anim Sci Biotechnol* 44: 100-103.
- MOHAMMADZADEH S, SHARRIATPANAHI M, HAMED M, AMANZADEH Y, EBRAHIMI SES AND OSTAD SN. 2007. Antioxidant power of Iranian propolis extract. *Food Chem* 103: 729-733.
- MORLOCK GE, RISTIVOJEVIC P AND CHERNETSOVA ES. 2014. Combined multivariate data analysis of high-performance thin-layer chromatography fingerprints and direct analysis in real time mass spectra for profiling of natural products like propolis. *J Chromatogr A* 1328: 104-112.
- OLDONI TLC, CABRAL ISR, D'ARCE MABR, ROSALEN PL, IKEGAKI M, NASCIMENTO AM AND ALENCAR SM. 2011. Isolation and analysis of bioactive isoflavonoids and chalcone from a new type of Brazilian propolis. *Sep Purif Technol* 77: 208-213.
- OLDONI TLC, OLIVEIRA SC, ANDOLFATTO S, KARLING M, CALEGARI MA, SADO RY, MAIA FMC, ALENCAR SM AND LIMA VA. 2015. Chemical Characterization and Optimization of the Extraction Process of Bioactive Compounds from Propolis Produced by Selected Bees. *J Braz Chem Soc* 26: 2054-2062.
- RANĐELOVIC SS, KOSTIC DA, ARSIC BB, MITIC SS, RASIC ID, MITIC MN, DIMITRIJEVIC DS AND STOJANOVIC GS. 2015. Chemometric Analysis of Grapes. *Open Chem* 13: 675-682.
- RE R, PELLEGRINI N, PROTEGGENTE A, PANNALA A, YANG M AND RICE-EVANS C. 1999. Antioxidant activity applying an improved ABTS radical cation decolorization assay. *Free Radic. Biol Med* 26: 1231-1237.
- SINGLETON VL, ORTHOFER R AND LAMUELA-RAVENTÓS RM. 1998. Analysis of total phenols and other oxidation substrates and antioxidants by means of folin-ciocalteu reagent. *Methods Enzymol* 299: 152-178.
- VALENZUELA-BARRA G, CASTRO C, FIGUEROA C, BARRIGA A, SILVA X, DE LAS HERAS B, HORTELANO S AND DELPORTE C. 2015. Anti-inflammatory activity and phenolic profile of propolis from two locations in Región Metropolitana de Santiago, Chile. *J Ethnopharmacol* 168: 37-44.
- WANG K, ZHANG J, PING S, MA Q, CHEN X, XUAN H, SHI J, ZHANG C AND HU F. 2014. Anti-inflammatory effects of ethanol extracts of Chinese propolis and buds from popular (*Populus×canadensis*). *J Ethnopharmacol* 155: 300-311.
- YANG H, DONG Y, DU H, SHI H, PENG Y AND LI X. 2011. Antioxidant Compounds from Propolis Collected in Anhui, China. *Molecules* 16: 3444-3455.

