



## Effects of different lipid levels on protozoa population, microbial protein synthesis and rumen degradability in cattle

Juliana Duarte Messana<sup>1\*</sup>, Telma Teresinha Berchielli<sup>1</sup>, Pedro Braga Arcuri<sup>2</sup>, Andressa Ferreira Ribeiro<sup>1</sup>, Giovani Fiorentini<sup>1</sup> and Roberta Carrilho Canesin<sup>1</sup>

<sup>1</sup>Departamento de Zootecnia, Faculdade de Ciências Agrárias e Veterinárias, Universidade Estadual Paulista "Júlio de Mesquita Filho", Via de Acesso Prof. Paulo Donato Castellane, s/n, 14884-900, Jaboticabal, São Paulo, Brazil. <sup>2</sup>Empresa Brasileira de Pesquisa Agropecuária, Setor Gado de Leite, Juiz de Fora, Minas Gerais, Brazil. \*Author for correspondence. E-mail: duarte\_juliana@hotmail.com

**ABSTRACT.** Protozoa population, microbial synthesis efficiency and rumen degradability of dry matter and neutral detergent fiber in cattle fed on diets with different lipid rates were evaluated. Nine 16-month-old Nelore young bulls, cannulated in the rumen and duodenum, weighing  $232 \pm 35$  kg, were used in the trial. Experimental design consisted of a 3 x 3 square in triplicate, comprising the following treatments: 2, 4 and 6% lipid in diet. In situ degradability was assessed by rumen incubation of corn silage, soybean, soybean meal and citrus pulp during 0, 3, 6, 12, 24, 48, 72 and 96h. The flow of microbial nitrogen and microbial efficiency were not influenced ( $p > 0.05$ ) by the inclusion of lipid levels in the diet. When the animals received diet with 4% lipid, there was a reduction ( $p < 0.05$ ) in the potential and effective degradation of dry matter of corn silage (64.10 and 55.04%, respectively), soybean meal (94.55 and 60.83%, respectively) and soybeans (98.45 and 76.44%, respectively). Since the degradability of neutral detergent fiber was not affected, 4 and 6% lipid levels in the diet may be used without altering the parameters of rumen degradation of fiber.

**Keywords:** degradability, fat, microorganisms, ruminants, soybean.

### Efeito de teores de lipídeos sobre a população de protozoários, síntese de proteína microbiana e a degradação ruminal em bovinos

**RESUMO.** Avaliaram-se a população de protozoários, a eficiência de síntese microbiana e a degradabilidade ruminal da matéria seca e da fibra em detergente neutro, em bovinos que consumiram dietas contendo diferentes teores de lipídeos. Foram utilizados nove novinhos da raça Nelore, canulados no rúmen e duodeno, com idade inicial 16 meses e  $232 \pm 35$  kg de peso corporal. O delineamento experimental utilizado foi quadrado latino 3 x 3 triplicado, sendo os tratamentos: 2; 4 e 6% de lipídeos na dieta. A degradabilidade "in situ" foi avaliada por meio da incubação ruminal da silagem de milho, soja grão, farelo de soja e polpa cítrica por 0, 3, 6, 12, 24, 48, 72 e 96h. O fluxo de nitrogênio microbiano e a eficiência de síntese microbiana não foram ( $p > 0,05$ ) influenciados pela inclusão de teores de lipídeos nas dietas. Quando os animais receberam 4% de lipídeo na dieta ocorreu redução ( $p < 0,05$ ) da degradação potencial e efetiva da matéria seca da silagem de milho (64,10 e 55,04%, respectivamente), farelo de soja (94,55 e 60,83%, respectivamente) e soja grão (98,45 e 76,44%, respectivamente). Teores 4 e 6% de lipídeos na dieta podem ser utilizados sem alterar os parâmetros de degradação ruminal da fibra.

**Palavras-chave:** degradabilidade, gordura, organismos, ruminantes, soja.

#### Introduction

Lipid supplementation in the diet has become a promising strategy to increase the efficiency in animal production by an increase in the diet's energetic density (HESS et al., 2008). However, the incorporation of lipids in diets is limited since they are modified by rumen microorganisms that transform them through two important processes, namely lipolysis and bio-hydrogenation. The processes' intensity largely depends on supplement sources and rates (MOATE et al., 2004).

Unsaturated fatty acids rates in the rumen above the saturation capacity of microorganisms produce adverse effects on rumen fermentation, such as decrease in fiber degradability, lowering of protozoa concentration, reduction in quantity and proportions of the short chain fatty acids (SILVA et al., 2007). The above effects may be due to the fiber's physical lining with lipids (JENKINS; MCGUIRE, 2006) and to the decrease in bacterial (in particular, gram-positive cellulolytic ones) and protozoa growth (TAMMINGA; DOREAU, 1991). Decrease in the number of protozoa is generally

associated with a reduction in bacterial N recycling in the rumen (MORAIS et al., 2006). Thus, an increase in the number of gram negative bacteria and a decrease in ammonia concentration occur (JOUANY, 1996), with a possible increase in the flow rate of rumen solids (CZERKAWSKI et al., 1975). These effects contribute towards an increase in the efficiency of microbial protein synthesis when lipids are supplemented for ruminants (DOREAU; FERLAY, 1995).

Diets with higher rates of ether extract are evaluated with regard to rumen protozoa, microbial protein synthesis and diet degradability.

### Material and methods

Experiment was conducted in the Food and Digestibility Evaluation Center of the Department of Animal Science of the Faculty of Agrarian and Veterinarian Sciences, UNESP, Jaboticabal, São Paulo State, Brazil. Nine 16-month-old gelded young Nelore bulls, cannulated in the rumen and the duodenum, mean initial weight  $232 \pm 35$  kg, were employed.

The animals were grouped within a 3 x 3 design with three simultaneous replications, three treatments and three experimental periods for 26 days each, or rather, 15 days for the animals' adaptation to diet, 7 days for sample collection and the last 4 days for in situ degradability evaluation. Experimental diets, formulated according to Fox et al. (2003) with Cornell Net Protein and Carbohydrate System, were as follows: 1) 2% ethereal extract in dry matter; 2) 4% ethereal extract in dry matter and 3) 6% ethereal extract in dry matter, composed of 50% corn roughage silage and of 50% concentrates. Table 1 shows the percentages of ingredients and the chemical composition of the diets.

Protozoa population was evaluated in samples of rumen contents collected 1h after feeding on the 18<sup>th</sup> day of each experimental period (VIDAL et al., 2007). Protozoa were counted directly in a Sedgewick-Rafter chamber (DEHORITY et al., 1989). Counts were obtained from aliquot of rumen contents conserved in formaldehyde, water solution and an equal volume of formaldehyde 37%. Samples were diluted with 20% glycerol and stained by Lugol solution for counting (D'AGOSTO; CARNEIRO, 1999).

Approximately 2 L of rumen contents from each animal were collected for bacterial isolation on the 19<sup>th</sup> day of each experimental period so that microbial protein synthesis may be evaluated (CECAVA et al., 1990). Purine bases were the microbial indicators and their determination in bacteria and duodenal digest was conducted according to recommendation by Zinn and Owens

(1986), with modifications by Ushida et al. (1985). Amount of microbial compounds in the duodenum was determined by N-RNA flow in the duodenum divided by the ratio N-RNA: total N of the rumen's isolated bacteria. Dry matter flux and microbial protein in the duodenum could be calculated.

Degradation of dry matter (DM) and of neutral detergent fiber (NDF) was evaluated by rumen incubation of corn silage, citric pulp, soybean meal and soybeans according to the in situ technique described by Orskov and McDonald (1979).

**Table 1.** Percentage of ingredients and chemical composition of experimental diets.

Ingredient (%)	Diet* (%DM)		
	2%	4%	6%
Corn silage	50.0	50.0	50.0
Citric pulp	28.0	26.0	24.0
Ground soybeans	0.0	12.0	23.0
Soybean meal	19.0	9.0	0.0
Mineral supplement <sup>1</sup>	3.0	3.0	3.0
<b>Chemical composition</b>			
Dry matter (DM)	63.74	63.76	63.78
Organic matter	91.60	91.81	92.00
Crude protein	15.26	14.28	13.45
Ether extract	1.93	4.23	6.32
Neutral detergent fiber	33.56	33.26	32.96
Acid detergent fiber	22.11	21.86	21.60
Lignin	2.72	2.67	2.63
Total Carbohydrates	74.26	73.15	72.06
Non-fibrous carbohydrates	40.70	39.90	39.10
	(Mcal kg <sup>-1</sup> )		
Gross energy	4.21	4.30	4.39
Digestible energy	3.09	3.15	3.09
Metabolized energy	2.53	2.58	2.53
Total digestible nutrients (%)	70.09	71.56	70.12

\*2, 4 and 6% = rate of lipids in diet. <sup>1</sup>Composition of mineral supplement: Ca:155 g; P: 80 g; Mg: 10g; S: 40 g; Na: 130 g; Cu: 1350 mg; Mn: 1040 mg; Zn: 5000 mg; I: 100 mg; Co: 80 mg; Se: 26 mg; F (maximum.): 800 mg; Solubility of P in citric acid at 2% (minimum): 90%.

Ingredients were incubated separately in the animals which received diets with increasing lipid rates: 2, 4 and 6% in DM so that their effect in the degradation of ingredients used in diets could be verified. During the last week of each experimental period nylon bags (7 x 14 cm), mean porosity 50  $\mu$ m, with approximately 5 g of sample, were incubated by cannulas in the animals' rumen for 3, 6, 12, 24, 48, 72 and 96h. Samples were pre-dried at 55°C during 72h and ground in a 5.0 mm sieve. During incubation, the bags were attached by stainless steel rings to nets and thus they could be fixed to the rumen cannula by nylon string with free movement within the rumen. After removal, the rumen bags were washed in running water till the wash liquid became colorless; they were then placed in buffers at 55°C for 72h for later weighing and DM and NDF analyses. Degradability at 0 time occurred when bags were immersed in a recipient with water at 39°C during 10 min. (CUMMINS et al., 1983).

Potential degradability (PD) was estimated by curves in the disappearance of the nutrients from the bags, adjusted according to model  $PD = a + b(1 - e^{-ct})$  suggested by Orskov and McDonald (1979), in which "a" is the soluble fraction; "b", the potentially degradable fraction; "c", the constant rate of degradation of "b" in time "t" of incubation. Real degradability was calculated by  $DE = a + (bc/c+k)$ , in which k is the flow rate of the rumen content at  $0.05 \text{ h}^{-1}$ .

Experimental design to evaluate protozoa population, microbial protein synthesis and degradability was a 3 x 3 square with three simultaneous repetitions, with three treatments and three periods, according to the following statistical model:

$$Y_{ijkl} = \hat{A}\mu + Q_i + T_j + P_k + Al(i) + QT_{ij} + e_{ijkl}$$

In which  $Y_{ijkl}$  is a dependent variable;  $\hat{A}$  is a mean of calculations;  $Q_i$  is the  $n^{\text{th}}$  effect of the square;  $T_j$  is the  $j^{\text{th}}$  of treatment or diet;  $P_k$  is the  $k^{\text{th}}$  effect of the line or period;  $Al(i)$  is the effect of the column or animal l, embedded in square i;  $QT_{ij}$  is the effect of the interaction between square i and treatment j;  $e_{ijkl}$  is the randomized error, presumed error distributed normally and independently (0, s<sup>2</sup>).

**Results and discussion**

Protozoa population was not influenced by 2, 4 and 6% lipid rates in the diet (Table 2). High coefficients of variation (CV) in the experiment are intrinsic to the method used (DEHORITY et al., 1989) since growth has not been reported in some counts, whereas a relatively large number of protozoa was observed in others (ORPIN, 1984). Gordon and Phillips (1989) and Messana et al. (2010) have reported coefficient variations similar to those in current experiment.

**Table 2.** Mean number of protozoa in the rumen of cattle fed on different lipid rates in the diet.

Mean population mL <sup>-1</sup>	Diet*			MSE**	VC***	Effects	
	2%	4%	6%			Linear	Square
Total Protozoa (no. x 10 <sup>5</sup> mL <sup>-1</sup> )	8.69	5.60	9.40	1.25	76.01	0.73	0.44
Entodinium	7.49	4.29	7.63	1.18	85.27	0.96	0.20
Epidinium	0.60	0.33	0.46	0.26	91.50	0.83	0.49
Isotricha	0.43	0.59	0.80	0.11	70.50	0.11	0.90
Dasytricha	0.17	0.39	0.51	0.19	70.79	0.51	0.18

\*2, 4 and 6% = Lipid rates in diet; \*\*Mean standard error; \*\*\*Variation Coefficient.

Rumen microorganisms are dependent on an ideal environment for their development with a temperature between 38 and 40°C and a pH range between 5.5 and 7.0 (HOOVER, 1986). The conditions reported in current analysis (pH between 6.6 and 6.4) may have been triggered to protozoa growth.

Protozoa of the genus *Entodinium* (Table 2) were found in absolute larger amounts in all treatments, as Valinote et al. (2006) have reported when they evaluated fatty acids as fat sources in cotton seeds and calcium salt, and the effect of monensin in diets with cotton seeds in diets for beef cattle. Predominance of *Entodinium* protozoa has also been reported by other authors who investigated protozoa populations in cattle with different food conditions, such as diet rich in sugarcane (FRANZOLIN; FRANZOLIN, 2000), in concentrate-rich diets, with or without fat addition (TOWNE et al., 1990), or with ionophores in diets rich in roughage or concentrates (GUAN et al., 2006; MARTINELE et al., 2008).

Several authors (VAN NEVEL; DEMEYER, 1988; CHAUDHARRY et al., 1995) reported that ciliated protozoa influence the degradation of structural carbohydrates, in particular, diets with high concentrate rates. Beside lipid adhesion to food particles and the toxic effect to fibrolytic bacteria, decrease in protozoa numbers may lessen the degradation of the diet fiber when lipids are inserted in the diet of ruminants, especially in diets with high concentrates. Consequently, treatments with increasing rates of lipids in the diet probably do not influence protozoa populations according to the proportion of fiber in the diet.

Results show that lipid rates did not affect (p > 0.05) intake of DM (IDM), nitrogen (IN) or quantities of organic matter (OMDR) or energy (EDR) digested in the rumen (Table 3).

**Table 3.** Efficiency of microbial protein synthesis in cattle fed on diets with different lipid rates.

Parameters	Diet*			MSE**	Effects	
	2%	4%	6%		Linear	Square
IDM (kg day <sup>-1</sup> )	5.39	6.07	5.65	0.24	0.58	0.21
IN (g day <sup>-1</sup> )	123.06	134.47	117.11	6.05	0.85	0.27
MODR (kg day <sup>-1</sup> )	3.39	4.04	3.67	0.18	0.43	0.13
EDR (Mcal day <sup>-1</sup> )	14.51	16.02	13.23	0.87	0.38	0.12
Flow to the duodenum						
MO (kg day <sup>-1</sup> )	1.74	1.78	1.82	0.09	0.29	0.70
N-mic (g day <sup>-1</sup> )	67.33	67.03	68.22	6.83	0.72	0.88
Efficiency of microbial synthesis						
g N kg <sup>-1</sup> MODR	18.97	17.19	19.34	0.22	0.95	0.72
g N Mcal <sup>-1</sup> of EDR	4.46	4.23	5.23	0.3	0.61	0.64

IMS = Ingestion of dry matter; IN = Ingestion of nitrogen; MODR = Organic matter digested in the rumen; EDR = Energy digested in the rumen; Nmic = Microbial Nitrogen. \*2, 4 and 6% = Lipid rates in diet. \*\*Mean standard error.

No difference in bacterial flow occurred; this was probably due to the fact that increasing lipid rates did not influence nitrogen consumption (Table 3). When ratio total N in the duodenum / total N ingested in diets with 2, 4 and 6% lipids is taken into account, 62.84, 60.55 and 64.60% of total N consumed reached the duodenum, respectively. Above results show that lipid rates did not significantly influence (p > 0.05) the efficiency of the microbial synthesis.

The provision of vegetal oils frequently causes an increase in the efficiency of microbial synthesis through a smaller predation by the oil's defaunated additive effects rather than by an increase in fermentable energy in the rumen (DEWHURST et al., 2000). On the other hand, results show that no defaunated effects caused by lipid rates were reported (Table 2). Thus, no difference was reported ( $p > 0.05$ ) in the efficiency of microbial synthesis (Table 3). Mean rate 18.50 g for Nmic kg<sup>-1</sup> of MODR of microbial efficiency synthesis is below the rates given by HAGEMEISTER et al. (1981). These authors registered efficiency rates of microbial synthesis of 18.0; 22.0; and 16.8 g 100 g<sup>-1</sup> of MODR for 0 to 20%, 30 to 70% and 70 to 100% rates of diet concentrates, respectively. These rates were higher than those in current analysis which employed different lipid rates in a diet with 50% concentrate. However, variation in efficiency rates of microbial synthesis in the literature (BÜRGER et al., 2000; HAGEMEISTER et al., 1981) is related to the type of microbiota and its growth phase, to nutrient availability, changes in diet and variations in feed frequency used in each experiment. The percentage of nutrients in the diet and food degradation synchronization which compose them is related to the efficiency of microbial synthesis (CLARK et al., 1992).

Lipid rates used influenced ( $p < 0.05$ ) the digestibility kinetics of DM degradability (Table 4).

**Table 4.** In situ degradability kinetics of DM and NDF of corn silage in cattle fed on different lipid rates in the diet.

Item <sup>1</sup>	Diet*				Effects	
	2%	4%	6%	EPM**	Linear	Square
	Dry Matter					
a (%) <sup>a</sup>	39.93	40.71	40.83	0.13	0.002	0.008
b (%) <sup>b</sup>	34.29	24.76	31.61	1.27	0.04	0.005
kd (% h <sup>-1</sup> ) <sup>c</sup>	2.29	2.50	3.42	0.20	0.01	0.08
DP (%) <sup>d</sup>	70.41	64.10	71.07	1.12	0.26	0.003
DE 0.05 (%) <sup>e</sup>	58.25	55.04	60.62	0.66	0.01	0.009
	Neutral detergent fiber					
b (%) <sup>f</sup>	58.10	57.18	52.10	0.94	0.004	0.02
kd (% h <sup>-1</sup> )	1.54	1.14	1.46	0.09	0.68	0.14
DP (%)	44.90	37.46	39.02	1.59	0.16	0.19
DE 0.05 (%)	25.32	20.52	21.89	1.02	0.19	0.18

<sup>1</sup>Soluble fraction (a), potentially degradable insoluble fraction (b), degradation rate of fraction b (kd), potential and (DP) and effective degradability (DE0.05) = effective degradability with presumed passage rate of 0.05 h<sup>-1</sup>; 2, 4 and 6% = Lipid rate of diet \*\*Mean standard error; <sup>a</sup>a = 38.37 + 0.98 a - 0.098 a<sup>2</sup>, r<sup>2</sup> = 0.97; <sup>b</sup>b = 57.20 - 15.05b + 1.79 b<sup>2</sup>, r<sup>2</sup> = 0.97; <sup>c</sup>kd = 1.63 + 0.28 kd, r<sup>2</sup> = 0.65; <sup>d</sup>DP = 90.0 - 13.11 DP + 1.66 DP<sup>2</sup>, r<sup>2</sup> = 0.97; <sup>e</sup>DE = 57.71 - 4.89 DE + 0.70 DE<sup>2</sup>, r<sup>2</sup> = 0.86; <sup>f</sup>b = 54.84 + 2.67 b - 0.52 b<sup>2</sup>, r<sup>2</sup> = 0.98.

Results for diets with 4 and 6% lipids agree with those by Palmquist and Jenkins (1980) who suggested a tolerance 4 to 5% more than total available energy by lipid supplement. A smaller potentially degradable fraction (b) in dry matter was reported when the animals were fed on diets with 4% lipids in DM. The same lipid rate also caused a lower potential degradability of DM of corn silage

when rates with 2 and 6% lipids are compared. Real degradability (DM) of corn silage was higher when animals received a diet with 6% lipids, perhaps due to their adjustment mode to the mathematic model used for the evaluation of results.

Although no significant difference in the degradability of NDF occurred, a significant decrease ( $p = 0.02$ ) occurred in the potentially degradable insoluble fraction (b) proportional to the increase of non-saturated fatty acids rates in the diet.

It seems that the mechanisms by which non-saturated fatty acids influenced the degradability of DM of corn silage occur regardless of the adsorption of bacteria and food particles, since NDF degradation was not altered. Similarly to current research, Broudiscou et al. (1990) reported that decrease in degradability was similar, regardless of covering or not of the cellulose with hydrolysed soybean oil.

Degradation decrease of silage DM with an increase in the diet's lipid rates found in current analysis failed to influence total digestibility of the DM of the diet. Even though a decrease in energy benefits of roughage feed in later treatment is taken into account, results show that, depending on the extension by which the fiber's rumen degradability is decreased, it may be compensated by a greater intestine digestion (WEISBJERG et al., 1992). Therefore, lipid dissociation in the rumen and their effects on the rumen degradability are closely related to pH rates of the rumen, or rather, with other diet components and their intake level (BALIEIRO NETO; MELOTTI, 2007).

When the kinetic degradation parameters of dry matter (DM) of citric pulp (Table 5) are taken into account, the soluble fraction in water at time 0 (a) was more appropriate for diet with 4% lipids. However, fraction "a" comprises two fractions: a soluble fraction in water and a fraction composed of solid particles that are released through the nylon bags' pores during washing. The difference between treatments may have been caused by this fact.

Although rumen degradability of dry matter of soybean meal decreased to the power of two when lipids were added to the diet (Table 6), degradability of neutral detergent fiber showed no difference among diet's lipid rates.

Real degradability of DM of soya meal was higher when the animals received 6% lipids in diet, perhaps due to adjustment to the mathematic model employed to evaluate results. When potential degradation is calculated for different timings, highest degradability occurred for the least lipid rate (2%) in diet.

**Table 5.** In situ degradability kinetics of DM and NDF of citric pulp in cattle fed on diets with different lipid rates.

Item <sup>1</sup>	Diet <sup>*</sup>			MSE <sup>**</sup>	Effects	
	2%	4%	6%		Linear	Square
	Dry matter					
a (%) <sup>a</sup>	45.17	45.78	45.65	0.12	0.04	0.12
b (%) <sup>b</sup>	53.02	47.49	51.37	0.15	0.02	0.07
kd (% h <sup>-1</sup> )	6.75	7.08	8.39	0.26	0.26	0.89
DP (%)	96.86	97.08	97.12	0.02	0.47	0.54
DE 0,05 (%)	73.10	77.50	77.73	1.40	0.27	0.65
	Neutral detergent fiber					
b (%) <sup>c</sup>	83.86	80.17	81.74	0.55	0.02	0.01
kd (% h <sup>-1</sup> ) <sup>d</sup>	3.74	3.49	3.76	0.05	0.76	0.04
DP (%) <sup>e</sup>	81.54	77.35	79.53	0.62	0.05	0.01
DE 0,05 (%) <sup>f</sup>	35.89	32.97	35.10	0.46	0.24	0.02

<sup>1</sup>Soluble fractions (a), potentially degradable non-soluble fraction (b), degradation rate of fraction b (kd), potential (DP) and real (DE0.05) degradability = real degradability with passage rate of 0.05 h<sup>-1</sup>; <sup>\*</sup>2, 4 and 6% = lipid rate in diet; <sup>\*\*</sup>Mean standard error <sup>a</sup> a = 44.96 + 0.15 a, r<sup>2</sup> = 0.51; <sup>b</sup> b = 52.32 - 0.18b, r<sup>2</sup> = 0.53; <sup>c</sup> c = 95.83 - 58.0 b + 0.66 b<sup>2</sup>, r<sup>2</sup> = 0.95; <sup>d</sup> kd = 4.51 - 0.51 kd + 0.065 kd<sup>2</sup>, r<sup>2</sup> = 0.84; <sup>e</sup> DP = 92.09 - 6.87 DP + 0.79 DP<sup>2</sup>, r<sup>2</sup> = 0.94; <sup>f</sup> DE = 43.87 - 5.26 DE + 0.63 DE<sup>2</sup>, r<sup>2</sup> = 0.89.

**Table 6.** In situ degradability kinetics of DM and NDF of soya meal in cattle fed on diet with different lipid rates.

Item <sup>1</sup>	Diet <sup>*</sup>			MSE <sup>**</sup>	Effects	
	2%	4%	6%		Linear	Square
	Dry Matter					
a (%)	33.66	33.62	33.64	0.14	0.93	0.96
b (%)	64.96	62.61	64.13	0.44	0.36	0.09
kd (% h <sup>-1</sup> ) <sup>a</sup>	5.63	4.88	7.36	0.53	0.007	0.002
DP (%) <sup>b</sup>	98.30	94.55	97.69	0.58	0.006	0.0001
DE 0,05 (%) <sup>c</sup>	68.02	60.83	71.79	1.68	0.02	0.003
	Neutral detergent fiber					
b (%)	96.75	97.74	97.84	0.20	0.06	0.22
kd (% h <sup>-1</sup> )	4.19	4.12	3.77	0.08	0.10	0.39
DP (%)	95.03	95.87	95.14	0.21	0.69	0.06
DE 0,05 (%)	44.14	44.17	42.01	0.49	0.13	0.28

<sup>1</sup>Soluble fraction (a), potentially degradable non-soluble fraction (b), degradation rate of fraction b (kd), potential (DP) and real (DE0.05) degradability = real degradability with estimated passage rate 0.05 h<sup>-1</sup>; <sup>\*</sup>2, 4 and 6% = diet's lipid rates; <sup>\*\*</sup>Mean standard error; <sup>a</sup>kd = 12.65 - 4.81 kd + 0.65 kd<sup>2</sup>, r<sup>2</sup> = 0.90; <sup>b</sup>DP = 108.9 - 7.04 DP + 0.86 DP<sup>2</sup>, r<sup>2</sup> = 0.99; <sup>c</sup>DE = 93.35 - 17.20 DE + 2.27 DE<sup>2</sup>, r<sup>2</sup> = 0.89.

Different rates of unsaturated fatty acids in the diet did not influence the degradability of soybeans (Table 7). In fact, it revealed degradation kinetics similar to that found by Silva et al. (2002) in research on animals supplemented with concentrated diet with ground soybeans.

The effect of unsaturated fatty acids rates tested on fiber degradability is extremely low or non-existent in most ingredients under analysis. Consequently, lipid rates did not influence diet degradation. Although ground soybean has been used, Silva et al. (2007) reported that release of lipids of oleaginous seeds is slow and provides small quantities of lipids within the rumen environment. This may cause fast bio-hydrogenation and thus may prevent the accumulation of unsaturated fatty acids. In fact, liability in rumen degradation, especially the diet fiber, is thus avoided. Further, the amount is not sufficient for the adhesion of food particle which may be a physical impairment to microorganism and microbial enzymes.

**Table 7.** In situ degradability kinetics of DM and NDF of soybeans in cattle fed on diet with different lipid rates.

Item <sup>1</sup>	Diet <sup>*</sup>			MSE <sup>**</sup>	Effects	
	2%	4%	6%		Linear	Square
	Dry matter					
a (%)	46.37	47.41	47.44	1.09	0.06	0.11
b (%)	53.71	51.13	51.51	1.14	0.07	0.08
kd (% h <sup>-1</sup> )	8.24	7.56	7.59	0.49	0.19	0.12
DP (%)	99.06	98.45	98.87	0.09	0.07	0.07
DE 0,05 (%)	78.72	76.44	78.41	0.49	0.75	0.11
	Neutral detergent fiber					
b (%)	96.39	94.82	96.56	0.31	0.75	0.06
kd (% h <sup>-1</sup> )	3.99	3.97	4.02	0.25	0.15	0.31
DP (%)	93.67	92.55	95.78	0.70	0.28	0.22
DE 0,05 (%)	42.39	41.85	48.38	1.56	0.18	0.30

<sup>1</sup>Soluble fractions (a), potentially degradable insoluble fraction (b), degradation rate of fraction b (kd), potential (DP) and real (DE 0.05) degradability rate = real degradability with estimated passage rate 0.05 h<sup>-1</sup>; <sup>\*</sup>2, 4 and 6% = lipid rates in diet; <sup>\*\*</sup>Mean standard error.

## Conclusion

Ground soybean with up to 6% lipids caused an increase in real degradability of silage roughage and a clear similarity in protozoa population, efficiency in microbial synthesis and fiber degradability. An inclusion of up to 6% lipids in the diet of Nelore young bulls may be thus employed.

## Acknowledgements

The authors would like to thank the Foundation for Research of São Paulo State (FAPESP) and Bellman Animal Nutrition for funding the research.

## References

- BALIEIRO NETO, G.; MELOTTI, L. Efeitos de níveis de sebo sobre a degradabilidade in situ do farelo de soja e do feno de Tifton (*Cynodon dactylon* (L.) Pers.) em vacas secas. **Brazilian Journal of Veterinary Research and Animal Science**, v. 44, n.4, p. 243-253, 2007.
- BROUDISCOU, L.; VAN NEVEL, C. J.; DEMEYER, D. L. Effect of soya oil hydrolysate on rumen digestion in defaunated and refaunated sheep. **Animal Feed Science and Technology**, v. 30, n. 1-2, p. 51-67, 1990.
- BÜRQUER, P. J.; PEREIRA, J. C.; VALADARES FILHO, S. C.; SILVA, J. F. C.; QUEIROZ, A. C.; CECON, P. R.; MAGIERO, D. Fermentação ruminal e eficiência microbiana em bezerras holandeses alimentados com dietas contendo diferentes níveis de concentrado. **Revista Brasileira de Zootecnia**, v. 29, n. 1, p. 215-224, 2000.
- CECAVA, M. J.; MERCHEN, N. R.; GAY, L. C.; BERGER, L. L. Composition of ruminal bacteria harvested from steers as influenced by dietary energy level, feeding frequency, and isolation techniques. **Journal of Dairy Science**, v. 73, n. 9, p. 2480-2488, 1990.
- CHAUDHARRY, L. C.; SRIVASTAVA, A.; SINGH, K. K. Rumen fermentation pattern and digestion of structural carbohydrates in buffalo (*Bubalus bubalis*) calves as affected by ciliate protozoa. **Animal Feed Science and Technology**, v. 56, n. 1-2, p. 111-117, 1995.

- CLARK, J. H.; KLUSMEYER, T. H.; CAMERON, M. R. Microbial protein synthesis and flows of nitrogen fractions to the duodenum of dairy cows. **Journal of Dairy Science**, v. 75, n. 8, p. 2304-2323, 1992.
- CUMMINS, K. A.; NOCEK, J. E.; POLAN, C. E.; HERBEIN, J. H. Nitrogen degradability and microbial protein synthesis in calves fed diets of varying degradability by the bag technique. **Journal of Dairy Science**, v. 66, n. 11, p. 2356-2364, 1983.
- CZERKAWSKI, J. W.; CHRISTIE, W. W.; BRECKENRIDGE, G.; HUNTER, M. L. Changes in the rumen metabolism of sheep given increasing mounts of linseed oil in their diet. **British Journal of Nutrition**, v. 34, n. 1, p. 25-44, 1975.
- D'AGOSTO, M. T.; CARNEIRO, M. E. Evaluation of lugol solution used for counting rumen ciliates. **Revista Brasileira de Zoologia**, v. 16, n. 3, p. 725-729, 1999.
- DEHORITY, B. A.; TIRABASSO, P. A.; GRIFO JR, A. P. Most-probable-number procedures for enumerating ruminal bacteria, including the simultaneous estimation of total and cellulolytic numbers in one medium. **Applied and Environmental Microbiology**, v. 55, n. 11, p. 2789-2792, 1989.
- DEWHURST, R. J.; DAVIES, D. R.; MERRY, R. J. Microbial protein supply from the rumen. **Animal Feed Science and Technology**, v. 85, n. 1-2, p. 1-21, 2000.
- DOREAU, M.; FERLAY, A. Effect of dietary lipids on the ruminal metabolism in the rumen: a review. **Livestock Production Science**, v. 43, n. 2, p. 97-110, 1995.
- FOX, D. G.; TYLUTKI, T. P.; TEDESCHI, L. O.; VAN AMBURGH, M. E.; CHASE, L. E.; PELL, A. N.; OVERTON, T. R.; RUSSELL, J. B. **The net carbohydrate and protein system for evaluating herd nutrition and nutrient excretion**. CNCPS version 5.0. Ithaca: Cornell University, 2003.
- FRANZOLIN, R.; FRANZOLIN, M. H. População de protozoários ciliados e degradabilidade ruminal em búfalos e bovinos zebuínos sob dieta a base de cana de açúcar. **Revista Brasileira de Zootecnia**, v. 29, n. 6, p. 1853-1861, 2000.
- GORDON, G. L. R.; PHILLIPS, M. W. Degradation and utilization of cellulose and straw by three different anaerobic fungi from the ovine rumen. **Applied and Environmental Microbiology**, v. 55, n. 7, p. 1703-1710, 1989.
- GUAN, H.; WITTENBERG, K. M.; OMINSKI, K. H. Efficacy of ionophores in cattle diets for mitigation of enteric methane. **Journal of Animal Science**, v. 84, n. 7, p. 1896-1906, 2006.
- HAGEMEISTER, H.; LÜPPING, W.; KAUFMANN, W. Microbial synthesis and digestion in the high-yielding dairy cow. In: HARESIGN, W. (Ed.) **Recent advances in animal nutrition**. London: Butterworths, 1981. p. 31-48.
- HESS, B. W.; MOSS, G. E.; HULE, D. C. A decade of developments in the area of fat supplementation research with beef cattle and sheep. **Journal of Animal Science**, v. 86, suppl., p. 188-204, 2008.
- HOOVER, W. H. Chemical factors involved in ruminal fiber digestion. **Journal of Dairy Science**, v. 69, n. 10, p. 2755-2766, 1986.
- JENKINS, T. C.; MCGUIRE, M. A. Major advances in nutrition: impact on milk composition. **Journal of Dairy Science**, v. 89, n. 4, p. 1302-1310, 2006.
- JOUANY, J. P. Effect of rumen protozoa on nitrogen utilization by ruminants. **The Journal of Nutrition**, v. 126, n. 4, p. 1335S-1346S, 1996.
- MARTINELE, I.; EIFERT, E. C.; LANA, R. P.; ARCURI, P. B.; D'AGOSTO, M. Efeito da monensina e do óleo de soja sobre os protozoários ciliados do rúmen e correlação dos protozoários com parâmetros da fermentação ruminal e digestivos. **Revista Brasileira de Zootecnia**, v. 37, n. 6, p. 1129-1136, 2008.
- MESSANA, J. D.; BERCHIELLI, T. T.; ARCURI, P. B.; REIS, R. A.; MALHEIROS, E. B.; RIVERA, A. R.; QUEIROZ, M. F. S. Effects of dry sugar cane yeast on the diet intake, digestibility and bovine rumen microbial populations. **Livestock Science**, v. 129, n. 1-3, p. 208-212, 2010.
- MOATE, P. J.; CHALUPA, W.; JENKINS, T. C.; BOSTON, R. C. A model to describe ruminal metabolism and intestinal absorption of long chain fatty acids. **Animal Feed Science and Technology**, v. 112, n. 1-4, p. 79-105, 2004.
- MORAIS, J. A. S.; BERCHIELLI, T. T.; REIS, R. A. Aditivos. In: BERCHIELLI, T. T.; PIRES, A. V.; OLIVEIRA, S. G. (Ed.). **Nutrição de ruminantes**. 1. ed. Jaboticabal: Funep, 2006. p. 539-570.
- ORPIN, C. G. The role of ciliate protozoa and fungi in the rumen digestion of plant cell walls. **Animal Feed Science and Technology**, v. 10, n. 2-3, p. 121-143, 1984.
- ORSKOV, E. R.; McDONALD, I. The estimation of protein degradability in the rumen from incubation measurements weighted according to rate of passage. **Journal of Agricultural Science**, v. 92, n. 2, p. 499-503, 1979.
- PALMQUIST, D. L.; JENKINS, T. C. Fat in lactation rations: review. **Journal of Dairy Science**, v. 63, n. 1, p. 1-14, 1980.
- SILVA, L. D. F.; RAMOS, B. M. O.; RIBEIRO, E. L. A.; MIZUBUTI, I. Y.; ROCHA, M. A.; MORAES, F. L. Z. Degradabilidade ruminal *In Situ* da matéria seca e proteína bruta de duas variedades de grão de soja com diferentes teores de inibidor de tripsina, em bovinos. **Revista Brasileira de Zootecnia**, v. 31, n. 3, p. 1251-1257, 2002.
- SILVA, R. M. C.; RODRIGUES, M. T.; BRANCO, R. H.; RODRIGUES, C. A. F.; SARMENTO, J. L. R.; QUEIROZ, A. C.; SILVA, S. P. Suplementação de lipídios em dietas para cabras em lactação: consumo e eficiência de utilização de nutrientes. **Revista Brasileira de Zootecnia**, v. 36, n. 1, p. 257-267, 2007.
- TAMMINGA, S.; DOREAU, M. Lipids and rumen digestion. In: JOUANY, J. P. (Ed.) **Rumen microbial metabolism and ruminant digestion**. Paris: Institut National de la Recherche Agronomique, 1991. p. 151-164.

- TOWNE, G.; NAGARAJA, T. G.; BRANDT JR., R. T.; KEMP, K. E. Ruminal ciliated protozoa in cattle fed finishing diets with or without supplemental fat. **Journal of Animal Science**, v. 68, n. 7, p. 2150-2155, 1990.
- USHIDA, K.; LASSALAS, B.; JOUANY, J. P. Determination of assay parameters for RNA analysis in bacterial and duodenal samples by spectrophotometry. Influence of sample treatment and preservation. **Reproduction and Nutrition Development**, v. 25, n. 6, p. 1037-1046, 1985.
- VALINOTE, A. C.; NOGUEIRA FILHO, J. C. M.; LEME, P. R.; SILVA, S. L.; CUNHA, J. A. Fontes de lipídio e monensina sódica na fermentação, cinética e degradabilidade ruminal de bovinos. **Pesquisa Agropecuária Brasileira**, v. 41, n. 1, p. 117-124, 2006.
- VAN NEVEL, C.; DEMEYER, D. I. Manipulation of rumen fermentation. In: HOBSON, H. D. (Ed.). **The rumen microbial ecosystem**. New York: Elsevier Science, 1988. p. 387-443.
- VIDAL, J. M.; PAIVA, P. C. A.; ARCURI, P. B.; LOPES, F. C. F.; ZAQUINI, L. Efeito de diferentes doses de enxofre no consumo voluntário e nas populações de protozoários do rúmen de novilhas mestiças alimentadas com capim-elefante de baixa qualidade. **Ciência e Agrotecnologia**, v. 31, n. 1, p. 218-222, 2007.
- WEISBJERG, M. R.; BORSTING, C. F.; HVELPLUND, T. The influence of tallow on rumen metabolism, microbial biomass synthesis and fatty acid composition of bacteria and protozoa. **Acta Agriculturae Scandinavica Section A - Animal Science**, v. 42, n. 1, p. 106-114, 1992.
- ZINN, R. A.; OWENS, F. N. A rapid procedure for purine measurement and its use for estimating net ruminal protein synthesis. **Canadian Journal of Animal Science**, v. 66, n. 1, p. 157-166, 1986.

*Received on March 10, 2011.*

*Accepted on July 14, 2011.*

License information: This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.