

# Physical Characteristics of Brazilian Wheat and their Utilization as an Indication of Flour Extraction

Rogério Germani\* and José Luiz Viana de Carvalho.

EMBRAPA Centro Nacional de Pesquisa de Tecnologia Agroindustrial de Alimentos - Av. das Américas, 29501, Guaratiba 23020-470 - Rio de Janeiro, RJ, Brazil.

## ABSTRACT

*An extensive work to determine the test weight (TW), the thousand kernel weight (TKW) and the hardness (HD) of Brazilian wheat, was undertaken for the crop years from 1990 to 1994, with the objective of drawing the Brazilian wheat physical characteristics and their relationship with flour extraction (FE). The results showed a median TW of 78.2kg/hl, a TKW of 34.2g and the HD was graded as hard. The values varied accordingly with the crop year, the local and year vs. local. Among the locals the wheat from the MS/DF/MG/GO States showed the best results. Correlation coefficients between FE and each of the parameters, TW, TKW and HD were low but significant ( $P < 5.0\%$ ) and, therefore, should not be used as a direct relationship but only as an indicator of FE. Segregation of TW, TKW and HD by specific ranges did not improve the correlation coefficient values.*

**Key words:** wheat; flour extraction; hectoliter weight, thousand kernel weight; hardness.

## INTRODUCTION

The measurement of the kernel apparent density (mass per volume unit) has been used as a parameter of wheat quality for commercial purpose in several countries (Zeleny, 1978). In Brazil, the kernel test weight (TW), measured as kilogram per hectoliter, was the only commercial quality parameter utilised until 1994. Although this parameter reflects very reasonably the grain sanity and its degree of development, it is also used as an indicator of the milling flour extraction (FE). Kremer (1993) found a high correlation between TW and FE while Johnson & Harris (1995) and Schuler *et al* (1995) did not find a significant correlation coefficient. Other parameters used as flour yield indicators are thousand kernel weight (TKW) (El-Shirbeeney & Allam, 1987) and kernel hardness (Baker & Dick, 1975). These works showed distinct results, however, they were conducted in countries where the conditions are very much different from Brazil and, therefore, can show different results. The objective of this work was to evaluate the relationship between the milling flour extraction and the kernel test weight,

the thousand kernel weight and the kernel hardness, from material grown in Brazil, as well as to relate the technological characteristics of Brazilian wheat to these parameters.

## MATERIALS AND METHODS

The wheat samples were obtained from the following experimental fields: FECOTRIGO-Federação da Cooperativas de Trigo de Cruz Alta-RS; EMBRAPA-CNPT (Centro Nacional de Pesquisa de Trigo), Passo Fundo-RS; Empresa de Pesquisa Agropecuária (EPAGRI), Santa Catarina state; Instituto Agrônômico do Paraná (IAPAR), Londrina-PR; Organização das Cooperativas do Paraná (OCEPAR), Cascavel-PR; EMBRAPA-CNPSO (Centro Nacional de Pesquisa de Soja), Londrina-PR; EMBRAPA-CPAO (Centro de Pesquisa Agropecuária do Oeste), Dourados-MS; Instituto Agrônômico de Campinas (IAC), SP and EMBRAPA-CPAC (Centro de Pesquisa Agropecuária do Cerrado), Brasília-DF. For the purpose of this work, the samples from MS, DF, MG and GO were considered together in one

---

\* Author for correspondence

group, since the recommended cultivars and the climatic conditions are similar. All the samples were conditioned to 15.0% moisture for 16-24hs prior to milling in a Brabender Qadrumat Senior mill, as described by Germani *et al* (1997). The moisture determination was done according to the method 44-15A from AACC (1983) and the ash content to the method 2.055 from AOAC (1984). Test weight was determined according to the equipment manufacturer (Dalle Molle) instructions, as described by Germani *et al* (1997) and, the thousand kernel weight, as described by SHUEY & GILLES (1972). The kernel hardness was tested using the Brabender micro hardness tester, as described by Kosmolak (1978), which measures the time required to grind 4g of wheat (for times below 32sec, the kernel is defined as hard, between 33 and 45sec, as semi-hard, between 46 and 63sec, as semi-soft, and above 63sec, as soft).


## RESULTS AND DISCUSSION

For this study 3250 samples of wheat from 1990, 1991, 1992, 1993 and 1994 crop year were used. Cumulative distribution of FE, TW, TKW and HD values are shown in Table 1. It could be seen that FE varied from 40.4 to 78.9%, with a median value of 64.5%. On the other hand, the TW showed the least variation, and its median was 78.2 kg/hl. For the TKW, the median value was 34.2g. The higher variation was observed with HD, which varied from 15 to 999sec, with a median value of 45sec. By the Kosmolak (1978) hardness scale, this median hardness value classified the wheat as hard. FE, TW and TKW distribution are illustrated in Figures 1, 2 and 3, respectively.


**Table 1.** Empirical cumulative distribution values regarding flour extraction (FE), test weight (TW), thousand kernel weight (TKW) and hardness (HD).

	Min.	5%	10%	25%	50%	75%	90%	95%	Máx.	CV%
FE	40.4	53.6	56.5	60.5	64.5	68.4	71.5	73.4	78.6	9.2
TW	60.2	69.8	72.1	75.2	78.2	80.4	82.1	83.1	84.9	5.3
TKW	17.3	25.2	27.0	30.3	34.2	38.2	41.7	44.0	56.0	16.9
HD*	15(VH)	17(VH)	18(VH)	22(VH)	41(H)	72(SS)	125(S)	211(S)	999(S)	132.9


\*VH=very hard. H=hard. SS=semi-soft. S=soft.



**Figure 1.** Flour extraction values distribution histogram.



**Figure 3.** Thousand kernel weight values distribution histogram



**Figure 2.** Test weight values distribution histogram.

Evaluating the local and crop year influence, it can be observed that there are significance differences among the samples cropped in the different states in different years as well as between the inter relation year/state for all parameters studied (Table 2).

A simple correlation analysis among FE, TW, TKW, HD and flour ash content are shown in Table 3. It could be observed that FE has a small

but positive correlation coefficient with TW and TKW, and a negative correlation with HD, showing that these parameters can not be straightly related, but only used as an indicator of FE. The low correlation found between FE and flour ash indicated that the FE values variation

was not caused by deficiency in the milling process. Schuler *et al* (1995) and Johnson & Harris (1995) did not find a significant correlation between FE and TW or TKW.

**Table 2.** Analysis of variance for the wheat kernel characteristics

Source of variation	D.F.	Chi square			
		Flour Extration	Test Weight	Thousand Kernel Weight	Hardness
Year	4	2115.2*	645.2*	976.6*	4306.7*
State	4	560.9*	109.7*	982.5*	14160.7*
Year/state	13	313.2*	184.3*	299.7*	5258.9*
Residous	25	4.9	3.4	5.1	20.3
C.V. (%)		7.69	4.4	149	41.4

\* Significance at P<5.0%.

To verify if the relations between FE and TW were influenced by the apparent kernel density, the coefficient of determination between these two parameters in several TW ranges was calculated as shown in Table 4. It was observed that the

correlation coefficients were still low and the higher ones were for the range of TW<70 and of TW<75 kg/hl, which were responsible for 19 and 12% of FE variation, respectively.

**Table 3.** Correlation coefficient among several parameters.

	Test Weght	Thousand		
		Kernel Weight	Hardness	Flour Ash
Flour Extration	0.317*	0.246*	-0.253**	0.162*
Test Weight		0.432*	-0.015 <sup>ns</sup>	-0.347*
Thousand Kernel Weight			0.025 <sup>ns</sup>	-0.175*
Hardness				-0.010 <sup>ns</sup>

• Significance at P<5.0%. ns = non significant.

**Table 4.** Coefficient of determination ( $R^2$ ) from the relation between flour extraction (FA) and test weight (TW)

TW range	$R^2$	Number of samples
TW<70	0.188*	168
70≤TW<75	0.040*	621
75≤TW<80	0.010*	1462
TW≥80	0.009*	950
TW<75	0.120*	789
TW<80	0.064*	2252
70≤TW<80	0.021*	2086
TW≥70	0.063*	3035
TW≥75	0.050*	2456
GLOBAL	0.102*	3202

\* Significance at  $P<5.0\%$ .

Similar segregation was performed with TKW (Table 5). In this case, any TKW value range gave expressive correlation coefficient with FE, since none of the ranges accounted for more than 6.0% of FE variation.

If the coefficient of determination was calculated between FE and HD, and the hardness values segregated by ranges (Table 6), it was observed that there was a great improvement in the  $R^2$  value which increased from 6.4% to 22.0%. Although the values of  $R^2$ , for the range of  $HD\leq 45$  and  $HD\leq 63$ , were similar to  $HD\leq 32$ , their contribution were irrelevant since the  $R^2$  values between these ranges were not significant.

The correlation between TW and TKW with FE was also calculated within hardness ranges (Table 7). It was observed that the influence of TW on FE was improved in the range of  $HD\leq 32$ sec, the range where it was classified as very hard. On the other hand, there was no expressive change for TKW.

**Table 7.** Variation of the coefficient of determination ( $R^2$ ) for the relation between flour extraction (FE) and test weight (TW), and between FE and thousand kernel weight (TKW), within the hardness ranges.

Range of hardness (sec)	(FE x TW)		(FE x TKW)	
	$R^2$	Number of samples	$R^2$	Number of samples
≤ 32	0.268*	1315	0.072*	1330
≤ 45	0.154*	1770	0.065*	1795
≤ 63	0.128*	2254	0.064*	2287
GLOBAL	0.102*	3202	0.060*	3239

\* Significance at  $P<5.0\%$ .

The material segregation by range of values, within each parameter, did not show any expressive improvement on the relationship with FE.

**Table 5.** Coefficient of determination ( $R^2$ ) for the relation between flour extraction (FE) and thousand kernel weight (TKW).

Range of TKW values	$R^2$	Number of Samples
TKW<30	0.059*	764
30≤TKW<35	0.010*	1049
35≤TKW<40	0.002 <sup>ns</sup>	922
TKW≥40	0.008*	505
TKW<35	0.054*	1813
TKW<40	0.053*	2735
30≤TKW<40	0.012*	1971
TKW≥30	0.020*	2477
TKW≥35	0.010*	1427
GLOBAL	0.060*	3240

\* Significance at  $P<5.0\%$ ; <sup>ns</sup>non significant.

**Table 6.** Coefficient of determination ( $R^2$ ) for the relation between flour extraction (FE) and hardness (HD).

Hardness range (sec)	$R^2$	Number of samples
$HD\leq 32$	0.215*	1298
$32<HD\leq 45$	0.006 <sup>ns</sup>	465
$45<HD\leq 63$	0.004 <sup>ns</sup>	491
$HD>63$	0.0002 <sup>ns</sup>	953
$HD\leq 45$	0.222*	1763
$HD\leq 63$	0.239*	2254
$32<HD\leq 63$	0.022*	956
$HD>45$	0.0008 <sup>ns</sup>	1478
$HD>32$	0.006*	1949
GLOBAL	0.064*	3207

\* Significance at  $P<5.0\%$ ; <sup>ns</sup>non significant.

Table 8 shows the median values obtained for the several crop years, within each parameter. It could be observed that there was a progressive increase in the kernel hardness from 1990 to 1994 crop year (Figure 4). This fact has been explained by the conscientiousness work, related to


technological quality of wheat, started in Brazil in 1989, since when several low quality cultivars were eliminated from planting recommendation and most of new releases were of proved high quality, as reported by Germani & Carvalho (1996).

**Table 8.** Median values for the wheat physical characteristics, for all states, by crop year.

Crop year	Number of samples	Median			
		Flour Extraction (%)	Test Weight (kg/hl)	Thousand Kernel Weight (gr)	Hardness* (sec)
1990	850	61.54 <sup>D</sup>	74.2 <sup>D</sup>	31.97 <sup>D</sup>	(56.6)SS <sup>A</sup>
1991	580	63.98 <sup>C</sup>	79.5 <sup>A</sup>	36.93 <sup>A</sup>	(52.2)SS <sup>B</sup>
1992	528	64.58 <sup>B</sup>	78.1 <sup>B</sup>	36.47 <sup>A</sup>	(48.3)SS <sup>C</sup>
1993	614	61.34 <sup>E</sup>	77.5 <sup>C</sup>	34.16 <sup>B</sup>	(45.9)SS <sup>D</sup>
1994	655	69.54 <sup>A</sup>	79.5 <sup>A</sup>	33.44 <sup>C</sup>	(40.0)H <sup>E</sup>

Mean followed by different letter are significantly different by the Duncan test at P<5.0%.

\* H=HARD, SS=semi-soft, S=soft.



**Figure 4.** Wheat hardness relative distribution. during the 90 to 94 crop years.

The parameters evaluation (by State) is shown in Table 9. It could be observed that the TW and TKW values were significantly higher for SP and MS/DF/MG/GO States, being the lowest for SC State. In terms of hardness, the wheat from MS/DF/MG/GO were the hardest, followed by those from SP, PR and RS States, and again being the softest those from SC State. For FE, the best results were also obtained from those sample from SP State, followed by MS/DF/MG/GO, PR and RS States, and those from SC with the lowest FE. This showed that there was a tendency of wheat of high TW, high TKW and high HD resulting in high FE. Therefore, samples from MS/DF/MG/GO States were generally of better quality than those from the other States (SP, PR and RS). Those from SC State showed the poorest quality.

Evaluating the correlation coefficient between FE and the other parameters by crop year and local (Table 10), it could be observed that there was variability among States by crop year, as already indicated by the interaction year/State (Table 7). This variability increased the difficulty of any indication of a tendency. In this segregation high correlation coefficient was obtained, as it was seen for TW in 1990, and for HD in 1992 and 1994 for the MS/DF/MG/GO States. The latter results were endorsed by Table 6, showing that the higher the hardness, the better the correlation with FE. Table 9 shows that the wheat grown in the MS/DF/MG/GO States was of high hardness. In spite of that the global evaluation by State did not show results as good as the ones obtained in distinct years.

**Table 9.** Median values for the wheat physical characteristics, for all years, by State.

State	Number of samples	Median			
		Flour Extraction (%)	Test Weight (kg/hl)	Thousand Kernel Weight (gr)	Hardness* (sec)
MS/ DF/ MG/GO	214	65.4 <sup>B</sup>	79.4 <sup>A</sup>	37.87 <sup>A</sup>	(33.9)H <sup>D</sup>
PR	1657	64.2 <sup>C</sup>	77.2 <sup>B</sup>	33.59 <sup>B</sup>	(46.7)SS <sup>C</sup>
SC	132	60.7 <sup>E</sup>	76.5 <sup>C</sup>	29.94 <sup>C</sup>	(63.0)S <sup>A</sup>
SP	282	68.0 <sup>A</sup>	79.0 <sup>A</sup>	38.57 <sup>A</sup>	(47.7)SS <sup>C</sup>
RS	955	63.2 <sup>D</sup>	77.2 <sup>B</sup>	34.13 <sup>B</sup>	(54.9)SS <sup>B</sup>

Median followed by different letter are significantly different by the Duncan test, at P<5.0%.

\* H=HARD, SS=semi-soft, S=soft.

**Table 10.** Correlation coefficient between flour extraction (FE) and test weight (TW), thousand kernel weight (TKW) and hardness (HD), by crop year and local.

Crop year		State				
		MS/DF/MG/GO	PR	RS	SC	SP
90	TW	0.834*	0.369*	0.245*	0.444*	- 0.519 <sup>ns</sup>
	TKW	0.614*	0.392*	0.266*	0.485*	0.242 <sup>ns</sup>
	HD	- 0.027 <sup>ns</sup>	- 0.095*	- 0.235*	- 0.104 <sup>ns</sup>	- 0.297 <sup>ns</sup>
91	TW	0.044 <sup>ns</sup>	0.009 <sup>ns</sup>	0.188*	N.s.a.	0.553*
	TKW	- 0.049 <sup>ns</sup>	0.085 <sup>ns</sup>	0.153*	N.s.a.	0.381 <sup>ns</sup>
	HD	- 0.676*	- 0.604*	- 0.368*	N.s.a.	- 0.479*
92	TW	- 0.393 <sup>ns</sup>	- 0.006 <sup>ns</sup>	- 0.312*	N.s.a.	0.067 <sup>ns</sup>
	TKW	- 0.176 <sup>ns</sup>	0.009 <sup>ns</sup>	0.218 <sup>ns</sup>	N.s.a.	0.166 <sup>ns</sup>
	HD	- 0.874*	- 0.287*	- 0.219 <sup>ns</sup>	N.s.a.	- 0.553*
93	TW	0.418 <sup>ns</sup>	0.147 <sup>ns</sup>	0.044 <sup>ns</sup>	0.191 <sup>ns</sup>	0.021 <sup>ns</sup>
	TKW	0.345 <sup>ns</sup>	0.317*	0.254*	0.060 <sup>ns</sup>	0.050 <sup>ns</sup>
	HD	- 0.448 <sup>ns</sup>	- 0.728*	- 0.123 <sup>ns</sup>	- 0.595*	0.243 <sup>ns</sup>
94	TW	0.142 <sup>ns</sup>	0.039 <sup>ns</sup>	0.261 <sup>ns</sup>	N.s.a.	0.259*
	TKW	0.479*	0.173*	0.370*	N.s.a.	0.197 <sup>ns</sup>
	HD	- 0.862*	- 0.565*	- 0.593*	N.s.a.	- 0.584*
GLOBAL	TW	0.442*	0.365*	0.137*	0.356*	- 0.022 <sup>ns</sup>
	TKW	0.200*	0.204*	0.187*	0.258*	0.084 <sup>ns</sup>
	HD	- 0.324*	- 0.255*	- 0.262*	- 0.022 <sup>ns</sup>	- 0.457*

\* Significance at P<5.0%. <sup>ns</sup>non significant. N.s.a.= no samples analysed.

## RESUMO

Um extensivo trabalho de determinação de peso do hectolitro grão (PH), peso de mil grãos (PMG) e dureza do grão (DZ), foi realizado durante as safras de 90 a 94, com o objetivo de traçar um perfil do trigo brasileiro e sua implicação na extração de farinha (EXFA). Obteve-se um valor médio de PH de 78,2kg/hl, de PMG de 34,2g e uma DZ classificada como semi-dura. Os valores encontrados variaram de acordo com o ano, estado

e, dentro do estado, nos diferentes anos. Entre os estados, os melhores resultados foram de amostras provenientes do MS/DF/MG/GO. Correlações entre EXFA e PH, PMG e DZ mostram ser significativas (P<5,0%) mas de baixo valor, não devendo, portanto, estes parâmetros físicos, ser utilizado em relação direta com a extração de farinha mas apenas como indicadores do potencial desta. Segregação dos valores de PH, PMG e DZ, por faixas, não melhoraram as correlações com EXFA.

## REFERENCES

- AACC - American Association Of Cereal Chemists (1983). *Approved methods of the American Association of Cereal Chemists*, 8. ed. St. Paul.
- AOAC - Association Of Official Analytical Chemists (1984). *Official methods of analysis of the Association of Official Analytical Chemists*, 15. ed. Gaithersburg.
- Baker, R.J.; Dyck, P.L. (1975), Relation of several quality characteristics to hardness in two spring wheat crosses. *Canadian Journal of Plant Science*, **55**, 625-627.
- El-Shirbeeney, A.; Allam, M. (1987), The thousand seed weight of wheat in relation to the technological properties of wheat. *Egyptian Journal of Food Science*, **15**(1), 83-90.
- Germani, R., Benassi, V.T.; Carvalho, J.L.V. de.(1997), *Metodologia de avaliação da qualidade tecnológica do grão e da farinha de trigo*. Rio de Janeiro: EMBRAPA-CTAA,. 43 p. Course apostille.
- Germani, R., Carvalho, J.L.V. (1996), Evolução do perfil da qualidade tecnológica das cultivares de trigo recomendadas para plantio no Brasil entre os anos de 91 e 96. Paper presented at 15<sup>th</sup> CONGRESSO BRASILEIRO DE CIÊNCIA E TECNOLOGIA DE ALIMENTOS, Poços de Caldas, MG.
- Johnson, J.M.; Harris, C.H. (1995), Test weight of soft red winter wheat related to flour yield and baking quality. Proceedings from *Institute of Food Technology Annual Meeting*, Chicago: Institute of Food Technology.
- Kosmolak, F.G. (1978), Grinding time: a screening test for kernel hardness in wheat. *Canadian Journal of Plant Science*, **58**, 415-420.
- Kremer, H. (1993), The effect of test weight on milling results. *Bulletin of the Association of Operative Millers*, 6189-6190.
- Schuler, S.F.; Bacon, R.K.; Finney, P.L.; Gbur, E.E. (1995), Characterization of test weight and its relationship to end-use quality of soft red winter wheat. *Crop Science*, **35**(4), 949-953.
- Shuey, W.C. & Gilles, K.A. (1972), Milling evaluation of hard red spring wheats. III. Relation of some physical characteristics of wheats to milling results. *The Northwestern Miller*, **279**(2), .141.
- Zeleny, L.( 1978), Criteria of wheat quality. In: *Wheat Chemistry and Technology*. ed. POMERANZ, Y. American Association of Cereal Chemists, St. Paul., pp. 19-50.

Received: June 02, 1998;  
Revised: November 16, 1998;  
Accepted: February 02, 1999.