

Expression analysis of three immune genes Interferon-gamma, Mx and Interferon regulatory factor-1 of Japanese flounder (*Paralichthys olivaceus*)

Feng Zhang¹, Xuemei Qiu^{1*}, Yang Liu¹, Juan Wang¹, Xipeng Li¹, Xiuli Wang¹.

¹ Dalian Ocean University - College of Fisheries and Life Science, Dalian, China.

ABSTRACT

IFN- γ (Interferon-gamma), *Mx* and *IRF-1* (Interferon regulatory factor-1) are main immune-related genes and they play important roles in the innate immune system of vertebrates. In this study, the expression level of the three immune-related genes in twelve tissues of normal adult Japanese flounder was detected using semi-quantitative RT-PCR. Thirteen time points (3h, 6h, 9h, 12h, 24h, 36h, 48h, 60h, 72h, 84h, 96h, 108h, 120h) were selected to analyze the expression of *IFN- γ* , *Mx* and *IRF-1* in spleen, head kidney, liver of Japanese flounder infected with Lipopolysaccharide (LPS). The Japanese flounder *IFN- γ* , *Mx* and *IRF-1* genes were differently expressed in these tissues and had high expression levels in classical fish immune organs like spleen and head kidney. It was found that the highest expression levels of the Japanese flounder *IFN- γ* were detected at 24h in spleen, 36h in head kidney and 48h in liver after challenge with LPS. Interestingly, the highest expression levels of *Mx* in spleen and head kidney were both at 36h and *IRF-1* in spleen and liver were both at 24h. The highest expression level of *Mx* in liver was at 48h and *IRF-1* in head kidney was at 12h. The study provides a basis for further research on immune mechanism of *IFN- γ* , *Mx*, *IRF-1* and the production of recombinant *IFN- γ* , *Mx* or *IRF-1* used in Japanese flounder cultivation in future.

Key words: Interferon-gamma, Mx, Interferon regulatory factor-1, *Paralichthys olivaceus*, tissue expression

* Author for correspondence: xuemeiqiu417@hotmail.com

Abbreviations: IFN- γ , Interferon-gamma; IRF-1, Interferon regulatory factor-1; LPS, Lipopolysaccharide; IPNV, infectious pancreatic necrosis virus; CCV, channel catfish herpesvirus; ISKNV, infectious spleen and kidney necrosis virus; ISRE, Interferon-stimulated response element; ISGs, interferon stimulated genes; DBD, DNA-binding domain; MHC, major histocompatibility complex; PS, physiological saline; IOD, integrated optical density; CK, uninfected adult's tissues; IHNV, infectious hematopoietic necrosis virus; HRV, Hiram rhabdovirus; HINAE, Hiram natural embryonic;

INTRODUCTION

The immune system is commonly divided into the innate (non-specific) and the adaptive (specific) immune system. The innate response generally precedes the adaptive response and determines the nature of the adaptive response¹ and². The fish immune system is very different from mammals and the innate immune system of fish is the primarily important defense barrier since the adaptive immune system is inefficient due to its evolutionary status³. Compared with the adaptive immune system, the innate immune response is instant and relatively temperature independent^{4,5}. The main parameters of the innate immune system of fish include lectins, lysozyme, complement proteins, antibacterial peptides, IgM, transferrin, interferon and Mx protein^{5,6,7}.

Interferon (IFN) which plays an important role in innate immune system is a protein secreted from cells in response to invasion by pathogens and virus. In mammals, IFN is classified into type-I IFN (α , β , ϵ , δ , τ and ω), type-II IFN (γ) and type-III IFN (λ) based on their differences in primary structure, serology and their most likely time of evolutionary divergence^{8,9,10}.

Interferon-gamma (IFN- γ) is a pleiotropic proinflammatory and antiviral cytokine, which was initially identified in PHA-activated mammalian lymphocyte supernatants¹¹. It was demonstrated that IFN- γ is a central participant in host resistance to various infections by using IFN- γ knockout mice infected with *Leishmania major*¹², *Listeria monocytogenes*¹³ and *Mycobacteria*¹⁴.

Mx proteins present a 70-80kDa GTPases of dynamin super family among vertebrates. The Mx gene in fish was first detected in 1989¹⁵ and then seven forms of Mx genes were found in zebrafish¹⁶, three forms in rainbow trout¹⁷, two forms in crucian carp (*Carassius auratus*)¹⁸ and a single form in Japanese flounder (*Paralichthys olivaceus*)¹⁹ and Pufferfish²⁰. However, the antiviral property of fish Mx proteins had been demonstrated only in Japanese flounder and Atlantic salmon against fish rhabdoviruses, infectious pancreatic necrosis virus, and viral haemorrhagic septicaemia virus^{21,22,23}.

Interferon regulatory factors (IRFs) were transcription factors which bound to the ISREs (Interferon-stimulated response element) and mainly regulated the expression of type-I IFN gene, interferon stimulated genes (ISGs) and other cytokines^{24,25}. Up to now as many as ten members of IRFs had been reported in vertebrates²⁶ and they all showed significant homology within the N-terminal 120 amino acids which contained the DNA-binding domain (DBD) forming a helix-turn-helix structure²⁷. Over the last two decades, these IRFs have been the focus of many immunological as well as medical studies²⁶ and it has been confirmed that they play important roles in antiviral activity and immune regulation^{28,29}.

Several IRFs have been characterized in model fish and some economically important fish species. For example, IRF-1 had been cloned in Japanese flounder³⁰ and later in rainbow trout³¹, turbot (*Scophthalmus maximus*)³² and mandarin fish (*Siniperca chuasti*)³³.

Japanese flounder is an important economic marine fish species in the world due to its rapid growth, good taste and high nutritive value. In recent years, the scale of Japanese flounder cultivation has been expanding continuously in China. In northern China, the seawater temperature is comparatively lower than in southern China, especially in winter. The lower temperature might affect the growth of Japanese flounder and its immune response. In this study, the expression regularity of the three immune genes (IFN- γ , Mx and IRF-1) of Japanese flounder at different time points and different tissues after infection with Lipopolysaccharide (LPS) at lower temperature was well revealed. This study might be helpful for further research on the antiviral mechanism of these three immune genes and for the production and application of relevant recombinant IFN- γ , Mx and IRF-1.

MATERIAL AND METHODS

Japanese flounder

Normal adult individuals weighing 700g on average were raised at Key Laboratory of Mariculture, Ministry of Agriculture, Dalian Ocean University. They were kept in two tanks (15m³) separately for one week before experimentation and the water temperature was 8 °C to 10 °C.

Challenge and sampling

Lipopolysaccharide (LPS) (sigma, L2880, 055:B5), the major constituent of the external layer of Gram-negative bacteria, was dissolved with physiological saline (PS) to a concentration of 1mg/ml. Thirty-nine Japanese flounders were injected intraperitoneally with LPS (400ug/kg)³⁴ as the experimental group and thirteen were injected with an equal dose of PS as the control group. Then three LPS-infected fish and one PS-injected fish were sacrificed at each sampling time (3h, 6h, 9h, 12h, 24h, 36h, 48h, 60h, 72h, 84h, 96h, 108h and 120h) after infection respectively. Tissues of interest (spleen, head kidney and liver) were dissected, quickly frozen in liquid nitrogen and stored at -80 °C until RNA extraction. Meanwhile, two normal adult fish were sacrificed and tissues of heart, blood, spleen, head kidney, brain, intestine, kidney, liver, gill, skin, muscle and gonad were removed and kept at -80°C until use³⁵. All procedures involving the handling and treatment of the individuals during this study were approved by the Animal Care and Use committee of Key Laboratory of Mariculture & Stock Enhancement in North China's Sea at Dalian Ocean University.

Primer design

According to IFN- γ (GenBank accession number AB435093), Mx (AB110446) and IRF-1 (AB005883) gene of Japanese flounder on NCBI, three pairs of primer (IFN-F/R, Mx-F/R, IRF-F/R) were designed separately (Table 1). The glyceraldehyde-3-phosphate dehydrogenase (GAPDH) (AB029337) gene of Japanese flounder was selected as the internal control and a pair of primer (GAPDH-F/R) was designed (Table 1).

Table1 Primer information of IFN- γ , Mx, IRF-1 and GAPDH gene of Japanese flounder used for semi-quantitative RT-PCR

Primer ID	Primer sequence	Gene	Accession number	Fragement length	Tm(°C)	Location*
IFN-F	5'-TGGTCTGTCTGTCCCTGTG-3'	IFN- γ gene	AB435093	371bp	56	26-396
IFN-R	5'-CTTGAAGCGATTTGTCCTC-3'					
Mx-F	5'-AAGAGGAAAAGGCATAGTG-3'	Mx gene	AB110446	395bp	57	177-571
Mx-R	5'-CCTCTGTGGTTGCTATGTC-3'					
IRF-F	5'-AGCAAAGGTCACAAAGCAG-3'	IRF-1 gene	AB005883	303bp	59	366-668
IRF-R	5'-ACAATGTCGTCGGAGTAGC-3'					
GAPDH-F	5'-ATGCTGGTGCCCACTATGT-3'	GAPDH gene	AB029337	577bp	56	260-836
GAPDH-R	5'-ACCTGGTGCTCGGTGTATG-3'					

*Location represents the position of the Japanese flounder IFN- γ , Mx, IRF-1 and GAPDH gene sequence, respectively.

Isolation of RNA

Total RNA was extracted using TRIzol reagent (Invitrogen) according to the reagent instructions. Briefly, 50mg samples were homogenized with 1ml TRIzol reagent and after incubation at 20°C for 5 min, 0.2ml of chloroform was added. The tubes were shaken vigorously by hand for 15 seconds and incubated them at 20°C for 3min. Then the mixture was centrifuged at 12000rpm, 4°C for 15 min. Subsequently, supernatant was collected and 0.5ml of isopropyl alcohol was added. After incubation at 20°C for 10min, the sample was centrifuged at 12000rpm, 4°C for 10 min. Finally, the precipitation was collected, washed with 1ml 75% ethanol once, then dried at room temperature and dissolved in RNase-free water. The dissolved RNA was kept at -80°C for cDNA synthesis.

Semi-quantitative RT-PCR analysis of IFN- γ , Mx and IRF-1 gene expression

cDNA synthesis was carried out using M-MLV Transcriptase (TaKaRa) according to the manufacture's instructions. Total RNA from the same time points or tissues were equally mixed for reverse transcription, separately.

Primers (IFN-F/R, Mx-F/R, IRF-F/R and GAPDH-F/R) were separately used for amplifying IFN- γ fragment of 371bp, Mx fragment of 395bp, IRF-1 fragment of 303bp and GAPDH fragment of 577bp. Polymerase chain reaction (PCR) were used for amplification with conditions as follows: 1 cycle of 5 min at 94°C; 20-25 cycles of 30s at 94°C, 30s at 56°C-59°C (Details seen in Table 1), 45s at 72°C; 1 cycle of 7 min at 72°C. PCR products were analyzed in a 1.0% agarose gel stained with ethidium bromide and the amplified DNAs were recorded with the Kodak Image Station 440CF. The integrated optical density (IOD) ratios of IFN- γ , Mx and IRF-1 versus GAPDH were determined with Gel-Pro analyzer to analyze the relative mRNA expression³⁶ and³⁷. Each column represents the average value of triple determinations.

RESULTS

IFN- γ , Mx and IRF-1 expression in twelve tissues of normal adult Japanese flounder

Semi-quantitative RT-PCR was conducted to analyze expressions of the IFN- γ , Mx and IRF-1 gene in normal Japanese flounders. The expression level of Japanese flounder IFN- γ was the highest in blood and high in spleen, head kidney, kidney and gill. The expression level was very faint in heart, brain, intestine, skin, liver and muscle. And IFN- γ was not detected in gonad (Fig.1 A).




Figure 1. Semi-quantitative RT-PCR analysis of IFN- γ , Mx and IRF-1 mRNA expression in various tissues of Japanese flounder. A, B, C: Expression of Japanese flounder IFN- γ , Mx, IRF-1 in various tissues, respectively; D:

Densitometric analysis of semi-quantitative RT-PCR was used to obtain the relative IOD of IFN- γ , Mx and IRF-1 versus GAPDH; M: molecular weight standard; 1-12: heart, blood, spleen, head kidney, brain, intestine, kidney, skin, Gill, liver, muscle, gonad. The Japanese flounder IFN- γ gene-specific RT-PCR products (371bp), Mx gene-specific RT-PCR products (395 bp) and IRF-1 gene-specific RT-PCR products (303 bp) are indicated on the right margin, along with GAPDH RT-PCR products (577 bp) as an internal control.

The expression level of Japanese flounder Mx was the highest in blood and head kidney and was high in heart, spleen, intestine, kidney and gill. The expression of Mx was very faint in liver and not detected in brain, skin, muscle and gonad (Fig.1 B).

Seen from Figure.1 C, the expression of Japanese flounder IRF-1 was detected in all tissues and was very high in blood, spleen, intestine, kidney, gill, muscle and gonad.

IFN- γ expression in infected Japanese flounder tissues

The expression of Japanese flounder IFN- γ was analyzed in the spleen, head kidney and liver after challenge with LPS. In the three tissues, the expression of IFN- γ was dramatically induced after infection. The highest expression levels of the IFN- γ were detected at 24h in spleen, 36h in head kidney and 48h in liver. The expression in spleen started at 12h, decreased gradually at 36h and returned to the level of background after 48h. The expression levels of IFN- γ in spleen at 24h, 36h and 48h were approximately 3.2-, 4.3- and 2.8-fold compared with the uninfected adult's tissues (CK), respectively. At the rest time points, no significant increases of the gene were observed (Fig.2 A, G). In head kidney, the expression levels kept high from 12h to 48h and then decreased gradually to background. The relative IOD value showed the expression levels from 12h to 48h were approximately 4.9-, 8.8-, 9- and 7.1-fold compared with CK (Fig.2 C, G). The high expression level in liver lasted from 24h to 84h and then became faint consistent with background. The highest expression level was at 48h and it was 4.5-fold compared with CK (Fig.2 E, G). We also injected the Japanese flounder with physiological saline (PS) as control and the results showed that the IFN- γ expression in spleen, head kidney and liver was consistent with CK, indicating the injection of PS was not harmful to Japanese flounder (Fig.2 B, D, F).




Figure 2. Semi-quantitative RT-PCR analysis of Japanese flounder IFN- γ in infected tissues (spleen, head kidney, liver). A, C, E: spleen, head kidney, liver of Japanese flounder injected with LPS, respectively; B, D, F: spleen, head kidney, liver of Japanese flounder injected with PS, respectively; G: the relative IOD of IFN γ versus GAPDH in infected spleen, head kidney and liver; M: molecular weight standard; CK: uninfected tissues as controls; Different time points after challenge with LPS are indicated on the above margin. The Japanese flounder IFN- γ gene-specific RT-PCR products (371 bp) and GAPDH RT-PCR products (577 bp) are indicated on the right margin.

Mx expression in infected Japanese flounder tissues

The expression of Mx in spleen, head kidney and liver was dramatically induced after challenge. The expression in spleen started at 9h and peaked at 36h and then decreased to background level at 60h. The expression levels of Mx in spleen from 9h to 48h were approximately 1.7-, 2-, 2.1-, 2.5- and 1.3-fold compared with CK (Fig.3 A, G). The highest expression level in head kidney was detected at 36h and went back to background level at 60h. The expression levels were approximately 5.1-, 5.1-, 7.9-, 8.6- and 4.3-fold compared with CK from 9h to 48h (Fig.3 C, G). Seen from Figure.3 E, the expression of Mx in liver was high at 36h, peaked at 48h and decreased at 60h and the expression of Mx at other time points was very faint. Compared with CK, the expression level peaked as high as 15-fold in liver at 48h (Fig.3 G). The control group also showed that the injection of PS did not influence the expression of Mx (Fig.3 B, D, F).




Figure 3. Semi-quantitative RT-PCR analysis of Japanese flounder Mx in infected tissues (spleen, head kidney, liver). A, C, E: spleen, head kidney, liver from the Japanese flounder injected with LPS, respectively; B, D, F: spleen, head kidney, liver from the Japanese flounder injected with PS, respectively; G: the relative IOD of Mx versus GAPDH in infected spleen, head kidney and liver; M: molecular weight standard; CK: uninfected tissues as controls; Different time points after the challenge with LPS are indicated on the above margin. The Japanese flounder Mx gene-specific RT-PCR products (395 bp) and GAPDH RT-PCR products (577 bp) are indicated on the right margin.

IRF-1 expression in infected Japanese flounder tissues

The highest expression levels of the Japanese flounder IRF-1 was detected at 12h in spleen and liver and at 24h in head kidney after infection. The expression in spleen decreased gradually to background level after 48h and the expression in head kidney had a similar trend after 60h (Fig. 4 A, C). As shown in Fig.4 G, the highest expression level in spleen at 12h was approximately 2.1-fold and in head kidney at 24h was approximately 3.3-fold compared with CK. The expression of IRF-1 in liver was not as obvious as in spleen and head kidney and the expression was very faint at most time points except at 12h. The expression of IRF-1 in liver was rapidly induced at 12h but decreased gradually to background after 60h. From 9h to 60h, the relative IOD value compared with CK was approximately 6- to 16-fold (Fig.4 E, G). The

expression of IRF-1 in spleen, head kidney and liver injected with PS was consistent with the expression of IFN- γ and Mx injected with PS (Fig.4 B, D, F).




Figure 4. Semi-quantitative RT-PCR analysis of Japanese flounder IRF-1 in infected tissues (spleen, head kidney, liver). A, C, E: spleen, head kidney, liver from the Japanese flounder injected with LPS, respectively; B, D, F: spleen, head kidney, liver from the Japanese flounder injected with PS, respectively; G: the relative IOD of IRF-1 versus GAPDH in infected spleen, head kidney and liver; M: molecular weight standard; CK: uninfected tissues as controls; Different time points after the challenge with LPS are indicated on the above margin. The Japanese flounder IRF-1 gene-specific RT-PCR products (303 bp) and GAPDH RT-PCR products (577 bp) are indicated on the right margin.

Expression of IFN- γ , Mx and IRF-1 in spleen, head kidney and liver at 12h, 24h, 36h and 48h

According to the results, the main expression time points of IFN- γ , Mx and IRF-1 were at 12h, 24h, 36h and 48h. So the four time points were selected to analyze the expression of the three genes in spleen, head kidney and liver.

The expression levels of IFN- γ , Mx and IRF-1 in liver was very faint compared with the expression levels in spleen and head kidney. The expression of IFN- γ in spleen was the same as in head kidney at 12h and 24h but became weaker than head kidney at 36h and 48h (Fig.5 A). The expression of Mx in spleen and head kidney was both high at 36h and the expression difference at the four time points was not significant (Fig.5 B). In spleen, the expression of IRF-1 was higher than head kidney at 12h and 36h, consistent with head kidney at 24h and 48h (Fig.5 C).

Expression analysis of three immune genes




Figure 5. Semi-quantitative RT-PCR analysis of IFN- γ , Mx and IRF-1 in infected tissues of Japanese flounder at 12h, 24h, 36h, 48h. A, B, C: Expression of Japanese flounder IFN- γ , Mx, IRF-1 in infected tissues (spleen, head kidney, liver) at different time points (12h, 24h, 36h, 48h). M: molecular weight standard; a-d: 12h, 24h, 36h, 48h. 1-3: spleen, head kidney, liver. D: the relative IOD of IFN- γ , Mx and IRF-1 versus GAPDH; The Japanese flounder IFN- γ gene-specific RT-PCR products (371bp), Mx gene-specific RT-PCR products (395bp) and IRF-1 gene-specific RT-PCR products (303bp) are indicated on the right margin, along with GAPDH RT-PCR products (577bp) as an internal control.

DISCUSSION

The innate immune system is primarily important to combat infections in fish. It is commonly stated that the innate immune system are relatively temperature independent because of its evolutionary status and poikilothermic nature^{3, 4, 5}. Interferon, Mx and IRF play important roles in immune response in fish. It was considered that innate parameters were active at the lower temperature range of the fish. It was proved that in cod the spontaneous haemolytic and antiprotease activity was more active at 1°C and 7°C than at 14°C³⁸. However, there were some examples of low temperature having adverse effects on innate parameters. Rainbow trout acclimatized at different temperatures showed lower phagocytic and complement activity at 5°C than at 20°C³⁹. Similarly, a drop in water temperature from 22°C to 10°C had adverse effects on the protective role of channel catfish mucus⁴⁰. In this study the Japanese flounders were kept for one week with lower water temperature at 8°C-10°C before the experiment was carried out. The results showed that the expression of IFN- γ , Mx and IRF-1 in spleen, head kidney and liver of Japanese flounders injected with LPS was different and regular at thirteen time points, which indicated the innate system of Japanese flounder could resist external infection well at lower temperature. Our results showed that the three immune-related genes were differently expressed in Japanese flounder tissues, which indicated the sites of function in the body. There were high expression levels of IFN- γ , Mx and IRF-1 in classical fish immune organs like spleen and head kidney (Fig. 1), which suggested their functional roles in the immune system. The expression of three genes was very intense in blood, which showed that abundant cytokines existed in blood that transcribes these genes. When organism was invaded by virus or bacteria, cytokines in blood could perform their function in time. Besides, the expression of IFN- γ and IRF-1 were also detected in muscle tissue of Japanese flounder but Mx was not detected. Interestingly, the expression of IFN- γ was also detected in goldfish muscle⁴¹, which was consistent with our results. This means that some immune factors

existed not only in immune organs but also in the organs which did not take part in immune response.

IFNs are best known for their vital roles in innate immune system. Recently, studies focused on the immune-related functions of IFN- γ had been carried out in many teleosts, such as goldfish⁴² and zebrafish¹⁰. We have carried out the expression analysis of the Japanese flounder IFN- γ in tissues infected with LPS and revealed the expression regularity. The Japanese flounder IFN- γ was highly induced in spleen at 24h, in head kidney at 36h and in liver at 48h (Fig. 2). The results were very similar to the expression of type-I IFN in head kidney of Atlantic Salmon⁴³, which suggested the Japanese flounder IFN- γ played important roles in immune responses to external invasion.

Mx induced by type-I IFNs (IFN- α and IFN- β) is a kind of anti-viral protein which specially prevents growth, in vivo and in vitro, of certain classes of virus. Up to date, many experiments have been carried out to study the immune-related functions of Mx in many teleosts. It was reported that Rainbow trout Mx1, Mx2 and Mx3 showed no significant inhibition against a fish rhabdovirus, infectious hematopoietic necrosis virus (IHNV)⁴⁴. The expression analysis of Japanese flounder Mx in infected tissues was carried out in the study. The Mx was highly expressed in spleen, head kidney at 36h and in liver at 48h after challenge with LPS (Fig. 3). The expression of Mx after challenge was instant, steady and abundant; therefore it could serve as a molecular indicator to the expression of type-I IFNs in Japanese flounder.

IRFs are best known for their vital roles in regulating the expression of IFN-I, ISGs and other cytokines^{24, 25}. Recently, studies on IRFs in teleosts also focused on their immune-related functions. We carried out the expression analysis of the Japanese flounder IRF-1 in infected tissues. Interestingly, the Japanese flounder IRF-1 was highly expressed at 12h in spleen, liver and at 24h in head kidney (Fig. 4). The result was very similar to the expression of IRF-1 in spleen of large yellow croaker⁴⁵. However, it was very different from the results of the relative expression of Japanese flounder IRF-1 infected with HRV and *Edwardsiella tarda* in 1998³⁰, which might be caused by the difference of fish, seawater temperature and area. The expression of IRF-1 gene may reflect its importance in acute stresses in Japanese flounders.

The expression of IRF-1 was found to be instant which could increase and peak in one day. It has been demonstrated that IRF-1 could recognize and bind to ISREs which were found in Mx promoters. IRF-1 could induce the expression of type-I IFN which could induce the expression of Mx, so the expression of IRF-1 should be earlier than Mx theoretically. Our results showed that the highest expression of IRF-1 was 24h earlier than Mx in spleen, 12h earlier than Mx in head kidney and 36h earlier than Mx in liver, which was consistent with the expression sequence of IRF-1 and Mx. We also found that the highest expression of IFN- γ and Mx were both at 36h in head kidney, 48h in liver. It indicated that the expression of Mx was not induced by IFN- γ , the expression patterns of the two genes were relatively consistent in time. We also studied the expression level of the three genes in spleen, head kidney and liver at different time points (12h, 24h, 36h and 48h). The results showed that the expression of the three genes in different organs changed with time after challenge and the expression level in liver was lower than in spleen and head kidney which meant that the immune response in liver was weaker than in spleen and head kidney after infection. For example, the expression of IFN- γ in spleen was basic equal with in head kidney at 12h and 24h but became weaker at 36h and 48h, which indicated the activity of IFN- γ in head kidney was much stronger. The expression of Japanese flounder Mx in spleen decreased earlier than in head kidney. However, the expression level of IRF-1 in spleen was higher than in head kidney, which suggested that the expression and activity of immune-related gene differed in tissues.

CONCLUSION

In conclusion, the highest expression level of the IFN- γ gene was detected at 24h in spleen, 36h in head kidney and 48h in liver after normal adult Japanese flounder challenged with lipopolysaccharide. In the same manner, the highest expression levels of Mx gene in spleen and head kidney were both at 36h and IRF-1 gene in spleen and liver were both at 24h. And the highest expression level of Mx gene in liver was at 48h and IRF-1 gene in head kidney was at 12h. This study provides a basis for further research on immune mechanism of IFN- γ , Mx, IRF-1 and the production of recombinant IFN- γ , recombinant Mx or recombinant IRF-1 used in Japanese flounder cultivation in future.

ACKNOWLEDGMENTS

This study was supported by the grant of Dalian Ocean University (No. 2012HYDX07). We thank John Wheaton (College of International Exchange and English Education, Jishou University) for polishing the manuscript.

REFERENCES

1. Fearon D.T. & Locksley R.M. The instructive role of innate immunity in the acquired immune response. *Science*. 1996; 272: 50-53.
2. Fearon D.T. Seeking wisdom in innate immunity. *Nature*. 1997; 388: 323-324.
3. Magnadóttir B. Innate immunity of fish (overview). *Fish Shellfish Immunol*. 2006; 20: 137-151.
4. Du Pasquier L. Antibody diversity in lower vertebrates--why is it so restricted? *Nature*. 1982;296: 311-313.
5. Ellis A.E. Innate host defence mechanism of fish against viruses and bacteria. *Dev. Comp. Immunol*.2001;25: 827-839.
6. Rombout J.H, Taverne N, van de Kamp M, Taverne-Thiele A.J. Differences in mucus and serum immunoglobulin of carp (*Cyprinus carpio* L.).*Dev. Comp. Immunol*. 1993; 17: 309-317.
7. Stafford J.L, Belosevic M. Transferrin and the innate immune response of fish: identification of a novel mechanism of macrophage activation. *Dev. Comp. Immunol*. 2003; 27: 539-554.
8. Muller U, Steinhoff U, Reis L.F, Hemmi S, Pavlovic J, Zinkernagel R.M, et al. Functional role of type I and type II interferons in antiviral defense. *Science*. 1994; 264: 1918-1921.
9. Roberts R.M, Liu L, Guo Q, Leaman D, Bixby J. The evolution of the type I interferons. *J. Interferon Cytokine Res*. 1998; 18: 805-816.
10. Sieger D, Stein C, Neifer D, van der Sar A.M, Leptin M. The role of gamma interferon in innate immunity in the zebrafish embryo. *Dis. Model Mech*. 2009; 2: 571-581.
11. Wheelock E.F. Interferon-like virus-inhibitor induced in human leukocytes by phytohemagglutinin. *Science*. 1965; 149: 310-311.
12. Wang Z.E, Reiner S.L, Zheng S, Dalton D.K, Locksley R.M. CD4⁺ effector cells default to the Th2 pathway in interferon gamma-deficient mice infected with *Leishmania major*. *J. Exp. Med*. 1994; 179: 1367-1371.
13. Huang S, Hendriks W, Althage A, Hemmi S, Bluethmann H, Kamijo R, et al. Immune response in mice that lack the interferon-gamma receptor. *Science*. 1993; 259: 1742-1745.
14. Cooper A.M, Dalton D.K, Stewart T.A, Griffin J.P, Russell D.G, Orme I.M. Disseminated tuberculosis in interferon gamma gene-disrupted mice. *J. Exp. Med*. 1993; 178: 2243-2247.
15. Staeheli P, Yu Y.S, Grob R, Haller O. A double stranded-RNA inducible fish gene homologous to the murine influenza virus resistance gene Mx. *Mol. Cell Biol*. 1989; 9: 3117-3121.

16. Altmann S.M, Mellon M.T, Johnson M.C, Paw B.H, Trede N.S, Zon L.I, et al. Cloning and characterization of an Mx gene and its corresponding promoter from the zebrafish, *Danio rerio*. *Dev. Comp. Immunol.* 2004;28: 295-306.
17. Trobridge G.D, Chiou P.P, Leong J.A. Cloning of the rainbow trout (*Oncorhynchus mykiss*) Mx2 and Mx3 cDNAs and characterization of trout Mx protein expression in salmon cells. *J. Virol.* 1997; 71: 5304-5311.
18. Zhang Y.B, Li Q, Gui J.F. Differential expression of two *Carassius auratus* Mx genes in cultured CAB cells induced by grass carp hemorrhage virus and interferon. *Immunogenetics.* 2004; 56: 68-75.
19. Lee J.Y, Hirono I, Aoki T. Cloning and analysis of expression of Mx cDNA in Japanese flounder, *Paralichthys olivaceus*. *Dev. Comp. Immunol.* 2000; 24: 407-415.
20. Yap W.H, Tay A, Brenner S, Venkatesh B. Molecular cloning of the pufferfish (*Takifugu rubripes*) Mx gene and functional characterization of its promoter. *Immunogenetics.* 2003; 54: 705-713.
21. Caipang C.M.A, Hirono I, Aoki T. In vitro inhibition of fish rhabdoviruses by Japanese flounder. *Paralichthys olivaceus* Mx, *Virology.* 2003; 317:373-382.
22. Larsen R, Rokenes T.P, Robertsen B. Inhibition of infectious pancreatic necrosis virus replication by atlantic salmon Mx1 protein. *J. Virol.* 2004; 78: 7938-7944.
23. Pakingking R. Jr, Okinaka Y, Mori K, Arimoto M, Muroga K, Nakai T. In vivo and in vitro analysis of the resistance against viral haemorrhagic septicaemia virus in Japanese flounder (*Paralichthys olivaceus*) preceedingly infected with aquabirnavirus. *Fish Shellfish Immunol.* 2004; 17: 1-11.
24. Mamane Y, Heylbroeck C, Génin P, Algarté M, Servant M.J, LePage C, et al. Interferon regulatory factors: the next generation. *Gene.* 1999; 237: 1-14.
25. Taniguchi T, Ogasawara K, Takaoka A, Tanaka N. IRF family of transcription factors as regulators of host defense. *Annu. Rev. Immunol.* 2001; 19:23-55.
26. Tamura T, Yanai H, Savitsky D, Taniguchi T. The IRF family transcription factors in immunity and oncogenesis. *Annu. Rev. Immunol.* 2008; 26: 535-584.
27. Escalante C.R, Yie J, Thanos D, Aggarwal A.K. Structure of IRF-1 with bound DNA reveals determinants of interferon regulation. *Nature.* 1998; 391: 103-106.
28. Barnes B, Lubyova B, Pitha P.M. On the role of IRF in host defense. *J. Interferon Cytokine Res.* 2002; 22: 59-71.
29. Nguyen H, Hiscott J, Pitha P.M. The growing family of interferon regulatory factors. *Cytokine Growth Factor Rev.* 1997;8: 293-312.
30. Yabu T, Hirose H, Hirono I, Katagiri T, Aoki T, Yamamoto E. Molecular cloning of a novel interferon regulatory factor in Japanese flounder *Paralichthys olivaceus*. *Mol. Mar. Biol. Biotechnol.* 1998; 7: 138-144.
31. Collet B, Hovens G.C, Mazzoni D, Hirono I, Aoki T, Secombes C.J. Cloning and expression analysis of rainbow trout *Oncorhynchus mykiss* interferon regulatory factor 1 and 2 (IRF-1 and IRF-2). *Dev. Comp. Immunol.* 2003; 27: 111-126.
32. Ordás M.C, Abollo E, Costa M.M, Figueras A, Novoa B. Molecular cloning and expression analysis of interferon regulatory factor-1 (IRF-1) of turbot and sea bream. *Mol. Immunol.* 2006; 43: 882-890.
33. Sun B.J, Chang M.X, Song Y, Yao W.J, Nie P. Gene structure and transcription of IRF-1 and IRF-7 in the mandarin fish *Siniperca chuatsi*. *Vet. Immunol. Immunopathol.* 2007; 116: 26-36.
34. Lee E.Y, Park H. H, Kim Y. T, Choi T. J. Cloning and sequence analysis of the interleukin-8 gene from flounder (*Paralichthys olivaceus*). *Gene.* 2001; 274: 237-243.
35. Liu Y, Chen S.L, Meng L, Zhang Y.X. Cloning, characterization and expression analysis of a novel CXC chemokine from turbot (*Scophthalmus maximus*). *Fish Shellfish Immunol.* 2007;23: 711-720.
36. Hayashi H, Nishimoto A, Oshima N, Iwamuro S. Expression of the estrogen receptor alpha gene in the anal fin of Japanese medaka, *Oryzias latipes*, by environmental concentrations of bisphenol A. *J. Toxicol Sci.* 2007; 32: 91-96.
37. Park D.Y, Lee C.H, Sol M.Y, Suh K.S, Yoon S.Y, Kim J.W. Expression and localization of the transforming growth factor-beta type I receptor and Smads in preneoplastic lesions during chemical hepatocarcinogenesis in rats. *J. Korean Med. Sci.* 2003; 18: 510-519.

38. Magnadóttir B, Jónsdóttir H, Helgason S, Björnsson B, Jørgensen T.O, Pilström L. Humoral immune parameters in Atlantic cod (*Gadus morhua* L.) I: The effects of environmental temperature. *Comp. Biochem Physiol B*. 1999; 122: 173-180.
39. Nikoskelainen S, Bylund G, Lilius E.M. Effect of environmental temperature on rainbow trout (*Oncorhynchus mykiss*) innate immunity. *Dev. Comp. Immunol*. 2004; 28: 581-592.
40. Quiniou S.M.A, Bigler S, Clem L.W, Bly J.E. Effects of water temperature on mucous cell distribution in channel catfish epidermis: a factor in winter saprolegniasis. *Fish Shellfish Immunol*. 1998; 8: 1-11.
41. Grayfer L. & Belosevic M. Molecular characterization, expression and functional analysis of goldfish (*Carassius auratus* L.) interferon gamma. *Dev. Comp. Immunol*. 2009; 33: 235-246.
42. Grayfer L, Belosevic M. Molecular characterization, expression and functional analysis of goldfish (*Carassius auratus* L.) interferon gamma. *Dev. Comp. Immunol*. 2009; 33: 235-246.
43. Robertsen B, Bergan V, Rokenes T, Larsen R, Albuquerque A. Atlantic salmon interferon genes: cloning, sequence analysis, expression, and biological activity. *J. Interferon Cytokine Res*. 2003; 23: 601-612.
44. Trobridge G.D, Chiou P.P, Leong J.A. Cloning of the rainbow trout (*Oncorhynchus mykiss*) Mx2 and Mx3 cDNAs and characterization of trout Mx protein expression in salmon cells. *J. Virol*. 1997; 71: 5304-5311.
45. Yao C.L, Kong P, Huang X.N, Wang Z.Y. Molecular cloning and expression of IRF1 in large yellow croaker, *Pseudosciaena crocea*. *Fish Shellfish Immunol*. 2010; 28: 654-660.

Received: February 03, 2016;

Accepted: July 14, 2016