



# The Brazilian Journal of INFECTIOUS DISEASES

www.elsevier.com/locate/bjid



## Brief communication

# Emergence of *Klebsiella pneumoniae*-producing KPC-2 carbapenemase in Paraíba, Northeastern Brazil

Lorena C.C. Fehlberg<sup>a,\*</sup>, Albalucia M.C. Carvalho<sup>b</sup>, Eloiza H. Campana<sup>a</sup>,  
Paulo P. Gontijo-Filho<sup>c</sup>, Ana C. Gales<sup>a</sup>

<sup>a</sup> Laboratório Alerta, Division of Infectious Diseases, Universidade Federal de São Paulo, São Paulo, SP, Brazil

<sup>b</sup> Hospital Universitário Lauro Wanderley, Universidade Federal da Paraíba, João Pessoa, PB, Brazil

<sup>c</sup> Biological Sciences Institute, Universidade Federal de Uberlândia, Uberlândia, MG, Brazil

## ARTICLE INFO

### Article history:

Received 14 February 2012

Accepted 19 July 2012

Available online 14 November 2012

### Keywords:

Beta-lactam resistance

Modified Hodge test

Carbapenem resistance

Enterobacteriaceae

## ABSTRACT

The emergence of KPC-2 producing *K. pneumoniae* in hospitalized patients at the intensive care unit (ICU) of a teaching hospital located in the city of João Pessoa, Paraíba, Brazil, is reported. Seven carbapenem-resistant *K. pneumoniae* recovered from different body sites of infection were analyzed. Most isolates showed a multidrug-resistance phenotype. Genotypic analysis demonstrated the presence of two genotypes, with the predominance of genotype A, which belongs to ST 437. These isolates also carry the encoding genes of five other beta-lactamases.

© 2012 Elsevier Editora Ltda. All rights reserved.

*Klebsiella pneumoniae* carbapenemases (KPC)-producing Enterobacteriaceae isolates have become a significant problem worldwide. KPC are enzymes that hydrolyze penicillins, cephalosporins, monobactams, and carbapenems, and are inhibited by clavulanic acid and tazobactam.<sup>1</sup> Of notice, genes encoding KPC enzymes are frequently located on transferable plasmids. KPC-producing clinical isolates have been associated with hospital outbreaks in the USA, Asia, Europe, South Korea, and Central and South America.<sup>2</sup> In Brazil, KPC-2-producing isolates have been initially reported in the cities of Recife, Rio de Janeiro, São Paulo, and Porto Alegre.<sup>3-9</sup> In this report, the emergence of KPC-producing-*K. pneumoniae* (KPC-KPN) clinical isolates recovered at the tertiary teaching

Hospital Lauro Wanderley, located in João Pessoa, Paraíba, Brazil is described.

In the period between 2009 and 2010, seven *K. pneumoniae* isolates were collected from different body sites of infection of five patients who were hospitalized at the intensive care unit (ICU) of the Hospital Lauro Wanderley. Initially, these isolates were identified as resistant to ertapenem by disc diffusion method, and were referred to the Laboratório Alerta, UNIFESP, São Paulo, Brazil for further characterization. Susceptibility testing was performed by the agar dilution method, according to the Clinical and Laboratory Standards Institute's (CLSI) guidelines<sup>10</sup> for amikacin, aztreonam, cefepime, cefoxitin, ceftazidime, ceftriaxone, ciprofloxacin, ertapenem,

\* Corresponding author at: Laboratório Alerta – Division of Infectious Diseases, Universidade Federal de São Paulo, Rua Leandro Dupret, 188, Vila Clementino – São Paulo/SP, 04025-010, Brazil.

E-mail address: [lorenafehlberg@gmail.com](mailto:lorenafehlberg@gmail.com) (L.C.C. Fehlberg).

1413-8670/\$ – see front matter © 2012 Elsevier Editora Ltda. All rights reserved.

<http://dx.doi.org/10.1016/j.bjid.2012.07.001>

**Table 1 – Clinical features of the patients who had *K. pneumoniae* KPC-2-producing isolates at Hospital University Lauro Wanderley, João Pessoa, Paraíba, Brazil.**

Patient	Isolate	PFGE pattern	ST	Gender	Age (y)	Underlying disease	Primary infection	Treatment	Antimicrobial resistant	MIC carbapenems (CLSI 2010) <sup>*</sup>		Bacteria isolation (day/mo/yr)	Beta-lactamases	ICU admission (day/mo/yr)	Death (day/mo/yr)
										IMI	MEM				
1	335	A1	437	F	52	Bowel cancer	Blood-stream	CS, MEM, AMG	ERT, CAZ, CRO, CFO, CIP, ATM	1 (S)	1 (S)	20/02/2010	KPC-2; TEM-1; OXA-1; CTX-M-15; SHV-11	19/02/2010	22/02/2010
2	336	A1	437	F	71	Diabetes	Blood-stream	CAZ, CLI, CIP, MEM	ERT, CAZ, CRO, CFO, CIP, ATM	2 (I)	2 (I)	09/09/2009	pc KPC-2; TEM-1; OXA-1; CTX-M-15; SHV-11	03/09/2009	09/09/2009
	339	A1	437	F	71	Diabetes	Urinary tract		ERT, IMI, CAZ, CRO, CFO CIP, ATM	16 (R)	0,5 (S)				
3	337	B1	70	F	29	HIV	Blood-stream	CS, MEM, AMG, FUN	ERT, CRO, CIP, ATM	2 (I)	1 (S)	13/02/2009	KPC-2; SHV-27	21/02/2009	16/03/2009
	338	B1	70	F	29	HIV	Pneumonia		ERT, CAZ, CRO, CFO, ATM, MEM, IMI	8 (R)	64 (R)	20/02/2009			
4	340	A1	437	M	65	Encephalopathy	Blood-stream	CS, MEM, AMG, VAN	ERT, CAZ, IMI, CRO, CFO, CIP, ATM	4 (R)	2 (I)	08/08/2009	KPC-2; TEM-1; OXA-1; CTX-M-15; SHV-11	18/08/2009	25/10/2009
5	341	A1	437	F	52	Post gastrectomy	Wound	MEM	ERT, CAZ, IMI, CRO, CFO, CIP, ATM	4 (R)	1 (S)	17/08/2009	KPC-2; TEM-1; OXA-1; CTX-M-15; SHV-11	12/08/2009	18/08/2009

AMG, aminoglycosides; ATM, aztreonam; CAZ, ceftazidime; CFO, cefoxitin; CIP, ciprofloxacin; CLI, clindamycin; CRO, ceftriaxone; CS, 3th and 4th cephalosporins; ERT, ertapenem; F, female; FUN, antifungal; HIV, human immunodeficiency virus infection; I, susceptibility reduced; ICU, intensive care unit; IMI, imipenem; M, male; MEM, meropenem; R, resistant; S, susceptible; ST, sequence type; VAN, vancomycin.

\* MICs were determined by agar dilution.

imipenem, meropenem, and piperacillin/tazobactam. Minimum inhibitory concentrations (MICs) for tigecycline and polymyxin B were determined by Etest (bioMérieux – Marcy l'Étoile, France) and interpreted according to the European Committee on Antimicrobial Susceptibility Testing's (EUCAST) guidelines.<sup>11</sup> The Modified Hodge Test (MHT) with ertapenem disk (10 µg) was used for phenotypic detection of carbapenemase activity.<sup>12</sup> Specific primers were used under standard polymerase chain reaction (PCR) conditions to detect the genes *bla*<sub>KPC</sub>, *bla*<sub>GES</sub>, *bla*<sub>CTX-M</sub>, *bla*<sub>SHV</sub>, *bla*<sub>TEM</sub>, *bla*<sub>OXA-1</sub>, *bla*<sub>OXA-2</sub>, *bla*<sub>OXA-10</sub>, *bla*<sub>OXA-18/45</sub>, and *bla*<sub>OXA-161</sub>, followed by DNA sequencing (ABI sequencer - Applied Biosystems, Foster City, CA). Clonal relatedness among isolates was examined by pulsed field gel electrophoresis (PFGE) using SpeI (New England, Beverly, MA). The band patterns were analyzed by visual interpretation, applying the criteria established by Tenover et al.<sup>13</sup> In addition, the PFGE patterns were analyzed using Bionumerics version 5.10 (Applied Maths, Sint-Martens – Latem, Belgium) for comparison with other KPC-2-producing strains previously identified in Brazil.<sup>3</sup> Multilocus sequence typing (MLST) of *K. pneumoniae* was performed as described previously.<sup>14</sup> Experimentally determined DNA sequences were uploaded into the MLST database (<http://www.pasteur.fr/recherche/genopole/PF8/mlst/Kpneumoniae.html>), and allelic numbers and sequence types (ST) were obtained.

*K. pneumoniae* isolates were more frequently collected from bloodstream infections (n = 4; 57.1%) (Table 1). Amikacin, polymyxin B, and tigecycline showed the highest susceptibility rates (100%) against the isolates studied, followed by meropenem (57.1%), imipenem (14.3%), and cefepime (14.3%). In contrast, 100% of the isolates were resistant to ertapenem, ceftriaxone, and aztreonam. Resistance to ceftazidime, cefoxitin, and ciprofloxacin was observed in six isolates (87.5%). All isolates showed positive MHT results and were found to carry the *bla*<sub>KPC-2</sub> gene. Two PFGE patterns, A and B, were found among the seven KPC-KPN; the genotype A was the most frequent (n = 5; 71.3%). Besides the presence of *bla*<sub>KPC-2</sub>, genotype A (ST 437) isolates also carried *bla*<sub>TEM-1</sub>, *bla*<sub>OXA-1</sub>, *bla*<sub>CTX-M-15</sub>, and *bla*<sub>SHV-11</sub>, while genotype B (ST 70) carried only *bla*<sub>SHV-27</sub>. Using the dendrogram, a similarity of nearly 79.3% was detected between genotype A and B. The similarity of the KPC-producing isolates evaluated in this study compared to those previously isolated in Recife<sup>3</sup> was 74.5%.

For the first time, *K. pneumoniae* isolates resistant to ertapenem were detected in the Hospital Lauro Wanderley, which were further confirmed as KPC producers. Among the isolates studied, one (335) showed susceptibility to meropenem and imipenem, whereas three isolates (337, 339, and 341) were only susceptible to meropenem, according to the CLSI breakpoints.<sup>10</sup> The discrepancy observed in the susceptible categories to carbapenems may be attributed to production of carbapenemases like KPC. The co-production of KPC with other beta-lactamases, such as TEM-1, OXA-1, SHV-11, SHV-27 and CTX-M-15, was observed among KPC-KPN isolates. These findings are in agreement with previous Brazilian reports which showed that KPC-KPN isolates also possessed the *bla*<sub>CTX-M</sub>, *bla*<sub>TEM</sub>, and *bla*<sub>SHV</sub> genes.<sup>3,5,15</sup> Although CTX-M-2 is a widespread variant in South America, the isolates evaluated carried *bla*<sub>CTX-M-15</sub>, a frequently found worldwide CTX-M-type.

Additionally, according to the authors' knowledge, this is the first description of co-production of KPC-2 and OXA-1 in Brazilian isolates. ST 437, the most prevalent among the KPC-KPN evaluated, is a ST related to the clonal complex 258, which is widely disseminated among KPC-KPN in Brazil, and associated with the dissemination of KPC worldwide.

Since the therapeutic options are limited and the appropriate empirical antimicrobial treatment is of crucial importance to patient outcome,<sup>16</sup> we suggest that the clinical laboratory perform accurate susceptibility testing for KPC producers, including the determination of MICs for tigecycline, aminoglycosides, polymyxins, and carbapenems, since KPC-KPN isolates with low carbapenems MICs and/or isolates with discrepancy for susceptible category among carbapenems tested were observed. In addition, regional surveillance studies that monitor the dissemination of ESBL and carbapenemase enzymes are crucial, since most of these genes are located on mobile genetic elements, which are easily transferred to other bacteria species. Because four of the five isolates were clonally related, suggesting a patient-to-patient transmission, implementation of infection control measures is necessary to restrain the dissemination of resistant genes.

## Conflict of interest

All authors declare to have no conflict of interest.

## Acknowledgements

The authors would like to thank Vinícius G. S. Oliveira and Adriana G. Nicoletti for performing the PFGE and MLST assays, respectively, and Ana Carolina R. Silva for her technical assistance with the PCR and sequencing.

Part of this study was presented at the II International Symposium of Clinical Microbiology, in Florianópolis, Brazil, September 29 to October 2, 2010.

## REFERENCES

1. Queenan AM, Bush K. Carbapenemases: the versatile beta-lactamases. *Clin Microbiol Rev.* 2007;20:440-58.
2. Bush K. Alarming beta-lactamase-mediated resistance in multidrug-resistant *Enterobacteriaceae*. *Curr Opin Microbiol.* 2010;13:558-64.
3. Monteiro J, Santos AF, Asensi MD, et al. First report of KPC-2 producing *Klebsiella pneumoniae* strains in Brazil. *Antimicrob Agents Chemother.* 2009;53:333-4.
4. Pavez M, Mamizuka EM, Lincopan N. Early dissemination of KPC-2-producing *Klebsiella pneumoniae* strains in Brazil. *Antimicrob Agents Chemother.* 2009;53:2702.
5. Peirano G, Seki LM, Passos VLV, et al. Carbapenem-hydrolyzing beta-lactamase KPC-2 in *Klebsiella pneumoniae* isolated in Rio de Janeiro. *Brazil J Antimicrob Chemother.* 2009;63:265-8.
6. Zavascki AP, Machado ABMP, Oliveira KRP, et al. KPC-2 producing *Enterobacter cloacae* in two cities from Southern Brazil. *Int J Antimicrob Agents.* 2009;34:281-91.
7. D'Alincourt Carvalho-Assef AP, Leão RS, da Silva RV, et al. *Escherichia coli* producing KPC-2 carbapenemase: first report in Brazil. *Diagn Microbiol Infect Dis.* 2010;68:337-8.

8. Leão RS, Pereira RH, Folescu TW, et al. KPC-2 Carbapenemase-producing *Klebsiella pneumoniae* isolates from patients with Cystic Fibrosis. *J Cyst Fibros*. 2011;10:140-2.
9. Almeida ACS, Vilela MA, Cavalcanti FL, et al. First description of KPC-2-producing *Pseudomonas putida* in Brazil. *Antimicrob Agents Chemother*. 2012;56:2205-6.
10. Clinical and Laboratory Standards Institute. Performance standards for antimicrobial susceptibility testing. 22th Informational Supplement Update. Clinical and Laboratory Standards (M100-S22): Wayne, PA; 2012.
11. EUCAST. Breakpoint tables for interpretation of MICs and zone diameters. Version 1.1, January 2012. Available from: [http://www.eucast.org/clinical\\_breakpoints/](http://www.eucast.org/clinical_breakpoints/)
12. Lee K, Chong Y, Shin HB, et al. Modified Hodge and ETDA-disk synergy testes to screen metallo-beta-lactamase-producing strains of *Pseudomonas aeruginosa* and *Acinetobacter* species. *Clin Microbiol Infect*. 2001;7:88-91.
13. Tenover FC, Arbeit RD, Goering RV, et al. Interpreting chromosomal DNA restriction patterns produced by pulsed-field gel electrophoresis: criteria for bacterial strain typing. *J Clin Microbiol*. 1995;33:2233-9.
14. Diancourt L, Passet V, Verhoef J, et al. Multilocus sequence typing of *Klebsiella pneumoniae* nosocomial isolates. *J Clin Microb*. 2005;43:4178-82.
15. Seki LM, Pereira PS, Souza MPAH, et al. Molecular epidemiology of KPC-2-producing *Klebsiella pneumoniae* isolates in Brazil: the predominance of sequence type 437. *Diag Microbiol Infect Dis*. 2011;70:274-7.
16. Rice LB. Emerging issues in the management of infections caused by multidrug-resistant Gram-negative bacteria. *Cleve Clin J Med*. 2007;74:S12-20.