



The Brazilian Journal of INFECTIOUS DISEASES

www.elsevier.com/locate/bjid



Letter to the Editor

Colistin resistance in non-fermenting Gram-negative bacilli in a university hospital

Dear Editor,

Indiscriminate use of antimicrobials has caused the worldwide emergence of multidrug resistant bacteria (MDR). Glucose non-fermenting Gram-negative bacilli (NF-GNB) that show resistance to several antimicrobials, including colistin, have been reported. The highest rates of this type of resistance have been registered in Asia and Europe.¹ This fact has created problems in therapy, since the colistin is used as last alternative to treat infections caused by MDR bacteria.² *Pseudomonas aeruginosa* is a skilful microorganism that causes severe opportunistic infections in immunocompromised patients due to its metabolic versatility.³ Isolation of *P. aeruginosa* resistant to almost all available antimicrobials is increasing in Intensive Care Units (ICU), leading to an increase in the number of healthcare-associated infections (HAI), significantly influencing morbidity, mortality, and high costs of treatment of infected patients.¹

In a prospective study developed between August 2015 and January 2016 we isolated 293 NF-GNB of various clinical specimens from patients admitted to a university hospital in the central region of Rio Grande do Sul, Brazil. Identification and antimicrobials sensitivity test were performed through the automated system VITEK®2 (bioMérieux). Data were interpreted according to the document M100-S25 of the Clinical and Laboratory Standards Institute [CLSI, 2015], considering resistant the isolated specimens which showed intermediate or resistant colistin profiles.

From the 293 NF-GNB assessed in this study, only two genera were identified: 78.83% (231) *P. aeruginosa* and 21.16% (62) *Acinetobacter* spp. Colistin resistance was identified in nine isolated strains of *P. aeruginosa* (9/231-3.89%) and none in *Acinetobacter* spp. The predominant age group of patients in whom colistin-resistant strains were identified was 41–69 years old (77.77%); this age-range also showed expressive resistance to carbapenems (66.67%) (Table 1).

Colistin-resistant *P. aeruginosa* were predominantly isolated from clinical specimens obtained from soft tissues (3/9-33.33%) and respiratory (sputum and tracheal secretion – 2/9-22.22%) secretions. The hospital wards in which resistant strains were initially identified were internal medicine

Table 1 – Sensitivity profile of 9 *P. aeruginosa* isolates resistant to colistin.

Antimicrobial	Sensitive (n-%)	Resistant (n-%)
Amikacin	7-77, 78%	2-22, 23%
Gentamicin	5-55, 56%	4-44, 45%
Cefepima	5-55, 56%	4-44, 45%
Ciprofloxacin	4-44, 45%	5-55, 56%
Piperacillin-Tazobactam	3-33, 34%	5-55, 56%
Ampicillin-Sulbactam	0	9-100%
Imipenem	3-33, 34%	6-66, 67%
Meropenem	3-33, 34%	6-66, 67%
Ceftazidime	3-33, 34%	6-66, 67%

(2/9-22.23%), surgery (2/9-22.23%), recovery room (2/9-22.23%), nephrology, adult ER, and ICU (1/9-11.11% each).

From the nine patients in whom colistin-resistant *P. Aeruginosa* was isolated, three progressed to death due to septicemia, four were oncologic patients under anti-neoplastic and radiotherapy regimens, one post-kidney transplant patient, two diabetes mellitus patients, and two chronic smokers/alcoholics. Therefore, we ratify that patients' immunodepression make them vulnerable to infections caused by colistin-resistant microorganisms.

The extensive use of colistin in veterinary medicine to treat gastrointestinal infections caused by Enterobacteriaceae has contributed to chromosomal mutations mediated by the plasmid *mcr-1* to codify resistance to colistin in Gram-negative bacilli. Fernandes and colleagues reported on the evidence of *mcr-1* originated from *Escherichia coli* isolated from food of animal origin in Brazil since 2012.⁴ This silent dissemination of *E. coli* resistance mediated by this plasmid is having an impact in Brazilian ICUs. The emergence and dissemination of MDR bacteria associated with high rates of treatment failure have fostered the use of poliximins, the main therapeutic option for the treatment of MDR infections, especially those caused by *P. aeruginosa*, *Acinetobacter baumannii*, and *Klebsiella pneumoniae*, producers of SPM-1, OXA-23 or KPC-2.⁵ Additional studies are being developed by our research group to identify the gene *mcr-1* in isolated strains that show resistance to colistin.

Conflicts of interest

The authors declare no conflicts of interest.

REFERENCES

1. Cai Y, Chai D, Wang R, Liang B, Bai N. Colistin resistance of *Acinetobacter baumannii*: clinical reports, mechanisms and antimicrobial strategies. *J Antimicrob Chemother*. 2012;22:1–9.
2. Olaitan AO, Morand S, Rolain JM. Emergence of colistin-resistant bacteria in humans without colistin usage: a new worry and cause for vigilance. *Int J Antimicrob Agents*. 2016;47:1–3.
3. Gellatly SL, Hancock REW. *Pseudomonas aeruginosa*: new insights into pathogenesis and host defenses. *Pathog Dis*. 2013;67:159–73.
4. Fernandes MR, Moura Q, Sartori L, et al. Silent dissemination of colistin-resistant *Escherichia coli* in South America could contribute to the global spread of the *mcr-1* gene. *Euro Surveill*. 2016;21:1–6.
5. Rossi F. The challenges of antimicrobial resistance in Brazil. *Clin Infect Dis*. 2011;52:1138–43.

Silvana Oliveira dos Santos^a, Sthefanine Martins La Rocca^b, Rosmari Hörner^{c,*}

^a Universidade Federal de Santa Maria, Programa de Pós-graduação em Ciências Farmacêuticas, Santa Maria, RS, Brazil

^b Universidade Federal de Santa Maria, Santa Maria, RS, Brazil

^c Universidade Federal de Santa Maria, Departamento de Análises Clínicas e Toxicológicas, Santa Maria, RS, Brazil

* Corresponding author.

E-mail address: rosmari.ufsm@gmail.com (R. Hörner).

Received 9 August 2016

Accepted 20 August 2016

1413-8670/© 2016 Sociedade Brasileira de Infectologia.

Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

<http://dx.doi.org/10.1016/j.bjid.2016.08.009>

Available online 15 October 2016