

## Emergence of Resistant *Candida* in Neutropenic Patients

Marcio Nucci and Arnaldo L. Colombo

*Mycology Laboratory, University Hospital,  
Federal University of Rio de Janeiro, and the Division  
of Infectious Diseases, Universidade Federal de São  
Paulo, SP, Brazil*

**Problems with resistance to antifungal drugs have emerged due to an increase in the incidence of systemic fungal infections and widespread use of antifungal agents. Accordingly, efforts have been made to develop adequate fungal susceptibility tests. The ideal test should have high intra and inter-laboratory reproducibility, good correlation with the clinical outcome, and should be easy to perform. While no such test has yet been developed, advances have been made. Over the past decade, many reports of fungal resistance have been published, most of them in AIDS patients. Though the frequency of resistant strains is still low in neutropenic cancer patients, and is mostly limited to *Candida glabrata* and *Candida krusei*, resistance to *Candida albicans* has also been reported.**

**Key Words:** *Candida*, resistance, neutropenic.

The incidence of systemic fungal infections has increased dramatically over the past 20 years [1]. Therapy for such infections has been difficult, because of the limited number of available antifungal agents. After the introduction of the azoles, which have a high oral bioavailability and a low incidence of side effects, a new era in the treatment of fungal infections begun. Indeed, fluconazole has proven highly effective in many clinical situations, such as oral and esophageal candidiasis in AIDS patients [2], systemic fungal infections in bone marrow [3] and liver transplant recipients [4], cryptococcosis [5,6] and candidemia [7,8]. However, in the case of oral candidiasis in AIDS patients, although the efficacy rate is very high, relapse frequently occurs. Therefore, many AIDS patients receive fluconazole for long periods of time. This

scenario has favored the development of resistance, and many reports of treatment failure have been published over the past decade [9-16]. Efforts have been made to standardize antifungal susceptibility tests for fungi. In this review, we will focus on the limitations of the tools for assessing antifungal susceptibility, and the magnitude of this problem in neutropenic cancer patients.

### Antifungal susceptibility tests

Resistance is a concept derived from an *in vitro* phenomenon that, theoretically, is associated with clinical failure, whereas susceptibility would indicate efficacy. However, this is not necessarily true, and in fact, besides *in vitro* antifungal resistance, there are many reasons for treatment failure. For instance, in the treatment of fungal infections in neutropenic patients, bone marrow recovery is a strong predictor of efficacy, regardless of whether the strain is susceptible or resistant [17-19]. Likewise, breakthrough candidemia, an increasingly reported phenomenon, seems not to be associated with resistant strains, but with the compromised immune status of the host [20]. Other

Received on 18 December 2001; revised 22 April 2002.

Address for correspondence: Dr. Marcio Nucci, M.D. Hospital Universitário Clementino Fraga Filho. Av. Brigadeiro Trompovsky s/n. 21941-590 Rio de Janeiro – Brasil. Phone: +5521 25622463. Fax: +5521 25622460. E-mail: mnucci@hucff.ufrj.br

**The Brazilian Journal of Infectious Diseases** 2002;6(3):124-128  
© 2002 by The Brazilian Journal of Infectious Diseases and Contexto Publishing. All rights reserved.  
1413-8670

possible reasons for efficacy failure include infection in non-vascular sites (abscesses, catheters, and prosthesis), poor absorption, and fast elimination or metabolism of the antifungal drug.

Since host factors are often much more important than drug susceptibility, susceptibility does not necessarily predict success. However, susceptibility tests would still be clinically useful if a lack of *in vitro* susceptibility was correlated with treatment failure. In addition, the test should be reproducible. Over the past decade, many methods have been developed, including disk diffusion methods [21], E-test [22], colorimetric methods [23], and broth macro and microdilution methods [24]. The broth macro and microdilution test is the NCCLS (National Committee for Clinical Laboratory Standards) reference method, and provides over 90% intralaboratory and interlaboratory reproducibility [25,26]. Interpretative break points have been established for azoles, based mainly on the analysis of the outcome of oropharyngeal candidiasis in AIDS patients [27]. Therefore, in the case of the azoles, a major limitation for the clinical usefulness of an antifungal susceptibility test is the very limited amount of data on the correlation between MICs and the outcome in invasive candidiasis, especially in neutropenic patients. Likewise, this method remains to be validated for other antifungal agents, including amphotericin B. The correlation between *in vitro* susceptibility tests for amphotericin B and the outcome in candidemia was assessed using the NCCLS method [28]. Among 105 candidemic episodes, 33 had microbiological failure, defined as persistence of *Candida* in the bloodstream after 3 days of amphotericin B treatment. The correlation between MIC and failure was poor. This was mainly because the MIC range was very narrow (0.06 – 2 µg/mL). A resistance breakpoint of =1 µg/mL was suggested.

Although reproducible, the NCCLS standardized method is cumbersome. Thus, alternatives to this method have been sought. Overall, disk diffusion and E-test are effective in identifying susceptible strains. However, these methods do not adequately discriminate

between resistant and intermediate or dose-dependent strains. Therefore, they may be useful as screening tests, but confirmation of resistant strains should be made with the NCCLS method.

### Antifungal resistance in clinical practice

The first reports of antifungal resistance occurred in patients with mucocutaneous candidiasis treated with ketoconazole [29], but since the AIDS epidemic began, this problem has gained great clinical relevance. The typical background for the development of azole resistance is the prolonged and repeated use of fluconazole for the management of oral and esophageal candidiasis in AIDS patients with low CD4+ cell counts [30]. Patients initially infected by susceptible strains of *Candida albicans* subsequently developed infection by the same genotype of *C. albicans*, but with high MICs. Another pattern of resistance is the acquisition of infection due to azole-resistant non-*albicans* species, such as *Candida krusei* and *Candida glabrata* [31]. However, during the late 90s, the introduction of highly effective antiretroviral therapy exerted a tremendous impact on the natural history of HIV infection and its complications. Indeed, the incidence of opportunistic infections, including oropharyngeal candidiasis due to resistant strains has decreased significantly.

Outside the setting of oropharyngeal candidiasis in AIDS patients, reports of infection by resistant strains have become more frequent. With the widespread use of azoles, especially fluconazole, for prophylaxis in neutropenic cancer patients, this problem has gained increased significance. A phenomenon that has been increasingly reported among patients receiving fluconazole is the shift from highly susceptible to less-susceptible species of *Candida*. Epidemiological studies performed in patients with cancer and fungemia have shown that while the number of cases caused by *Candida albicans* has decreased, the frequency of infection due to *Candida krusei* and *Candida glabrata* has increased substantially [32-35]. While *C. krusei* is considered resistant to fluconazole, the MIC values of fluconazole for the *C. glabrata* isolates

are variable, but are much higher than those reported for *C. albicans*, *Candida tropicalis* and *Candida parapsilosis*.

This increase in the incidence of infection due to less-susceptible *Candida* species has been attributed to the widespread use of fluconazole. For instance, in a large prospective survey conducted in European institutions, antifungal prophylaxis was a strong predictor for non-*albicans* candidemia [35]. In another prospective study, 349 *Candida* isolates obtained from colonization and systemic infections were analyzed [36]. Resistance to fluconazole was observed in 3.4% of *C. albicans* isolates and in 30.7% of *C. glabrata* isolates. Only 2 strains (*C. glabrata* and *C. krusei*) were resistant to amphotericin B (MIC=1 µg/mL). In this study, previous azole exposure was associated with the isolation of azole-resistant *C. albicans* strains, but this association was not observed with *C. glabrata*.

The influence of fluconazole use on the development of azole resistance was further evaluated in 585 bone marrow transplant recipients [37]. Weekly mouthwash samples were obtained, and yielded *Candida* in 256 patients. While *C. albicans* was the most frequent species obtained before fluconazole exposure, the majority of patients who were colonized by *C. glabrata* and *C. krusei* had received fluconazole for a median of 36 days. Ninety-nine percent of *C. glabrata* isolates were resistant to fluconazole. Among 30 candidemias occurring during fluconazole prophylaxis, 14 were caused by *C. glabrata*, 6 were due to *C. krusei*, and 2 were due to *C. albicans*, and all strains were highly resistant to fluconazole. *C. albicans* strains that acquire resistance after exposure to fluconazole have been reported by other authors [38], but as shown in other studies [36,39,40], this seems to be rare.

The development of candidemia during antifungal treatment (breakthrough candidemia) is another phenomenon that can be associated with infection due to resistant strains. During the past decade, reports of the occurrence of breakthrough candidemia have been published [20,41-45]. Unfortunately, in the majority of such reports no information on the MICs was provided. In a recent study of 29 breakthrough candidemias occurring among 270 candidemias, MICs

were obtained from all patients [46]. The MIC 50s for fluconazole and amphotericin B were similar in breakthrough and non-breakthrough candidemias. Furthermore, the frequency of strains with high MICs in the entire cohort was very low (4 patients with amphotericin B MIC =2, and 4 patients with fluconazole MIC ≥64). Therefore, it seems that in most cases, breakthrough candidemia is not related to the development of infection due to resistant strains.

In Brazil, antifungal susceptibility tests of 200 bloodstream isolates of *Candida* were performed using the NCCLS method [47]. Cancer was the most frequent underlying disease, accounting for 33% of cases, and about 50% of these were neutropenic. The incidence of fluconazole resistance was 2.5%, and resistance was limited to *C. krusei* and *C. glabrata* isolates. Data from an epidemiological study conducted by the same authors showed that the incidence of these two species was low [48]. The low incidence of resistant strains was attributed to the infrequent use of fluconazole in these Brazilian hospitals.

In summary, during the past decade, fluconazole has been extensively used in neutropenic patients. This has resulted in a marked decrease in the incidence of invasive candidiasis, especially in allogeneic bone marrow transplants. However, this was accompanied by a shift from highly susceptible to less susceptible *Candida* species, by a process of selection. Indeed, *C. glabrata* has emerged as an important pathogen, causing fungemia in many countries. In addition, resistance has also emerged as a result of the acquisition of resistance in otherwise susceptible *Candida* species, but at a much smaller magnitude. Finally, the limitations of the present methods for assessing antifungal susceptibility hamper any conclusion about resistance to amphotericin B and preclude their use for therapy guidance in the clinical setting.

## References

1. Beck-Sague C., Jarvis W.R. Secular trends in the epidemiology of nosocomial fungal infections in the United States, 1980-1990. National Nosocomial Infections Surveillance System. *J. Infect. Dis.* **1993**;167:1247-51.

2. De Wit S., Urbain D., Rahir F., et al. Efficacy of oral fluconazole in the treatment of AIDS associated oesophageal candidiasis. *Eur J Clin Microbiol Infect Dis* **1991**;10:503-5.
3. Marr K.A., Seidel K., Slavin M.A., et al. Prolonged fluconazole prophylaxis is associated with persistent protection against candidiasis-related death in allogeneic marrow transplant recipients: long-term follow-up of a randomized, placebo-controlled trial. *Blood* **2000**;96:2055-61.
4. Winston D.J., Pakrasi A., Busuttill R.W. Prophylactic fluconazole in liver transplant recipients. A randomized, double-blind, placebo-controlled trial. *Ann Intern Med* **1999**;131:729-37.
5. Saag M.S., Cloud G.A., Graybill J.R., et al. A comparison of itraconazole versus fluconazole as maintenance therapy for AIDS-associated cryptococcal meningitis. National Institute of Allergy and Infectious Diseases Mycoses Study Group. *Clin Infect Dis* **1999**;28:291-6.
6. Menichetti F., Fiorio M., Tosti A., et al. High-dose fluconazole therapy for cryptococcal meningitis in patients with AIDS. *Clin Infect Dis* **1996**;22:838-40.
7. Rex J.H., Bennett J.E., Sugar A.M., et al. A randomized trial comparing fluconazole with amphotericin B for the treatment of candidemia in patients without neutropenia. Candidemia Study Group and the National Institute. *N Engl J Med* **1994**;331:1325-30.
8. Phillips P., Shafran S., Garber G., et al. Multicenter randomized trial of fluconazole versus amphotericin B for treatment of candidemia in non-neutropenic patients. Canadian Candidemia Study Group. *Eur J Clin Microbiol Infect Dis* **1997**;16:337-45.
9. Dewsnup D.H., Stevens D.A. Efficacy of oral amphotericin B in AIDS patients with thrush clinically resistant to fluconazole. *J Med Vet Mycol* **1994**;32:389-93.
10. Redding S., Smith J., Farinacci G., et al. Resistance of *Candida albicans* to fluconazole during treatment of oropharyngeal candidiasis in a patient with AIDS: documentation by *in vitro* susceptibility testing and DNA subtype analysis. *Clin Infect Dis* **1994**;18:240-2.
11. Vuffray A., Durussel C., Boerlin P., et al. Oropharyngeal candidiasis resistant to single-dose therapy with fluconazole in HIV-infected patients. *AIDS* **1994**;8:708-9.
12. Sangeorzan J.A., Bradley S.F., He X., et al. Epidemiology of oral candidiasis in HIV-infected patients: colonization, infection, treatment, and emergence of fluconazole resistance. *Am J Med* **1994**;97:339-46.
13. Lyon R., Woods G. Comparison of the BacT/Alert and Isolator blood culture systems for recovery of fungi. *Am J Clin Pathol* **1995**;103:660-2.
14. Revankar S.G., Kirkpatrick W.R., McAtee R.K., et al. A randomized trial of continuous or intermittent therapy with fluconazole for oropharyngeal candidiasis in HIV-infected patients: clinical outcomes and development of fluconazole resistance. *Am J Med* **1998**;105:7-11.
15. Saag M.S., Fessel W.J., Kaufman C.A., et al. Treatment of fluconazole-refractory oropharyngeal candidiasis with itraconazole oral solution in HIV-positive patients. *AIDS Res Hum Retroviruses* **1999**;15:1413-7.
16. Fichtenbaum CJ, Koletar S, Yiannoutsos C, et al. Refractory mucosal candidiasis in advanced human immunodeficiency virus infection. *Clin Infect Dis* **2000**;30:749-56.
17. Nucci M., Silveira M.I., Spector N., et al. Risk factors for death among cancer patients with fungemia. *Clin Infect Dis* **1998**;27:107-11.
18. Anaissie E.J., Rex J.H., Uzun O., Vartivarian S. Predictors of adverse outcome in cancer patients with candidemia. *Am J Med* **1998**;104:238-45.
19. Uzun O., Anaissie E.J. Predictors of outcome in cancer patients with candidemia. *Ann Oncol* **2000**;11:1517-21.
20. Uzun O., Asciglu S., Anaissie E.J., Rex J.H. Risk factors and predictors of outcome in patients with cancer and breakthrough candidemia. *Clin Infect Dis* **2001**;32:1713-7.
21. Kirkpatrick W.R., Turner T.M., Fothergill A.W., et al. Fluconazole disk diffusion susceptibility testing of *Candida* species. *J Clin Microbiol* **1998**;36:3429-32.
22. Colombo A.L., Barchiesi F., McGough D.A., et al. Evaluation of the E test system versus a microtitre broth method for antifungal susceptibility testing of yeasts against fluconazole and itraconazole. *J Antimicrob Chemother* **1995**;36:93-100.
23. Li R.K., Elie C.M., Clayton G.E., Ciblak M.A. Comparison of a new colorimetric assay with the NCCLS broth microdilution method (M-27A) for antifungal drug MIC determination. *J Clin Microbiol* **2000**;38:2334-8.
24. National Committee for Clinical Laboratory Standards. Reference method for broth dilution. Antifungal susceptibility testing of yeasts; Approved Standard. Document M27-A **1997**;17:1-29.
25. Espinel-Ingroff A., Kerkering T.M., Goldson P.R., Shadomy S. Comparison study of broth macrodilution and microdilution antifungal susceptibility tests. *J Clin Microbiol* **1991**;29:1089-94.
26. Fromtling R.A., Galgiani J.N., Pfaller M.A., et al. Multicenter evaluation of a broth macrodilution antifungal susceptibility test for yeasts. *Antimicrob Agents Chemother* **1993**;37:39-45.
27. Rex J.H., Pfaller M.A., Galgiani J.N., et al. Development of interpretive breakpoints for antifungal susceptibility testing: conceptual framework and analysis of *in vitro-in vivo* correlation data for fluconazole, itraconazole, and candida infections. Subcommittee on Antifungal Susceptibility Testing of the National Committee for Clinical Laboratory Standards. *Clin Infect Dis* **1997**;24:235-47.

28. Nguyen M.H., Clancy C.J., Yu V.L., et al. Do *in vitro* susceptibility data predict the microbiologic response to amphotericin B? Results of a prospective study of patients with Candida fungemia. *J Infect Dis* **1998**;177:425-30.
29. Horsburgh C.R.Jr., Kirkpatrick C.H. Long-term therapy of chronic mucocutaneous candidiasis with ketoconazole: experience with twenty-one patients. *Am J Med* **1983**;74:23-9.
30. Ruhnke M., Eigler A., Tennagen I., et al. Emergence of fluconazole-resistant strains of *Candida albicans* in patients with recurrent oropharyngeal candidosis and human immunodeficiency virus infection. *J Clin Microbiol* **1994**;32:2092-8.
31. Redding S.W., Kirkpatrick W.R., Dib O., et al. The epidemiology of non-albicans *Candida* in oropharyngeal candidiasis in HIV patients. *Spec Care Dentist* 2000;20:178-81.
32. Wingard J.R., Merz W.G., Rinaldi M.G., et al. Increase in *Candida krusei* infection among patients with bone marrow transplantation and neutropenia treated prophylactically with fluconazole. *N Engl J Med* **1991**;325:1274-7.
33. Wingard J.R., Merz W.G., Rinaldi M.G., et al. Association of *Torulopsis glabrata* infections with fluconazole prophylaxis in neutropenic bone marrow transplant patients. *Antimicrob Agents Chemother* **1993**;37:1847-9.
34. Abi-Said D., Anaissie E., Uzun O., et al. The epidemiology of hematogenous candidiasis caused by different *Candida* species. *Clin Infect Dis* **1997**;24:1122-8.
35. Viscoli C., Girmenia C., Marinus A., et al. Candidemia in cancer patients: a prospective, multicenter surveillance study by the Invasive Fungal Infection Group (IFIG) of the European Organization for Research and Treatment of Cancer (EORTC). *Clin Infect Dis* **1999**;28:1071-9.
36. Safdar A., Chaturvedi V., Cross E.W., et al. Prospective study of candida species in patients at a comprehensive cancer center. *Antimicrob Agents Chemother* **2001**;45:2129-33.
37. Marr K.A., Seidel K., White T.C., Bowden R.A. Candidemia in allogeneic blood and marrow transplant recipients: evolution of risk factors after the adoption of prophylactic fluconazole. *J Infect Dis* **2000**;181:309-16.
38. Nolte F.S., Parkinson T., Falconer D.J., et al. Isolation and characterization of fluconazole- and amphotericin B-resistant *Candida albicans* from blood of two patients with leukemia. *Antimicrob Agents Chemother* **1997**;41:196-9.
39. Hoban D.J., Zhanel G.G., Karlowsky J.A. *In vitro* susceptibilities of *Candida* and *Cryptococcus neoformans* isolates from blood cultures of neutropenic patients. *Antimicrob Agents Chemother* **1999**;43:1463-4.
40. Pfaller M.A., Diekema D.J., Jones R.N., et al. International surveillance of bloodstream infections due to *Candida* species: frequency of occurrence and *in vitro* susceptibilities to fluconazole, ravuconazole, and voriconazole of isolates collected from 1997 through 1999 in the SENTRY Antimicrobial Surveillance Program. *J Clin Microbiol* **2001**;39:3254-9.
41. Nguyen M.H., Peacock J.E. Jr., Morris A.J., et al. The changing face of candidemia: emergence of non-*Candida albicans* species and antifungal resistance. *Am J Med* **1996**;100:617-23.
42. Blumberg E.A., Reboli A.C. Failure of systemic empirical treatment with amphotericin B to prevent candidemia in neutropenic patients with cancer. *Clin Infect Dis* **1996**;22:462-6.
43. Nucci M., Colombo A.L., Spector N., et al. Breakthrough candidemia in neutropenic patients. *Clin Infect Dis* **1997**;24:275-6.
44. Girmenia C., Martino P., Cassone A. Breakthrough candidemia during antifungal treatment with fluconazole in patients with hematologic malignancies. *Blood* **1996**;87:838-9.
45. Kremery V. Jr., Sejnova D., Pichnova E. Breakthrough *Candida tropicalis* fungemia during ketoconazole prophylaxis in cancer patients. *Acta Oncol* **1999**;38:663-5.
46. Nucci M., Colombo A.L. Risk factors for breakthrough candidemia. *Eur J Clin Microbiol Infect* **2002**;21:209-11.
47. Nakagawa Z., Nucci M., Branchini M.L., et al. *In vitro* susceptibility patterns of 200 recent bloodstream isolates of *Candida* species to four antifungal drugs determined by the NCCLS microbroth procedure. Programs and Abstracts of the 40th Interscience Symposium on Antimicrobial Agents and Chemotherapy **1988**;September 24-27:488.
48. Colombo A.L., Nucci M., Salomao R., et al. High rate of non-albicans candidemia in Brazilian tertiary care hospitals. *Diagn Microbiol Infect Dis* **1999**;34:281-6.