



## Environmental Microbiology

# Polycyclic aromatic hydrocarbons degradation by marine-derived basidiomycetes: optimization of the degradation process



Gabriela A.L. Vieira<sup>a</sup>, Mariana Juventina Magrini<sup>b</sup>, Rafaella C. Bonugli-Santos<sup>c</sup>,  
Marili V.N. Rodrigues<sup>b</sup>, Lara D. Sette<sup>a,b,\*</sup>

<sup>a</sup> Universidade Estadual Paulista Júlio de Mesquita Filho (UNESP), Instituto de Bociências, Departamento de Bioquímica e Microbiologia, Rio Claro, SP, Brazil

<sup>b</sup> Universidade Estadual de Campinas (UNICAMP), Centro Pluridisciplinar de Pesquisas Químicas, Biológicas e Agrícolas, Paulínia, SP, Brazil

<sup>c</sup> Universidade Federal da Integração Latino-Americana (UNILA), Instituto Latino Americano de Ciências da Vida e da Natureza, Foz do Iguaçu, PR, Brazil

## ARTICLE INFO

## Article history:

Received 5 November 2017

Accepted 13 April 2018

Available online 3 May 2018

Associate Editor: Vânia Melo

## Keywords:

Marine biotechnology

PAHs

Fungi

Acute toxicity

Experimental design

## ABSTRACT

Pyrene and benzo[a]pyrene (BaP) are high molecular weight polycyclic aromatic hydrocarbons (PAHs) recalcitrant to microbial attack. Although studies related to the microbial degradation of PAHs have been carried out in the last decades, little is known about degradation of these environmental pollutants by fungi from marine origin. Therefore, this study aimed to select one PAHs degrader among three marine-derived basidiomycete fungi and to study its pyrene detoxification/degradation. *Marasmiellus* sp. CBMAI 1062 showed higher levels of pyrene and BaP degradation and was subjected to studies related to pyrene degradation optimization using experimental design, acute toxicity, organic carbon removal (TOC), and metabolite evaluation. The experimental design resulted in an efficient pyrene degradation, reducing the experiment time while the PAH concentration applied in the assays was increased. The selected fungus was able to degrade almost 100% of pyrene (0.08 mg mL<sup>-1</sup>) after 48 h of incubation under saline condition, without generating toxic compounds and with a TOC reduction of 17%. Intermediate metabolites of pyrene degradation were identified, suggesting that the fungus degraded the compound via the cytochrome P450 system and epoxide hydrolases. These results highlight the relevance of marine-derived fungi in the field of PAH bioremediation, adding value to the blue biotechnology.

© 2018 Published by Elsevier Editora Ltda. on behalf of Sociedade Brasileira de Microbiologia. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

\* Corresponding author at: Departamento de Bioquímica e Microbiologia – IB, Universidade Estadual Paulista Júlio de Mesquita Filho (UNESP), Av. 24A, 1515, 13506-900, Rio Claro, SP, Brazil.

E-mail: [larasette@rc.unesp.br](mailto:larasette@rc.unesp.br) (L.D. Sette).

<https://doi.org/10.1016/j.bjm.2018.04.007>

1517-8382/© 2018 Published by Elsevier Editora Ltda. on behalf of Sociedade Brasileira de Microbiologia. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## Introduction

Polycyclic aromatic hydrocarbons (PAHs) are organic compounds composed of two or more fused benzene rings. These compounds are formed during the combustion of organic molecules and their subsequent recombination<sup>1,2</sup> and are widely distributed in the environment.<sup>3,4</sup>

PAHs discharged from various activities, such as combustion of fossil fuels, shipping, use and disposal of petroleum products, agricultural burning, coal, and wood-preserving products are persistently hazardous pollutants to the environment.<sup>5</sup> Through atmospheric deposition and municipal as well as industrial waste water discharges, PAHs are distributed in the marine environment and finally enriched in the sediment due to their great ability to bind to particulate matter.<sup>6</sup>

Low molecular weight (LMW) PAHs (composed by 2–3 aromatic rings) are predominant in petrogenic sources and can be introduced into aquatic environments through oil spills, discharge from tanker operations, and municipal and urban runoff. High molecular weight (HMW) PAHs (composed by 4–6 aromatic rings), such as pyrene and BaP, are most important in pyrogenic sources and are released into the environment mainly in the form of exhaust and solid residues.<sup>7,8</sup> Both of these compounds are classified by the US Environmental Protection Agency (EPA) as priority pollutants<sup>9,10</sup> and have been accumulating in sediments due to their limited water solubility and high affinity for particulate matter.<sup>11</sup>

Microbial degradation can be considered as an attractive biotechnological alternative for achieving possible mineralization of the pollutant<sup>12</sup> and its transformation into less toxic products with greater solubility in water, which can then be degraded by the action of other microorganisms.<sup>13,14</sup>

Hydrocarbon degradation using fungi from marine origin was reported by Ahearn and Meyers.<sup>15</sup> However, their application in this field is still poorly studied. The use of marine-derived fungi for the bioremediation of polluted saline environments<sup>2</sup> is facilitated by their tolerance to saline conditions, setting these microorganisms as important microbial resources for biotechnological application in the degradation of PAH polluted environments, such as ocean and marine sediments.<sup>16</sup>

In the field of environmental pollutants biodegradation studies related to toxicity of metabolites generated during the process is required. For that end, crustaceans and rotifers larvae are applied in ecotoxicological assays due to their ecological relevance and their easy maintenance under laboratory conditions.<sup>17–19</sup> Additionally, PAHs biodegradation generally is incomplete, resulting in intermediate compounds. Determinations of PAHs degradation metabolic pathways enable to predict the destiny of such compounds in the environment and elucidate toxicity loss.<sup>20,21</sup>

Considering that PAHs are pollutants commonly found in the marine environment and that little is known about their degradation by marine microorganism, the present study aimed to screen (among the three marine-derived basidiomycetes) the best fungus able to degrade pyrene and BaP and evaluated its capacity to detoxify and degrade pyrene under saline condition.

## Material and methods

### Microorganisms

The marine derived fungal strains *Tinctoporellus* sp. CBMAI 1061, *Marasmiellus* sp. CBMAI 1062, and *Peniophora* sp. CBMAI 1063 were isolated from the Brazilian sponges *Dracmacidon reticulatum* and *Amphimedon viridis* collected in the town of São Sebastião, São Paulo State, Brazil.<sup>22</sup> All of them were taxonomically identified as previously reported by Bonugli-Santos et al.<sup>23</sup> and deposited in the Brazilian Collection of Microorganisms from Environment and Industry – CBMAI (CPQBA, UNICAMP).

### Fungal growth and initial screening

The three isolates were cultivated in 2% (m/v) malt extract agar (MA2) for ten days at 28 °C. For initial screening of PAHs degradation (pyrene and BaP), three fungal culture cylinders (0.5 cm diameter) from the edge of the colony were transferred to 125 mL Erlenmeyer flasks containing 50 mL of 2% malt extract broth (MB2) and cultured for 72 h at 28 °C and 120 rpm (initial biomass). Pyrene and BaP were dissolved in dimethylsulfoxide and were added separately at the concentration of 0.04 and 0.02 mg mL<sup>-1</sup>, respectively. The concentration of PAHs applied in this work were chosen based in previous study reported by Passarini.<sup>9</sup> The flasks were incubated for 7, 14 and 21 days under the same conditions. No saline conditions were applied for the initial screening.

### Quantitative analyses of PAHs degradation

PAHs degradation were quantified using gas chromatography–mass spectrometry (GC–MS) analyses, as previously described by Passarini et al.<sup>9</sup> The extraction of the PAHs and their degradation products in fermentative medium was accomplished using an Ultra-Turrax system. Ethyl acetate was used as solvent to rupture the fungal cellular walls and remove adsorbed compounds. To each Erlenmeyer flask containing the fungus in fermentative medium, 50 mL of ethyl acetate was added, and the fungus was subjected to cellular disintegration at 14,000 rpm for 1 min. The material was then filtered, transferred to a 250 mL separating funnel and shaken vigorously for 1 min. The organic phase was filtered by anhydrous sodium sulfate saturated with ethyl acetate directly into a 250 mL, round-bottom flask; the aqueous phase was re-extracted with an additional 50 mL of ethyl acetate. The organic phases were combined and evaporated under vacuum at 40–45 °C until the volume was reduced to approximately 2 mL. The contents of the flask were analytically transferred into a 10 mL volumetric flask and diluted to the mark with ethyl acetate. An aliquot of 1 mL of this solution was then diluted in 10 mL of ethyl acetate containing 1 mL of the internal standard solution prior to GC–MS analysis.

For the GC–MS analyses, an Agilent (Palo Alto, CA) 6890N GC that was equipped with an Agilent 7683B autosampler coupled to an Agilent 5975 mass-selective detector was used. Data acquisition and analyses were performed using

the standard software supplied by the manufacturer. The compounds were separated on a fused-silica capillary column (HP-5MS, 30 m × 0.25 mm i.e., 0.25- $\mu$ m film thickness) (J&W Scientific, Cologne, Germany). The GC temperature program was as follows: 180 °C, 5 °C min<sup>-1</sup> up to 310 °C, held for 10 min. The temperatures for the injection port and detector were set at 290 °C and 300 °C, respectively. The splitless injection mode was used with the split outlet opened after 3 min, and helium, with a flow rate of 1.0 mL min<sup>-1</sup>, was used as the carrier gas. The retention times and characteristic mass fragments were recorded, and the chosen diagnostic mass fragments were monitored in the selected ion monitoring (SIM) mode. The characteristic ions that were used for quantification were as follows: benzo[a]pyrene (*m/z* 252, 126, 250), pyrene (*m/z* 202, 101, 200) and dibutyl phthalate (*m/z* 149, 205, 223) as an internal standard. For quantification, the peak area ratios of the analytes to the internal standard were calculated as a function of the compound concentration.

### Metabolites identification

The GC-MS technique was applied for metabolites detection formed during pyrene degradation. The basidiomycete *Marasmiellus* sp. was incubated in 50 mL of liquid medium (MB2) for 5 and 15 days in the presence of pyrene (0.04 mg mL<sup>-1</sup>) at 28 °C and 120 rpm. The culture medium containing the grown mycelium was extracted in Ultra-Turrax homogenizer using 60 mL of dichloromethane as the extractor agent. After two extractions, organic phases were pooled and dried in vacuum using a rotary evaporator in a water bath at 30 °C. For each culture time, five replicas were grouped in order to obtain a higher concentration of the formed products. The final residue was dissolved in 2 mL of ethyl acetate and 1  $\mu$ L was injected into the GC-MS system. Sample components were separated on a HP-5MS capillary column (30 m × 0.25 mm × 0.25  $\mu$ m) following a temperature program: from 60 °C to 280 °C, up 5 °C per minute, kept at 280 °C for 26 min, and from 280 °C to 310 °C, up 5 °C per minute and held at 310 °C for 5 min. The injector temperature was 250 °C (splitless injection) and the interface temperature was 300 °C. Samples were analyzed in Full Scan mode in the 40–600 *m/z* range. Peaks were identified by comparing the mass spectrum and the data available in the library software NIST MS SEARCH version 2.0 and independently through fragmentation pattern interpretation. Samples were also analyzed in the Selected Ion Monitoring mode (SIM). Metabolites were identified by one identification ion and at least one additional confirmation ion per compound, as mass spectra described in literature. Control experiments, without PAH, were also analyzed.

### Experimental design and optimization of pyrene degradation

In order to investigate the culture conditions effect on pyrene degradation process by *Marasmiellus* sp., and to obtain the necessary conditions for efficient degradation, a series of experimental designs were established.<sup>24</sup> Initially, the Plackett-Burman design (Matrix with 12 assays) (PB12) was performed for the analyses of the following independent variables evaluation: temperature, pH, salinity and

initial concentrations of malt extract, glucose, peptone, yeast extract and PAH. The coded values are presented in Table 2. Salinity was adjusted with artificial seawater that was prepared according to Kester et al.<sup>25</sup> All of the different variables were prepared on two levels, designated as -1 (lower) and +1 (upper) (Table 2). Three central point assays were added to the matrix in order to determine the standard error. In a total of 15 trials, the first design allowed to evaluate the effect of 8 variables on PAH degradation after 7 days of incubation. Initial biomass was obtained at 120 rpm without the presence of pyrene, then a dimethylsulfoxide solution containing the pollutant was added in separately after 72 h of growth. The degradation quantification was performed as previously described. Variables effects on the PAH biodegradation responses were evaluated using the program STATISTICA 7.0.

Based on the results from PB12, a fractional factorial design 2<sup>4-1</sup> with three central points was purposed to evaluate the effect of four variables on pyrene biodegradation (Table 3): glucose, malt extract, peptone and PAHs after 4 days of incubation. Other growing conditions were 28 °C, pH 8, 35 ppm salinity and the absence of yeast extract. The variables glucose and peptone were selected to carry out a Central Composite Design 2<sup>2</sup> (CCD) containing 4 axial points and three repetitions at the central point in order to improve pyrene degradation rates (concentration set at 0.08 mg mL<sup>-1</sup>) after 2 days of incubation (Table 4). The malt extract concentration was set to 25 g L<sup>-1</sup> and other growth conditions were maintained as described in fractional factorial design 2<sup>4-1</sup>.


After culture conditions optimization for pyrene degradation the metabolites formed were identified as described above. Additionally, acute toxicity and the total organic carbon reduction (TOC) were analyzed as described below.

### Acute toxicity


*Artemia* sp. (brine shrimp) dry eggs were purchased from a fish store. *Artemia* sp. larvae were obtained after hatching from dry eggs in artificial seawater with constant light. Recently hatched (48 h) larvae were used for the tests, which were performed in triplicate, using test tubes containing 10% (v/v) NaCl 30% and 4.5 mL of samples, totalizing 5 mL of test solution. After preparation of the tubes, 30 larvae were added and were exposed to the samples for 24 h under light. After that period aspects as *Artemia* sp. motility and survival were observed. The mortality rate was calculated to determine toxicity and statistics tests (meta-analysis – multiple proportions – *p*-value <0.01) were run using the software BioStat 5.0.

### TOC analysis

The total organic carbon reduction (TOC) analysis was performed with a total organic carbon analyzer, TOC-L Shimadzu, as previously described by Bonugli-Santos et al.<sup>26</sup> TOC measurement is based on the combustion of the total carbon (TC) by catalytic oxidation at high temperature. The standard deviation is less than 2% in



**Fig. 1 – Pyrene (A) and BaP (B) degradation by the three marine-derived basidiomycetes after 7, 14 and 21 days of incubation at 28 °C and 120 rpm. The values are triplicate averages and vertical bars show the standard error.**



**Fig. 2 – Pyrene and BaP degradation by *Marasmiellus sp. CBMAI 1062* after 7 days of incubation at 28 °C and 120 rpm. The values are triplicate averages and vertical bars show the standard error.**

each sample analyzed in triplicate. The % TOC reduction was calculated using Eq. (1):

$$\% \text{ TOC reduction} = \left( \frac{\text{TOC}_{\text{initial}} - \text{TOC}_{(t)}}{\text{TOC}_{\text{initial}}} \right) \times 100 \quad (1)$$

where  $\text{TOC}_{\text{initial}}$  and  $\text{TOC}_{(t)}$  represent the initial TOC value and the TOC value at time 't' (h), respectively. Samples containing optimized conditions for pyrene degradation in 48 h were analyzed. Abiotic controls (without microorganism) were included during the experiment.

## Results

### PAHs degradation by marine-derived fungi

The three marine-derived basidiomycete fungi were able to degrade pyrene (Fig. 1A) and BaP (Fig. 1B) at different rates. *Marasmiellus sp.* was able to degrade more than 90% of the initial amount of both pollutants (0.02 and 0.04 mg mL<sup>-1</sup> BaP and pyrene, respectively) after 7 days of incubation. After 21 days, *Tinctoporellus sp.* reached almost 50% of

degradation, and *Peniophora sp.* was able to degrade 30% of both PAHs.

*Marasmiellus sp.* reached high levels of pyrene and BaP degradation after 7 days of incubation and was selected and submitted to additional PAHs degradation studies that were carried out after 1, 3, 5, and 7 days of incubation (Fig. 2). After 24 h of incubation, 79.4% of the BaP were degraded, and at the end of the 7th day 97.2% of BaP degradation was achieved. Pyrene was degraded at a slower rate, reaching 18.5% of initial concentration degradation after 24 h of incubation. However, after 7 days of incubation 92.8% of pyrene was degraded. Since the results from pyrene degradation were more consistent indicating a degradation curve (Fig. 2), this environmental pollutant was selected and subjected to the studies related to the optimization of degradation, including the analyses of metabolites, toxicity, and total organic carbon reduction (TOC).

### Pyrene degradation and detoxification by *Marasmiellus sp. CBMAI 1062*

Analyses through the SIM mode and Full Scan enabled the detection and identification of four metabolites after pyrene degradation by the fungus *Marasmiellus sp.* (Table 1). Metabolite I presented a molecular ion (M+)  $m/z$  218 and fragment ion  $m/z$  189, resulting in the loss of a -CHO group. It was the most abundant metabolite identified and its standard mass spectrum is characteristic of the hydroxypyrene compound. Metabolites II and III presented molecular ion (M+)  $m/z$  236 and base peak  $m/z$  218. This profile represents H<sub>2</sub>O unit loss (M+–18). Fragment ion  $m/z$  189,  $m/z$  176 and  $m/z$  94 were also

**Table 1 – Monitored ions during pyrene degradation by *Marasmiellus sp. CBMAI 1062* through the SIM mode and Full Scan.**


Compound	Monitored ions ( $m/z$ )	
	Identification	Confirmation
Hydroxypyrene	218	189
Dihydroxypyrene	234	205, 176
Pyrene dihydrodiol	236	218, 189, 176, 94



**Table 2 – Plackett-Burman (matrix 12) (PB12) experimental design applied to the evaluation of the effects of 8 factors in pyrene degradation by *Marasmiellus* sp. CBMAI 1062 after 7 days of incubation.**

Assay	Temperature (°C)	pH	Salinity (ppm)	Glucose (g/50 mL)	Malt extract (g/50 mL)	Peptone (g/50 mL)	Yeast extract (g/50 mL)	Pyrene (mg mL <sup>-1</sup> )	Pyrene degradation (%)
1	+1 (28)	-1 (7)	+1 (35)	-1 (0.25)	-1 (0.25)	-1 (0)	+1 (0.1)	+1 (0.08)	45.9
2	+1 (28)	+1 (9)	-1 (21)	+1 (0.75)	-1 (0.25)	-1 (0)	-1 (0)	+1 (0.08)	48.9
3	-1 (22)	+1 (9)	+1 (35)	-1 (0.25)	+1 (0.75)	-1 (0)	-1 (0)	-1 (0.04)	91.0
4	+1 (28)	-1 (7)	+1 (35)	+1 (0.75)	-1 (0.25)	+1 (0.1)	-1 (0)	-1 (0.04)	86.2
5	+1 (28)	+1 (9)	-1 (21)	+1 (0.75)	+1 (0.75)	-1 (0)	+1 (0.1)	-1 (0.04)	96.7
6	+1 (28)	+1 (9)	+1 (35)	-1 (0.25)	+1 (0.75)	+1 (0.1)	-1 (0)	+1 (0.08)	82.4
7	-1 (22)	+1 (9)	+1 (35)	+1 (0.75)	-1 (0.25)	+1 (0.1)	+1 (0.1)	-1 (0.04)	93.7
8	-1 (22)	-1 (7)	+1 (35)	+1 (0.75)	+1 (0.75)	-1 (0)	+1 (0.1)	+1 (0.08)	49.4
9	-1 (22)	-1 (7)	-1 (21)	+1 (0.75)	+1 (0.75)	+1 (0.1)	-1 (0)	+1 (0.08)	62.7
10	+1 (28)	-1 (7)	-1 (21)	-1 (0.25)	+1 (0.75)	+1 (0.1)	+1 (0.1)	-1 (0.04)	99.2
11	-1 (22)	+1 (9)	-1 (21)	-1 (0.25)	-1 (0.25)	+1 (0.1)	+1 (0.1)	+1 (0.08)	69.2
12	-1 (22)	-1 (7)	-1 (21)	-1 (0.25)	-1 (0.25)	-1 (0)	-1 (0)	-1 (0.04)	52.8
13 (C)	0 (25)	0 (8)	0 (28)	0 (0.5)	0 (0.5)	0 (0.05)	0 (0.05)	0 (0.06)	66.6
14 (C)	0 (25)	0 (8)	0 (28)	0 (0.5)	0 (0.5)	0 (0.05)	0 (0.05)	0 (0.06)	67.3
15 (C)	0 (25)	0 (8)	0 (28)	0 (0.5)	0 (0.5)	0 (0.05)	0 (0.05)	0 (0.06)	69.0

In parenthesis, real values.



**Fig. 3 – Standardized Pareto chart showing the effects of 8 factors (Plackett-Burman matrix 12) in pyrene degradation by *Marasmiellus* sp. CBMAI 1062, after 7 days of incubation.**


detected. The pattern of the mass spectrum obtained suggests that metabolites II and III are dihydrodiol. The two identified peaks probably correspond to different ring cleavage position. The metabolite IV presented a molecular ion (M<sup>+</sup>) *m/z* 234 and fragment ions *m/z* 205 (M<sup>+</sup>–COH) and *m/z* 176 (M<sup>+</sup>–2COH). The mass spectrum pattern obtained suggests that the metabolite IV is dihydroxypyrene.<sup>27</sup> The hydroxyl groups' position was not determined. None of these metabolites were detected in the experimental control without PAH.

The fungus *Marasmiellus* sp. was submitted to experimental design studies in order to evaluate the influence of different variables in pyrene degradation and to improve the degradation process. In this context, the influence of 8 independent factors on pyrene degradation by *Marasmiellus* sp. was investigated using PB12 and the results are shown in Table 2. Pyrene biodegradation ranges from 45.9% to 99.2%. The pH and temperature showed positive effects on pyrene degradation, while

the PAH concentration demonstrated a negative effect (Fig. 3). Salinity did not interfere significantly in the pyrene degradation (Fig. 3). However, higher degradation values were observed in saline and alkaline conditions (assays 5 and 7, Table 2).

Considering the results obtained in the PB12 experiment some variables were chosen for the next stage, a fractional factorial design 2<sup>4-1</sup>: glucose, malt extract, peptone and PAH. Since higher amounts of pyrene were degraded at high rates, the incubation time was reduced from seven to four days. High level of pyrene degradation (98.6%) was also obtained after four days of incubation (assay 8, Table 3), with the three nutrients set at their upper level.

Since the malt extract had a positive effect on pyrene degradation in the fractional factorial design 2<sup>4-1</sup> it was set at 25 g L<sup>-1</sup> (Fig. 4). The PAH concentration was set at 0.08 mg mL<sup>-1</sup>, and incubation time was reduced to two days. Central Composite Design 2<sup>2</sup> (CCD) was used to evaluate the



**Fig. 4 – Standardized Pareto chart showing the effects of 4 factors (Fractional factorial design 2<sup>4-1</sup>) in pyrene degradation by *Marasmiellus* sp. CBMAI 1062, after 4 days of incubation.**

**Table 3 – Fractional factorial design 2<sup>4-1</sup> applied to the evaluation of the effects of 4 factors in pyrene degradation by *Marasmiellus* sp. CBMAI 1062 after 4 days of incubation.**

Assay	Glucose (g/50 mL)	Malt extract (g/50 mL)	Peptone (g/50 mL)	Pyrene (mg mL <sup>-1</sup> )	Pyrene degrad. (%)
1	-1 (0)	-1 (0.75)	-1 (0.1)	-1 (0.04)	94.2
2	+1 (0.26)	-1 (0.75)	-1 (0.1)	+1 (0.08)	92.2
3	-1 (0)	+1 (1.25)	-1 (0.1)	+1 (0.08)	97.8
4	+1 (0.26)	+1 (1.25)	-1 (0.1)	-1 (0.04)	98.0
5	-1 (0)	-1 (0.75)	+1 (0.2)	+1 (0.08)	97.2
6	+1 (0.26)	-1 (0.75)	+1 (0.2)	-1 (0.04)	97.7
7	-1 (0)	+1 (1.25)	+1 (0.2)	-1 (0.04)	98.5
8	+1 (0.26)	+1 (1.25)	+1 (0.2)	+1 (0.08)	98.6
9 (C)	0 (0.13)	0 (1.0)	0 (0.15)	0 (0.06)	95.3
10 (C)	0 (0.13)	0 (1.0)	0 (0.15)	0 (0.06)	96.2
11 (C)	0 (0.13)	0 (1.0)	0 (0.15)	0 (0.06)	97.9

In parenthesis, real values.

effects of two factors, glucose and peptone. High rate of degradation (98.2%) was obtained using this strategy (assay 10, Table 4). No significant differences were observed among the assays (listed in Table 4). Thus, it was not possible to propose a mathematical model and to get the response surface.

Assays of the central points, which had the highest pyrene degradation rate were analyzed by GC-MS for the metabolites identification. The same four metabolites previously identified when the fungus was cultured under the non optimized condition (malt extract – non saline condition) after 5 and 15 days of incubation were formed when the fungus were cultured under the optimized condition (malt extract, glucose and peptone – saline condition) after 48 h of incubation.

Acute toxicity using *Artemia* sp. was carried out after the growth of *Marasmiellus* sp. under pyrene optimized degradation conditions. In the medium containing pyrene (abiotic control) the percentage of survivors was 67.2%, and low motility for the living individuals was observed. In the presence of *Marasmiellus* sp. and pyrene (bioassay) 95% of survivors with active motility were detected. Statistical tests (meta-analysis – multiple proportions) revealed significant difference between both conditions considering *p*-value <0.01. Results indicate

that there was a loss of toxicity after pyrene biodegradation by *Marasmiellus* sp.

Studies related to the organic carbon consumption (TOC analysis) were also performed using pyrene optimized degradation conditions. *Marasmiellus* sp. was able to reduce 17% of TOC after 48 h of incubation, indicating that the fungus used the PAHs as carbon sources.

## Discussion

Data derived from the initial screening revealed the potential of the three marine-derived basidiomycete fungi for PAHs degradation. However, results contradicted data from literature,<sup>28</sup> where initial degradation rates were higher for the compound with lower molecular weight (less ring numbers). The toxicity of BaP could result in an induction of BaP degradation by *Marasmiellus* sp., as a surviving strategy. Additionally, it is worthwhile to mention that the concentration of BaP (0.02 mg mL<sup>-1</sup>) was lower than the pyrene concentration (0.04 mg mL<sup>-1</sup>).

In a previous study reported by our research group<sup>9</sup> the best result of PAHs degradation was achieved by *Aspergillus sclerotiorum* CBMAI 849 that was able to degrade 99.7% of pyrene (0.07 mg mL<sup>-1</sup>) and 76.6% of BaP (0.03 mg mL<sup>-1</sup>) after 8 and 16 days of incubation, respectively. Hadibarata and Kristanti<sup>29</sup> also evaluated the pyrene biodegradation by the basidiomycete fungus *Armillaria* sp. F022 isolated from a tropical rain forest in Indonesia. About 63% of the initial amount of pyrene (0.005 mg mL<sup>-1</sup>) was biodegraded after 30 days of incubation. In another study the maximum pyrene (0.01 mg mL<sup>-1</sup>) degradation rate (99%) was achieved by *Pleurotus pulmonarius* F043 after 30 days of incubation.<sup>5</sup>

Considering the results obtained in the screening experiments the fungus *Marasmiellus* sp. was selected and cultured for 5 and 15 days aiming to detect and identify the metabolites formed during pyrene degradation. The presence of metabolites I, II, III and IV suggest that pyrene was degraded by the cytochrome P450 system and epoxide hydrolases activity. Although the hydroxyl group position was not determined, the 1-hydroxypyrene has been described as a metabolite of fungal pyrene degradation.<sup>30,31</sup> The metabolites formed by pyrene biotransformation found in this study were reported as pyrene

**Table 4 – Central composite design (2<sup>2</sup>) applied to the evaluation of the effects of two factors in pyrene degradation by *Marasmiellus* sp. CBMAI 1062 after 48 h of incubation.**

Assay	Glucose (g/50 mL)	Peptone (g/50 mL)	Pyrene degradation (%)
1	-1 (0.0175)	-1 (0.165)	92.5
2	+1 (0.1025)	-1 (0.165)	95.1
3	-1 (0.0175)	+1 (0.235)	98.0
4	+1 (0.1025)	+1 (0.235)	97.5
5	-1.41 (0)	0 (0.2)	96.9
6	+1.41 (0.12)	0 (0.2)	98.0
7	0 (0.06)	-1.41 (0.15)	97.9
8	0 (0.06)	+1.41 (0.25)	97.8
9	0 (0.06)	0 (0.2)	97.1
10	0 (0.06)	0 (0.2)	98.2
11	0 (0.06)	0 (0.2)	98.0

In parenthesis, real values.

degradation products by several fungi, including representatives of the genus *Marasmiellus*.<sup>30,32</sup> Lange et al.<sup>32</sup> reported that the isolated *Marasmiellus ramealis* JK 314 was able to degrade 76.5% of pyrene producing 1-hydroxypyrene, 1-pyrenylsulfate, 1,6-hydroxypyrene, 1,6-pyrenequinone and trans-4,5-dihydro-4,5-dihydroxypyrene metabolites after 14 days of incubation.

It is worth emphasizing that the initial biotransformation of a compound could facilitate the attack by other organisms.<sup>33</sup> In addition, the metabolites formed initially as phenols and dihydrodiol can be detoxified by conjugation to other molecules. The conjugates produced by fungal strains include sulfates, glycosides, glucuronides and xilosides.<sup>34</sup> These conjugates are generally more soluble in water than the parent compound and can be excreted from the body.

Results from statistical experimental designs showed that efficient pyrene degradation was reached by *Marasmiellus* sp. in a shorter period of time (48 h), higher concentration (0.08 mg mL<sup>-1</sup>), and under saline conditions. Statistical experimental designs can be considered as an efficient approach to determine the best culture conditions in biotechnological processes,<sup>35,36</sup> saving cost and time due to the reduction in the total number of experiments.<sup>37</sup> Considering that marine-derived fungi are adapted to the ocean conditions, the great performance of the fungus during pyrene degradation under saline condition was expected. However, several studies showed that biodegradation is negatively affected by the increasing in salt concentration.<sup>38–40</sup>

After pyrene degradation by *Marasmiellus* sp. there was a loss of toxicity, highlighting the biotechnological potential of this fungus to degrade and detoxify this environmental pollutant. Similar result was reported by Munusamy et al.<sup>41</sup> where representatives of the white rot fungus *Pycnoporus sanguineus* were able to degrade PAH with the production of non toxic compounds based on toxicity analysis using *Artemia* sp.

*Marasmiellus* sp. also showed capability to use pyrene as carbon source. Considering the short period of time (48 h) applied for pyrene degradation by the fungus, result from TOC reduction can be considered very satisfactory. A higher TOC reduction could be obtained after a longer incubation time. According to Russo et al.<sup>42</sup> mushroom compost submitted to combined ozonation and aerobic BaP biodegradation presented from 33% to 72% of TOC removals after 30 days of incubation (soil samples).

## Conclusions

Results from the present study revealed the potential of the marine-derived fungi for environmental pollutants degradation. In particular, the fungus *Marasmiellus* sp. CBMAI 1062 showed great ability to degrade PAHs and can be considered a new microbial resource for bioremediation. Experimental design was successfully applied to optimize the PAH degradation by *Marasmiellus* sp., resulting in a very high rate (almost 100%) of pyrene degradation after only 48 h of incubation. Intermediate metabolites generated suggested that the fungus used the cytochrome P450 system and epoxide hydrolases and highlight the importance of this organism in the early

stages of pyrene degradation. Pyrene toxicity was around 28% reduced and total organic carbon was 17% consumed after the degradation process under optimized condition, showing that the fungus was able to detoxify and use the pollutant as carbon source. Additionally, since this fungus was isolated from the marine environment and was able to degrade PAHs under saline conditions, biological advantages can be provided for its use in saline environments and/or processes.

## Conflicts of interest

The authors declare no conflicts of interest.

## Acknowledgements

Gabriela A. L. Vieira was supported by a PhD scholarship from the São Paulo Research Foundation – FAPESP (2012/12622-3) and Mariana J. Magrini was supported by a Master's Degree scholarship from the Coordination for the Improvement of Higher Education Personnel – CAPES. The authors thank FAPESP for its financial support (Grants 2010/50190-2, 2013/19486-0 and 2016/07957-7). Lara D. Sette thanks the National Council for Scientific and Technological Development (CNPq) for the Productivity Fellowship (304103/2013-6).

## REFERENCES

1. Haritash AK, Kaushik CP. Biodegradation aspects of Polycyclic Aromatic Hydrocarbons (PAHs): a review. *J Hazard Mater.* 2009;169(1–3):1–15.
2. Raghukumar C, D'Souza-Ticlo D, Verma A. Treatment of colored effluents with lignin-degrading enzymes: an emerging role of marine-derived fungi. *Crit Rev Microbiol.* 2008;34(3–4):189–206.
3. Shuttleworth KL, Cerniglia E. Environmental aspects of PAH biodegradation. *Appl Biochem Biotechnol.* 1995;54(1):291–302.
4. Abdel-Shafy HI, Mansour MSM. A review on polycyclic aromatic hydrocarbons: source, environmental impact, effect on human health and remediation. *Egypt J Pet.* 2016;25(1):107–123.
5. Hadibarata T, Teh ZC. Optimization of pyrene degradation by white-rot fungus *Pleurotus pulmonarius* F043 and characterization of its metabolites. *Bioprocess Biosyst Eng.* 2014;37(8):1679–1684.
6. Lang S-C, Hursthouse A, Mayer P, et al. Equilibrium passive sampling as a tool to study polycyclic aromatic hydrocarbons in Baltic Sea sediment pore-water systems. *Mar Pollut Bull.* 2015;101(1):296–303.
7. Souza HML, Taniguchi S, Bicego MC, et al. Polycyclic aromatic hydrocarbons in superficial sediments of the Negro River in the Amazon Region of Brazil. *J Braz Chem Soc.* 2015;26:1438–1449.
8. Syed JH, Iqbal M, Zhong G, et al. Polycyclic aromatic hydrocarbons (PAHs) in Chinese forest soils: profile composition, spatial variations and source apportionment. *Sci Rep.* 2017;7:2692.
9. Passarini MRZ, Rodrigues MVN, da Silva M, Sette LD. Marine-derived filamentous fungi and their potential application for polycyclic aromatic hydrocarbon bioremediation. *Mar Pollut Bull.* 2011;62(2):364–370.
10. Zhang S, Yao H, Lu Y, et al. Uptake and translocation of polycyclic aromatic hydrocarbons (PAHs) and heavy metals

- by maize from soil irrigated with wastewater. *Sci Rep*. 2017;7(1):12165.
11. Rothermich MM, Hayes LA, Lovley DR. Anaerobic, sulfate-dependent degradation of polycyclic aromatic hydrocarbons in petroleum-contaminated harbor sediment. *Environ Sci Technol*. 2002;36(22):4811–4817.
  12. Raghukumar C, Shailaja M, Parameswaran P, Singh S. Removal of polycyclic aromatic hydrocarbons from aqueous media by the marine fungus NIOCC# 312: involvement of lignin-degrading enzymes and exopolysaccharides. *Indian J Mar Sci*. 2006.
  13. Cerniglia CE. Fungal metabolism of polycyclic aromatic hydrocarbons: past, present and future applications in bioremediation. *J Ind Microbiol Biotechnol*. 1997;19(5):324–333.
  14. Cerniglia CE, Sutherland JB. *Bioremediation of Polycyclic Aromatic Hydrocarbons by Lignolytic and Non-Lignolytic Fungi*. vol. 23; 2001:136–187.
  15. Ahearn DG, Meyers SP. The role of fungi in the decomposition of hydrocarbons in the marine environment. In: Walters AH, Hueck-van der Plas EH, eds. *Biodeterioration of Materials*. London: Applied Science; 1972:12–18.
  16. Bonugli-Santos RC, Vasconcelos R dos S, Passarini MRZ, et al. Marine-derived fungi: diversity of enzymes and biotechnological applications. *Front Microbiol*. 2015;6, <http://dx.doi.org/10.3389/fmicb.2015.00269>.
  17. Garaventa F, Gambardella C, Di Fino A, Pittore M, Faimali M. Swimming speed alteration of *Artemia* sp. and *Brachionus plicatilis* as a sub-lethal behavioural end-point for ecotoxicological surveys. *Ecotoxicology*. 2010;19(3):512–519.
  18. Piazza V, Ferioli A, Giacco E, et al. A standardization of *Amphibalanus (Balanus) amphitrite* (Crustacea, Cirripedia) larval bioassay for ecotoxicological studies. *Ecotoxicol Environ Saf*. 2012;79:134–138.
  19. Costa E, Piazza V, Gambardella C, et al. Ecotoxicological effects of sediments from Mar Piccolo, South Italy: toxicity testing with organisms from different trophic levels. *Environ Sci Pollut Res*. 2016;23(13):12755–12769.
  20. Carmona M, Zamarró MT, Blázquez B, et al. Anaerobic Catabolism of Aromatic Compounds: a Genetic and Genomic View. *Microbiol Mol Biol Rev MMBR*. 2009;73(1):71–133.
  21. Ghosal D, Ghosh S, Dutta TK, Ahn Y. Current state of knowledge in microbial degradation of polycyclic aromatic hydrocarbons (PAHs): a review. *Front Microbiol*. 2016;7:1369.
  22. Menezes CBA, Bonugli-Santos RC, Miqueletto PB, et al. Microbial diversity associated with algae, ascidians and sponges from the north coast of São Paulo state, Brazil. *Microbiol Res*. 2010;165(6):466–482.
  23. Bonugli-santos RC, Durrant LR, Sette LD. Laccase activity and putative laccase genes in marine-derived basidiomycetes. *Fungal Biol*. 2010;114(10):863–872.
  24. Rodrigues M, Iemma A. *Planejamento de experimentos e otimização de processos*. 2ª Campinas; 2009.
  25. Kester DR, Duedall IW, Connors DN, Pytkowicz RM. Preparation of artificial seawater. *Limnol Oceanogr*. 1967;12(1):176–179.
  26. Bonugli-Santos RC, Vieira GAL, Collins C, et al. Enhanced textile dye decolorization by marine-derived basidiomycete *Peniophora* sp. CBMAI 1063 using integrated statistical design. *Environ Sci Pollut Res*. 2016;23(9):8659–8668.
  27. Lambert M, Kremer S, Sterner O, Anke H. Metabolism of pyrene by the basidiomycete *Crinipellis stipitaria* and identification of pyrenequinones and their hydroxylated precursors in strain JK375. *Appl Environ Microbiol*. 1994;60(10):3597–3601.
  28. Wang C, Sun H, Li J, Li Y, Zhang Q. Enzyme activities during degradation of polycyclic aromatic hydrocarbons by white rot fungus *Phanerochaete chrysosporium* in soils. *Chemosphere*. 2009;77(6):733–738.
  29. Hadibarata T, Kristanti RA. Biodegradation and metabolite transformation of pyrene by basidiomycetes fungal isolate *Armillaria* sp. F022. *Bioprocess Biosyst Eng*. 2013;36(4):461–468.
  30. Lange B, Kremer S, Sterner O, Anke H. Pyrene metabolism in *Crinipellis stipitaria*: identification of trans-4,5-dihydro-4,5-dihydroxyppyrene and 1-pyrenylsulfate in strain JK364. *Appl Environ Microbiol*. 1994;60(10):3602–3607.
  31. Cerniglia CE, Sutherland JB. Degradation of polycyclic aromatic hydrocarbons by fungi. In: Timmis KN, ed. *Handbook of Hydrocarbon and Lipid Microbiology*. Berlin, Heidelberg: Springer Berlin Heidelberg; 2010:2079–2110.
  32. Lange B, Kremer S, Anke H, Sterner O. Metabolism of pyrene by basidiomycetous fungi of the genera *Crinipellis*, *Marasmius*, and *Marasmiellus*. *Can J Microbiol*. 1996;42(11).
  33. Keck J, Sims RC, Coover M, Park K, Symons B. Evidence for cooxidation of polynuclear aromatic hydrocarbons in soil. *Water Res*. 1989;23(12):1467–1476.
  34. Sutherland JB. Detoxification of polycyclic aromatic hydrocarbons by fungi. *J Ind Microbiol*. 1992;9(1):53–61.
  35. Pearce AR, Rizzo DM, Watzin MC, Druschel GK. Unraveling associations between cyanobacteria blooms and in-lake environmental conditions in Missisquoi Bay, Lake Champlain, USA, using a modified self-organizing map. *Environ Sci Technol*. 2013;47(24):14267–14274.
  36. Zeng T, Arnold WA. Clustering chlorine reactivity of haloacetic acid precursors in inland lakes. *Environ Sci Technol*. 2014;48(1):139–148.
  37. Chauhan K, Trivedi U, Patel KC. Statistical screening of medium components by Plackett–Burman design for lactic acid production by *Lactobacillus* sp. KCP01 using date juice. *Bioresour Technol*. 2007;98(1):98–103.
  38. Piedad Díaz M, Grigson JWS, Peppiatt JC, Burgess GJ. Isolation and characterization of novel hydrocarbon-degrading euryhaline consortia from crude oil and mangrove sediments. *Mar Biotechnol*. 2000;2(6):522–532.
  39. Valentín L, Feijoo G, Moreira MT, Lema JM. Biodegradation of polycyclic aromatic hydrocarbons in forest and salt marsh soils by white-rot fungi. *Int Biodeterior Biodegrad*. 2006;58(1):15–21.
  40. Minai-Tehrani D, Minoui S, Herfatmanesh A. Effect of salinity on biodegradation of polycyclic aromatic hydrocarbons (PAHs) of heavy crude oil in soil. *Bull Environ Contam Toxicol*. 2009;82(2):179–184.
  41. Munusamy U, Sabaratnam V, Muniandy S, Abdullah N, Pandey A, Jones E. Biodegradation of polycyclic aromatic hydrocarbons by laccase of *Pycnoporus sanguineus* and toxicity evaluation of treated PAH. *Biotechnology*. 2008;7(4):669–677.
  42. Russo L, Rizzo L, Belgiorno V. Ozone oxidation and aerobic biodegradation with spent mushroom compost for detoxification and benzo(a)pyrene removal from contaminated soil. *Chemosphere*. 2012;87(6):595–601.