

EFFECT OF *CINNAMOMUM ZEYLANICUM* BLUME ESSENTIAL OIL ON THE GROWTH AND MORPHOGENESIS OF SOME POTENTIALLY PATHOGENIC *ASPERGILLUS* SPECIES

Egberto Santos Carmo¹; Edeltrudes de Oliveira Lima²; Evandro Leite de Souza^{3*}; Frederico Barbosa de Sousa⁴

¹Laboratório de Tecnologia Farmacêutica, Universidade Federal da Paraíba, João Pessoa, PB, Brasil; ²Laboratório de Micologia, Departamento de Ciências Farmacêuticas, Centro de Ciências da Saúde, Universidade Federal da Paraíba, João Pessoa, PB, Brasil; ³Laboratório de Microbiologia de Alimentos, Departamento de Nutrição, Centro de Ciências da Saúde, Universidade Federal da Paraíba, João Pessoa, PB, Brasil; ⁴Laboratório de Microscopia e Imagem Biológica, Centro de Ciências da Saúde, Universidade Federal da Paraíba, João Pessoa, PB, Brasil

Submitted: July 05, 2007; Returned to authors for corrections: November 26, 2007; Approved: January 20, 2008.

ABSTRACT

Cinnamomum zeylanicum Blume is known for a wide range of medicinal properties. This study aimed to assess the interference of *C. zeylanicum* essential oil on the growth and morphogenesis of some potentially pathogenic *Aspergillus* species. The essential oil presented strong antifungal effect causing the growth inhibition of the assayed strains and development of large growth inhibition zones. MIC₅₀ and MIC₉₀ values were 40 and 80 µL/mL, respectively. 80, 40 and 20 µL/mL of the oil strongly inhibited the radial mycelial growth of *A. niger*, *A. flavus* and *A. fumigatus* along 14 days. 80 and 40 µL/mL of the oil caused a 100% inhibition of the fungal spore germination. Main morphological changes observed under light microscopy provided by the essential oil in the fungal strains were decreased conidiation, leakage of cytoplasm, loss of pigmentation and disrupted cell structure indicating fungal wall degeneration. It is concluded that *C. zeylanicum* essential oil could be known as potential antifungal compound, particularly, to protect against the growth of *Aspergillus* species.

Key words: *Aspergillus*, *Cinnamomum zeylanicum* Blume, essential oil, antifungal activity.

INTRODUCTION

Aspergilli are ubiquitous fungi found in air, soil, plants and decayed organic materials. Human contamination by *Aspergillus* has been related to the inhalation of conidia, colonization of wounds and/or penetration in human tissues through surgical interventions (13,30). Aspergillosis is an opportunistic infection which can attack the lungs, ears, eyes, digestive system, kidneys and brain (12,19).

Aspergillosis generally rises as a respiratory disease characterized for the presence of granulomatous lesions in lungs or bronchi, followed by dissemination to other organs throughout the bloodstream (3,14). Cutaneous aspergillosis, aspergillar otomycosis, aspergillar onychomycosis, invasive lung aspergillosis, aspergillar sinusitis, immunoallergic

aspergillosis, aspergilloma and mycotoxicosis are some clinical forms of aspergillosis (19,30).

The growth of Aspergilli in foodstuffs is toxicologically significant since some species are known to produce mycotoxins. Aflatoxins-B₁, B₂, G₁ and G₂ (produced by *A. flavus* and *A. parasiticus*), aspergillic acid (produced by *A. flavus*), hydroxyaspergillic acid (produced by *A. flavus*), ochratoxins (produced by *A. ochraceus*), oxalic acid (produced by *A. flavus* and *A. glaucus*), terreic acid (produced by *A. terreus*) are some mycotoxins produced by *Aspergillus* species (6,37). The consumption of mouldy products can cause human or animal mycotoxicoses, and more importantly some mycotoxins are potent carcinogens (23,26).

Resistance of *Aspergilli* to some clinically used antifungals brings a worrying clinical prognostic in people attacked by

*Corresponding Author. Mailing address: Laboratório de Microbiologia de Alimentos, Departamento de Nutrição, Centro de Ciências da Saúde, Universidade Federal da Paraíba, João Pessoa, Paraíba, Brasil. E-mail: evandroleitesouza@ccs.ufpb.br

aspergillosis (11,14). For over fifty years antibiotics have been applied for treating or inhibiting infections. The wide use and sometimes misuse of chemo-antimicrobials in both human and animal medicine has been responsible for the selection of resistant strains (17,21).

Regarding the increasing clinical importance given to fungi causing infections and the development of drug resistance many scientific studies focusing the antifungal properties of plant products have been carried out (4,41). Antifungal effect of medicinal plants and derivatives has been scientifically proven in assays with essential oils, extracts and isolated phytochemicals (20,26,42). Essential oils as antimicrobial agents present two main characters: their natural origin generally means more safety to people and environment; and they can be considered at low risk for development of microbial resistance since they are mixtures of compounds which may present different mechanisms of antimicrobial activity (15,29).

Cinnamomum zeylanicum Blume, Lauraceae, has many biological properties as analgesic, antiseptic, antispasmodic, aphrodisiac, astringent, carminative, haemostatic, insecticidal and parasiticide. Barks from branches, without the epidermis and subereous layer, is marketed as the commercial cinnamon which has long use in perfumery, culinary and native medicine fields (2,20). Previous research has revealed interesting antimicrobial effect in *C. zeylanicum* essential oil (10,27,28). Camphene, linalool, α -phelendrene, α -terpinene, limonene, β -cymene, α -cariophyllene, cinnamaldheyde and eugenol are some of the compounds found in *C. zeylanicum* essential oil (24,44).

The present study aimed to evaluate the effect of *C. zeylanicum* Blume essential oil on the growth and morphogenesis of some *Aspergillus* species known as potential etiological agent of fungal infections.

MATERIAL AND METHODS

Essential oil

C. zeylanicum Blume essential oil was supplied by Ferquima Ind. Com. Ltda. (Vargem Grande Paulista, São Paulo, Brazil) and its quality parameters (appearance, color, purity, odor, density - 20°C, refraction index -20°C) were described in an accompanying technical report. The essential oil was assayed at concentrations ranging from 320 to 5 μ L/mL and the solutions were prepared according to Souza *et al.* (42).

Fungal strains

Aspergillus fumigatus (ATCC-16913 and ATCC-40640), *A. niger* (P-03 and LM-257), *A. flavus* (ATCC-16013 and LM-247), *A. parasiticus* (ATCC-15517 and NRRL-2999), *A. terreus* (UP-03 and ATCC-7860) and *A. ochraceus* (ATCC-7860 and LM06) strains were used as test microorganisms. These strains were obtained from the Microorganisms Collection, Laboratory of Clinical Mycology, Department of Pharmaceutical Sciences,

Health Sciences Center, Federal University of Paraíba, Brazil. Stock cultures were maintained on Sabouraud agar (SA) slants and stored in a refrigerator (7°C, \pm 1°C).

Minimum Inhibitory Concentration - MIC

The MIC of the essential oil was determined by a qualitative method using the solid medium diffusion procedure (22). For this, 1 mL of the fungal homogenous suspension (approximately 10⁶ spores/mL) prepared according to Rana *et al.* (32) was uniformly spread on sterile SA Petri dishes. After inoculum absorption by SA, wells were made using sterile glass stems (diameter 6 mm) which were filled with 50 μ L of the essential oil solution. The incubation period was 7-10 days at 25-28°C. At the end of the incubation period, the MIC was the lowest essential oil concentration showing growth inhibition zones with diameter equal to or greater than 10 mm (41,42).

Amphotericin B (100 μ g/mL) and ketoconazole (50 μ g/mL) were used as control using the solid medium diffusion procedure using filter paper discs (Cecon, diameter 6 mm) (7). The control of viability of the assayed fungal strains was carried out by observing their capability of growing on SA without adding the essential oil or standard antifungals.

Mycelial growth inhibition

Inhibition of the fungal mycelial growth was determined using the poisoned substrate technique by the daily measure of the radial mycelial growth on SA added of 80, 40 and 20 μ L/mL of the essential oil (1,15). For this, a 2 mm plug taken from a 10-days-old fungal colony cultivated on SA slants was placed on the center of the sterile SA Petri dishes containing the essential oil and incubated at 25-28°C. At different time intervals (1, 2, 4, 6, 9 and 14 days) of incubation, the radial mycelial growth was measured (mm) using calipers. The control was the observation of the radial mycelial growth on SA added of ketoconazole (50 μ g/mL) and without adding the essential oil.

Spore germination assay

An aliquot of 0.2 mL of each concentration of the essential oil (80, 40 and 20 μ L/mL) was mixed with 0.2 mL of the fungal spore suspension (approximately 10⁶ spores/mL). The mixture was placed on separated glass slides which were incubated in a moisture chamber at 25-28°C for 24 hours. At the end of the incubation period, each slide was fixed with lacto-phenol-cotton blue stain and observed under the microscope for spore germination. Control without essential oil was tested in the same way. About 200 spores were counted and the per cent of spore germination was calculated in comparison with the control assay (32,38).

Fungal morphogenesis study

For evaluating morphological alterations caused by the essential oil in *A. niger* P-03 a sample of mycelium was taken

from the periphery of a 10-days-old fungal colony grown on SA at 25–28°C containing the essential oil (80 µL/mL). The samples were fixed in lacto-phenol–cotton blue stain and observed under the microscope at 400 x to examine morphological abnormalities. Control assay without essential oil was tested in the same way (38).

All antifungal assays were carried out in duplicate and the results were expressed as an average of the two parallel assays.

Statistical analysis

Statistical analysis was performed to determine significant differences ($P < 0.05$) by the Tukey test in the mycelial radial growth assays. For this the Sigma stat 2.03 computer program was used.

RESULTS

Results of the inhibitory effect of *C. zeylanicum* essential oil in a solid medium on some potentially pathogenic *Aspergillus* species are shown in Table 1. The oil at 320 - 80 µL/mL strongly inhibited the growth of all assayed strains. 40 µL/mL was the MIC₅₀ (lowest concentration causing a growth inhibition of 50 percent or more of the assayed strains), while 80 µL/mL was the MIC₉₀ (lowest concentration causing a growth inhibition of 90 percent or more of the assayed strains) found for *C. zeylanicum* essential oil. The essential oil at 320 to 40 µL/mL provided growth inhibition zones with diameter equal to or higher than the ones caused by standard antifungals (amphotericin B and ketoconazole).

Fig. 1, 2 and 3 show the effect of *C. zeylanicum* essential oil (80, 40 and 20 µL/mL) and ketoconazole on the radial mycelial growth of *A. flavus* LM-247, *A. fumigatus* ATCC-40640 and *A. niger* P-03 using the poisoned substrate technique. The essential oil provided a fungicidal effect noted by total inhibition of the mycelial growth along 14 days of exposure. The oil provided significant ($P < 0.05$) inhibitory effect on the mycelial growth when compared with the control assay and ketoconazole. Only *A. niger* showed small mycelial growth up to 8 days of exposure when exposed to 20 µL/mL of the oil.

Ketoconazole showed no significant ($P < 0.05$) reduction in the mycelial growth after 14 days of exposure in comparison with the control assay. Ketoconazole was inserted in the radial mycelial growth assay because no tested strain showed resistance to it in the MIC assay.

Results obtained from the effect of *C. zeylanicum* essential oil (80, 40 and 20 µL/mL) on spore germination of *A. flavus* LM-247, *A. fumigatus* ATCC-40640 and *A. niger* P-03 are shown in Table 2. The oil at different concentrations caused an interesting inhibition of the spore germination. A 100% inhibition was found at 80 and 40 µL/mL of the oil, while it was over 90 and 80% at 20 µL/mL for *A. flavus* and *A. fumigatus*, respectively. *A. niger* presented a 25% of spore germination inhibition at 20 µL/mL.




Figure 1. Effect of *C. zeylanicum* essential oil and ketoconazole on the radial mycelial growth kinetic of *A. flavus* LM-247.





Figure 2. Effect of *C. zeylanicum* essential oil and ketoconazole on the radial mycelial growth kinetic of *A. fumigatus* ATCC-40640.

Table 1. Inhibitory effect of *C. zeylanicum* essential oil on the growth of some *Aspergillus* species (results expressed in millimeters of mould growth inhibition zones).

Moulds	C. zeylanicum essential oil ($\mu\text{L}/\text{mL}$)								Control	
	320	160	80	40	20	10	5	Amp B ^a (100 $\mu\text{g}/\text{mL}$)	Ketoc ^b (50 $\mu\text{g}/\text{mL}$)	Viab ^c
<i>A. fumigatus</i> ATCC-16913	22	20	18	12	8	0	0	8	18	+
<i>A. fumigatus</i> ATCC-40640	25	20	17	13	0	0	0	0	10	+
<i>A. niger</i> P-03	24	17	12	0	0	0	0	12	10	+
<i>A. niger</i> LM-257	22	20	15	6	0	0	0	8	10	+
<i>A. flavus</i> ATCC-16013	20	17	10	11	0	0	0	0	24	+
<i>A. flavus</i> LM-247	18	16	11	10	0	0	0	7	15	+
<i>A. parasiticus</i> ATCC-15517	17	11	8	0	0	0	0	8	20	+
<i>A. parasiticus</i> NRRL-2999	20	17	12	13	0	0	0	7	20	+
<i>A. terreus</i> UP-03	17	15	13	10	0	0	0	0	18	+
<i>A. terreus</i> ATCC-7860	20	18	14	12	0	0	0	0	20	+
<i>A. ochraceus</i> ATCC-7860	26	20	17	15	0	0	0	5	12	+
<i>A. ochraceus</i> LM-06	23	20	17	15	0	0	0	0	12	+

^a amphotericin B; ^b ketoconazole; ^c strain viability: ability of the strain to grow in Sabouraud agar without adding essential oil or synthetic antibiotic.

**Figure 3.** Effect of *C. zeylanicum* essential oil and ketoconazole on the radial mycelial growth kinetic of *A. niger* P-03.

Spores which germinated when exposed to 20 $\mu\text{L}/\text{mL}$ of essential oil produced smaller germ tubes (early growing hyphae) in comparison with the control assay (data not showed).

Table 2. Inhibition of *C. zeylanicum* essential oil on spore germination of *Aspergillus* species (results expressed in percent of spore germination inhibition in comparison with the control assay).

Moulds	C. zeylanicum essential oil		
	20 $\mu\text{L}/\text{mL}$	40 $\mu\text{L}/\text{mL}$	80 $\mu\text{L}/\text{mL}$
<i>A. flavus</i> LM-247	92%	100%	100%
<i>A. fumigatus</i> ATCC-40640	86%	100%	100%
<i>A. niger</i> P-03	25%	100%	100%

Observations of *A. niger* examined under the light microscope at 400 x magnification after exposure to 80 $\mu\text{L}/\text{mL}$ of *C. zeylanicum* essential oil showed some morphological abnormalities (Fig. 4). Microscopic examination of the control mycelium (untreated cell) showed a regular cell structure with homogenous cytoplasm, clearly visible sterigmata bearing conidia and profuse conidiation on a large and radiated conidial head. The mycelial growth of *A. niger* in the medium containing the essential oil appeared to present morphological changes with a heterogeneous mycelial structure. In addition, the alterations observed included decreased conidiation (lack of sporulation), visible loss of cytoplasm content, loss of pigmentation, aberrant development of hyphae and fragmentation. The essential oil clearly caused reduction in conidial heads, with distorted presence of conidiophores.




Figure 4. Light microphotographs of *A. niger* mycelium growing on AS without or with *C. zeylanicum* essential oil during 7 days of incubation at 25–28°C. (I) Control conidial head of *A. niger*, large and radiated, development of vesicle on conidiophore, conidia clearly visible, Bar 100 µm. (II-III) Modifications of conidial head of *A. niger* induced by 80 µL/mL of *C. zeylanicum* essential oil showing clear decrease in conidiation, Bar 100 µm.

DISCUSSION

This report describes the effectiveness of *C. zeylanicum* essential oil against *Aspergillus* species by different testing procedures. A wide range of antifungal activities was found. Preliminary experiments were carried out *in vitro* using the solid medium diffusion procedure in order to find the MIC of the essential oil on 12 *Aspergillus* strains on the basis of the diameter of mould growth inhibition zones. The essential oil up to 80 µL/mL caused a strong mould growth inhibition. Anti-*Aspergillus* activity presented a dose dependent effect.

The poisoned substrate technique showed a strong fungicidal effect of *C. zeylanicum* essential oil at 80 and 40 µL/mL with a sustained and broad spectrum of inhibition over time. However, at 20 µL/mL of the oil occurred a steady slow growth rate during the first part of the period and disappearing after 8 days of exposure. Soliman & Badea (40) reported complete inhibition of *A. flavus*, *A. parasiticus* and *A. ochraceus* by cinnamon oil (< 500 ppm). It has been reported that the inhibition of mould mycelial growth (fungistatic or fungicidal effect) in solid or liquid medium by essential oils over a wide range of concentrations has been accompanied by concomitant decease or total inhibition of mycotoxins production (e.g. by *Aspergillus parasiticus*, *A. ochraceus*, *Fusarium graminearum*, *F. proliferatum*) (4,5,15,26,33).

The essential oil presented an intense suppressing effect on the fungal spore germination. In agreement with earlier researches (32,38) the inhibition of spore germination caused by *C. zeylanicum* essential oil was in a dosage response manner. The capability of essential oils to inhibit fungal spore

germination has been noted in other studies using different testing methods (31,34,38).

Earlier reports showed that some essential oils are able to inhibit the mycelial growth of fungi in laboratorial media and foodstuffs (4,33,43). However, the antifungal property of different essential oils ranges from a narrow to wide spectrum depending on the assayed essential oil, its concentration and the target fungal (8,15,18,36,46).

The antifungal property of phytochemicals found in *C. zeylanicum* essential oil (e.g. mono- and sesqui-terpenes, phenols, cynamaldheyde) could involve inhibition of extracellular enzymes synthesis and the disruption of the cell wall structure resulting in lack of cytoplasm, damage of integrity and ultimately the mycelial death. Cytoplasm granulation, cytoplasm membrane rupturing, cytoplasm hyperacidity, break down of the electron transport chain, H⁺-ATPase and channel inhibition are some structural and metabolic events possibly related to the antifungal property of essential oils (9,10,25). It is also reported that essential oils are able to interfere into the mitochondrial membrane system by a membrane-disruptive activity closely associated with the enzymatic reactions, such as respiratory electron transport, protein transport and coupled phosphorylation (4,35).

Velluti *et al.* (11) report that eugenol (a phenolic compound known as major component of *C. zeylanicum* essential oil) presents its antimicrobial activity attributed to the presence of an aromatic nucleus and a phenolic OH group known to be reactive and to form hydrogen bonds with active sites of target enzymes. Although, the antimicrobial activity of an essential oil is attributed mainly to its major components, the synergistic or antagonistic effect of compounds in minor percentage in the mixture has to be considered (15,42).

The observations of light microscopy showed that the main morphological changes caused by *C. zeylanicum* essential oil on *A. niger* were associated with the degeneration of fungal hyphae causing leakage of cytoplasm content and loss of conidiation. These modifications in the cytological structure may be related to the interference of the essential oil with the enzymes responsible for wall synthesis as previously cited by other researchers (39,45). De Billerbeck *et al.* (16) noted that *Cymbopogon nardus* (L.) W. Watson essential oil was able to cause morphological changes in *A. niger* including heterogeneous mycelium structure, granular/vesicular aspect of the cytoplasm content and complete degeneration of fungal hyphae resulting in an empty hyphae tip and cytoplasm retraction. Ultrastructural changes were smooth cell walls, depressions (like craters) on the cell surface. Such changes could be related to the interference of its components on enzymatic reactions of wall synthesis, which affects fungal morphogenesis and growth.

Sharma and Tripathi (38) noted that *Citrus sinensis* (L.) Osberk essential oil caused reduction in the conidial heads, poorly developed sterigmata and distorted (squashed and

flattened) conidiophores in *A. niger*. Anomalous empty, budded and flattened hyphae and cell wall destruction was also found. These effects results in the death of hyphae suggesting that the essential oil antifungal property is a result of its attack on the cell wall and retraction of cytoplasm. Rasooli and Owlia (34) observed that thyme oils provided irreversible damage to cell wall (degenerative changes), cytoplasm membrane (irregular, dissociated from cell wall, invaginated) and nuclear membrane (folding) of *Aspergillus parasiticus*.

In conclusion, our results indicate that essential oils could find a practical and rational use in the inhibition of mould growth. Particularly, *C. zeylanicum* essential oil possesses strong anti-*Aspergillus* activity inhibiting the growth, spore germination and causing deleterious cellular morphological changes of different *Aspergillus* species. The broad inhibition of fungal growth and sporulation by *C. zeylanicum* essential oil, in addition to its availability as natural volatile product, justifies its possible rational use as an alternative antifungal compound to control the growth and dissemination of pathogen *Aspergillus* species.

RESUMO

Efeito do óleo essencial de *Cinnamomum zeylanicum* Blume sobre o crescimento e morfogênese de algumas espécies de *Aspergillus* potencialmente patogênicas

Cinnamomum zeylanicum Blume é uma planta conhecida por apresentar ampla variedade de propriedades medicinais. Portanto, este estudo teve por objetivo avaliar a interferência do óleo essencial *C. zeylanicum* sobre o crescimento e morfogênese de algumas espécies de *Aspergillus* potencialmente patogênicas. O óleo essencial testado apresentou potente efeito antifúngico demonstrado pela visualização de grandes zonas de inibição de crescimento de todas as linhagens testadas. Os valores de CIM₅₀ e de CIM₉₀ foram 40 e 80 µL/mL, respectivamente. Nas concentrações de 80, 40 e 20 µL/mL o óleo demonstrou um potente efeito fumigante, inibindo o crescimento micelial radial de *A. niger*, *A. flavus* e *A. fumigatus* ao longo de 14 dias de exposição. A 80 e 40 µL/mL o óleo essencial promoveu inibição de 100% da germinação de esporos, das três espécies de *Aspergillus* citadas anteriormente. Além disso, alterações morfológicas no crescimento fúngico foram observadas sob microscopia óptica após exposição ao óleo essencial, como diminuição da conidiação, perda citoplasmática, perda de pigmentação e rompimento da estrutura fúngica (hifa) indicando degeneração da parede celular. Diante do exposto, conclui-se que o óleo essencial de *C. zeylanicum* poderia ser empregado como potente composto antifúngico, particularmente, prevenindo crescimento de espécies de *Aspergillus*.

Palavras-chave: *Aspergillus*, *Cinnamomum zeylanicum* Blume, óleo essencial, atividade antifúngica.

REFERENCES

- Adam, K.; Sivropoulou, A.; Kokkini, S.; Lanaras, T.; Arsenakis, M. (1998). Antifungal activities of *Origanum vulgare* subsp. *hirtum*, *Mentha spicata*, *Lavandula angustifolia*, and *Salvia fruticosa* essential oils against human pathogenic fungi. *J. Agric. Food Chem.*, 46, 1739-1745.
- Almeida, E.R. (1993). *Plantas medicinais brasileiras*. Hemus Editora, São Paulo.
- Almeida, M.B.; Bussamra, M.H.; Rodrigues, J.C. (2006). Allergic bronchopulmonary aspergillosis in paediatric cystic fibrosis patients. *Paed. Resp. Rev.*, 7, 67-72.
- Atanda, O.O.; Akpan, I.; Oluwafemi, F. (2006). The potential of some essential oils in the control of *A. parasiticus* CFR 223 and aflatoxin production. *Food Cont.*, 18, 601-607.
- Basilico, M.Z.; Basilico, J.C. (1999). Inhibitory effects of some spices essential oils on *Aspergillus ochraceus* NRRL 3174 growth and ochratoxin A production. *Lett. Appl. Microbiol.*, 29, 238-241.
- Battilani, P.; Magan, N.; Logrieco, A. (2006). European research on ochratoxin A in grapes and wine. *Int J. Food Microbiol.*, 11, S2-S4.
- Bauer, A.W.; Kirby, W.M.M. (1966). Antibiotic susceptibility testing by standardized single disk method. *Am. J. Clin. Pathol.*, 45, 493-496.
- Baydar, H.; Sagdiç, O.; Ozkan, G.; Karadogan, T. (2004). Antibacterial activity and composition of essential oils from *Origanum*, *Thymbra* and *Satureja* species with commercial importance in Turkey. *Food Cont.*, 15, 169-172.
- Brul, S.; Coote, P. (1999). Preservative agents in foods: mode of action and microbial resistance mechanisms. *Int. J. Food Microbiol.*, 50, 1-17.
- Burt, S. (2004). Essential oils: their antibacterial properties and potential applications in foods – a review. *Int. J. Food Microbiol.*, 94, 223-253.
- Canuto, M.M.; Rodero, F.G. (2002). Antifungal drug resistance to azoles and polyenes. *Lancet. Infect. Dis.*, 2, 550-563.
- Chakrabarti, A.; Marak, R.S.K.; Sing, S.; Gupta, S.O.; Hurst, S.E.; Padhye, A.A. (2006). Brain abcess due to *Aspergillus nidulans*. *J. Med. Mycol.*, 16, 100-104.
- Chamilos, G.; Kontoyianis, D.P. (2005). Update on antifungal drug resistance mechanisms of *Aspergillus fumigatus*. *Drug Res. Update*, 8, 344-358.
- Curtis, L.; Conroy, L.; Coli, S.; Baker, K.; Our, C.H.; Hershow, R.; Norlock-Cruz, F.; Scheff, P. (2005). *Aspergillus* surveillance project at a large tertiary-care hospital. *J. Hosp. Infect.*, 59, 188-196.
- Daferera, D.J.; Ziogas, B.N.; Polissiou, M.G. (2003). The effectiveness of plant essential oils on the growth of *Botrytis cinerea*, *Fusarium* sp. and *Clavibacter michiganensis* subsp. *michiganensis*. *Crop. Protect.*, 22, 39-34.
- de Billerberck, V.G.; Roques, C.R.; Bessière, J.M.; Fonvieille, J.L.; Dargent, R. (2001). Effect of *Cymbopogon nardus* (L.) W. Watson essential oil on the growth and morphogenesis of *Aspergillus niger*. *Can. J. Microbiol.*, 47, 9-17.
- Desselberger, V. (2000). Emerging and re-emerging infectious disease. *J. Inf. Dis.*, 40, 3-15.
- Duarte, M.C.T.; Figueira, G.M.; Sartoratto, A.; Rehder, V.L.G.; Delarmelina, C. (2005). Anti-*Candida* activity of Brazilian medicinal plants. *J. Ethnopharm.*, 97, 305-311.
- Dubey, A.; Patwardhan, R.V.; Samph, S.; Santoshi, V.; Kolluri, S.; Nanda, A. (2006). Intracranial fungal granuloma: analysis of 40 patients and review of the literature. *Surg. Neurol.*, 63, 254-260.
- Gayoso, C.W.; Lima, E.O.; Trajano, V.N.; Pereira, F.O.; Souza, E.L.; Lima, I.O.; Navarro, D.F. (2005). Sensitivity of fungi isolated from onychomycosis to *Eugenia cariophyllata* essential oil and eugenol. *Fitoterapia*, 76, 247-249.
- Georgopapadakou, N.H. (2002). Infectious diseases 2001: drug resistance, new drugs. *Drug Res.*, 5, 181-191.

22. Hadacek, F.; Greger, H. (2000). Testing of antifungal natural products: methodologies, comparability of results and assay choice. *Phytochem. Anal.*, 11, 137-148.
23. Kawashima, L.M.; Valente, S.L.M. (2006). Incidência de fumonisina B₁, aflatoxinas B₁, B₂, G₁ e G₂, ocratoxina A e zearalenona em produtos de milho. *Cienc. Tecnol. Alim.*, 26, 516-521.
24. Lima, I.O.; Oliveira, R.A.G.; Lima, E.O.; Souza, E.L.; Farias, N.P.; Navarro, D.F. (2005). Inhibitory action of some phytochemicals on yeasts potentially causing of opportunistic infections. *Rev. Bras. Cien. Farm.*, 41, 199-203.
25. Lopez Díaz, T.M.L.; González, C.J.; Moreno, B.; Otero, A. (2002). Effect of temperature, water activity, pH and some antimicrobials on the growth of *Penicillium olsonii* isolated from the surface of Spanish fermented meat sausage. *Food Microbiol.*, 19, 1-7.
26. Marín, S.; Velluti, A.; Ramos, A.J.; Sanchis, V. (2004). Effect of essential oils on zearalenone and deoxynivalenol production by *Fusarium graminearum* in non-sterilized maize grain. *Food Microbiol.*, 21, 313-318.
27. Mishra, N.; Upma, K.; Shukla, D. (2000). Antifungal activity of essential oil of *Cinnamomum zeylanicum*. *J. Essent. Oil Res.*, 3, 97-110.
28. Moreira, A.C.P.; Lima, E.O.; Souza, E.L.; Van Dingenen, M.A.; Trajano, V.N. (2007). Inhibitory effect of *Cinnamomum zeylanicum* Blume (Lauraceae) essential oil and β-pinene on the growth of dematiaceous moulds. *Braz. J. Microbiol.*, 38, 33-38.
29. Nostro, A.; Blanco, A.R.; Cannatelle, M.A.; Enea, V.; Flamini, G.; Morelli, I.; Roccaro, A.S.; Alonso, V. (2004). Susceptibility of methicillin-resistant staphylococci to oregano essential oil, carvacrol and thymol. *FEMS Microbiol. Lett.*, 230, 191-195.
30. Pasquier, F.; Croxo, C.; Melliez, H.; Porte, H.; Bourgeois-Petit, E.; Cambier, N.; Rose, S. (2006). L'aspergillome pulmonaire: une complication possible de la drepanocytose. *La Revue de Medicine Interne*, 27, 260-263.
31. Pattnaik, S.; Subramanyan, V.R.; Kole, C. (1996). Antibacterial antifungal activity of ten essential oils in vitro. *Microbios*, 86, 121-126.
32. Rana, B.K.; Singh, U.P.; Taneja, V. (1997). Antifungal activity and kinetics of inhibition by essential oil isolated from leaves of *Aegle marmelos*. *J. Ethnopharm.*, 57, 29-34.
33. Rasooli, I.; Abyaneh, M.R. (2004). Inhibitory effect of Thyme oils on growth and aflatoxin production by *Aspergillus parasiticus*. *Food Cont.*, 15, 479-483.
34. Rasooli, I.; Owlia, P. (2005). Chemoprevention by thyme oils of *Aspergillus parasiticus* growth and aflatoxin production. *Phytochemistry*, 66, 2851-2856.
35. Rasooli, I.; Rezaei, M.B.; Allameh, A. (2006). Growth inhibition and morphological alterations of *Aspergillus niger* by essential oils from *Thymus eriocalyx* and *Thymus x-porlock*. *Food Conti.*, 17, 359-364.
36. Sahin, F.; Gulluce, M.; Daferera, D.; Sokmen, A.; Polissiou, M.; Agar, G.; Ozer, H. (2004). Biological activities of the essential oils and methanol extract of *Origanum vulgare* ssp. *vulgare* in the Eastern Anatolia region of Turkey. *Food Cont.*, 15, 549-557.
37. Saleemulla, A.I.; Khalil, I.A.; Shah, H. (2006). Aflatoxin contents of stored and artificially inoculated cereals and nuts. *Food Chem.*, 98, 699-703.
38. Sharma, N.; Tripathi, A. (2006). Effects of Citrus (L.) Osbeck epicarp essential oil on growth and morphogenesis of *Aspergillus niger* (L.) Van Tieghem. *Microbiol. Res.*, In press, 2007.
39. Shukla, A.C.; Shahi, S.K.; Dixit, A. (2000). Epicarp of *Citrus sinensis*: a potential source of natural pesticides. *Ind. Phytopath.*, 53, 468-471.
40. Soliman, K.M.; Badeae, R.I. (2002). Effect of oil extracted from some medicinal plants on different mycotoxicogenic fungi. *Food Chem. Toxicol.*, 44, 1669-1675.
41. Souza, E.L.; Lima, E.O.; Freire, K.R.L.; Sousa, C.P. (2005). Inhibition action of some essential oils and phytochemicals on the growth of moulds isolated from foods. *Braz. Arch. Biol. Technol.*, 48, 245-250.
42. Souza, E.L.; Stamford, T.L.M.; Lima, E.O.; Trajano, V.N. (2007). Effectiveness of *Origanum vulgare* L. essential oil to inhibit the growth of food spoiling yeasts. *Food Cont.*, 18, 409-413.
43. Thyagaraja, N.; Hosono, A. (1996). Effect of spice extract on fungal inhibition. *Lebensm-Wiss u-Technol.*, 29, 286-288.
44. Velluti, A.; Sanchis, V.; Ramos, A.J.; Egido, J.; Marin, S. (2003). Inhibitory effect of cinnamon, clove, lemongrass, oregano and palmarosa essential oils on growth and fumonisin B₁ production by *Fusarium proliferatum* in maize grain. *Int. J. Food Microbiol.*, 89, 145-154.
45. Zambonelli, A.; Zechini d'Aulerio, A.; Bianchi, A.; Albasini, A. (1996). Effects of essential oil on phytopathogenic fungi. *Phytopath.*, 144, 491-494.
46. Zygallo, J.A.; Grossi, N.R. (1995). Comparative study of the antifungal activity of essential oils from aromatic plants growing wild in the central region of Argentina. *Flavor and Fragrance Journal.*, 10, 113-118.