



Fungal and Bacterial Physiology

Influence of iron and copper on the activity of laccases in *Fusarium oxysporum* f. sp. *lycopersici*



Wendy Susana Hernández-Monjaraz, César Caudillo-Pérez,
Pedro Ulises Salazar-Sánchez, Karla Lizbeth Macías-Sánchez*

Instituto Politécnico Nacional, Unidad Profesional Interdisciplinaria de Ingeniería campus Guanajuato, Silao de la Victoria, Guanajuato, Mexico

ARTICLE INFO

Article history:

Received 28 October 2017

Accepted 22 June 2018

Available online 13 August 2018

Associate Editor: Jerri Zilli

Keywords:

Chelators

Cooper

Fusarium

Iron

Laccases

ABSTRACT

Fusarium oxysporum f. sp. *lycopersici* is a phytopathogenic fungus that causes vascular wilt in tomato plants. In this work we analyze the influence of metal salts such as iron and copper sulphate, as well as that of bathophenanthrolinedisulfonic acid (iron chelator) and bathocuproinedisulfonic acid (copper chelator) on the activity of laccases in the intra (IF) and extracellular fractions (EF) of the wild-type and the non-pathogenic mutant strain (*rho1::hyg*) of *F. oxysporum*. The results show that laccase activity in the IF fraction of the wild and mutant strain increased with the addition of iron chelator (53.4 and 114.32%; respectively). With copper, it is observed that there is an inhibition of the activity with the addition of CuSO_4 for the EF of the wild and mutant strain (reduction of 82 and 62.6%; respectively) and for the IF of the mutant strain (54.8%). With the copper chelator a less laccase activity in the IF of the mutant strain was observed (reduction of 53.9%). The results obtained suggest a different regulation of intracellular laccases in the mutant strain compared with the wild type in presence of CuSO_4 and copper chelator which may be due to the mutation in the *rho* gene.

© 2018 Sociedade Brasileira de Microbiologia. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Fusarium oxysporum is a highly relevant phytopathogenic fungus due to its economic importance.¹⁵ The special form of *lycopersici* is the causal agent of vascular fusariosis disease in tomato culture. This fungus reaches the vascular system of

the plant and subsequently achieves the colonization of the host plant by spreading through the xylem vessels.²⁸

Laccases (benzenediol: oxygen oxidoreductase, EC 1.10.3.2) are a type of polyphenoloxidase enzymes that are part of the ligninolytic system and with ascorbate oxidases, ceruplasmin and other enzymes belong among blue multicopper oxidases.^{23,1} The participation of laccases is associated with

* Corresponding author.

E-mail: kmaciass@ipn.mx (K.L. Macías-Sánchez).

<https://doi.org/10.1016/j.bjm.2018.06.002>

1517-8382/© 2018 Sociedade Brasileira de Microbiologia. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

many biological processes, such as spore resistance, pigmentation, virulence factors,^{2,37} lignification of plant cell walls,²⁰ lignin biodegradation⁵ and the protection of the fungus against phenolic compounds released by plants, such as phytoalexins,²⁴ among others. It is known that the manifestation of laccases in fungi is influenced by several factors, including the presence of metals.²⁵ Copper has been demonstrated to regulate the induction transcriptional of the laccases in *Trametes versicolor* and *Trametes pubescens*,^{11,14} *Pleurotus*,^{32,21} among others.

On the other hand, FeSO₄ at a concentration of 1 mM did not induce laccase activity with respect to control in *Pleurotus pulmonarius* at 7 days of growth but there was an inhibition of activity at 10 days of growth. A similar response was observed in *Pleurotus ostreatus*, since there was an inhibition of activity at 10 days, but in this case there a stimulatory effect of activity was observed at 7 days of growth.³³ It was obtained a complete inhibition of activity of a purified laccase of the fungus *Trematosphaeria mangrovei* with FeSO₄ at 1 mM.¹⁷

In *F. oxysporum* f. sp. *lycopersici* the presence of six genes encoding laccase proteins (*lcc1*, *lcc2*, *lcc3*, *lcc4*, *lcc5* and *lcc9*) has been reported. The mutants *lcc1*, *lcc3* and *lcc5* were implemented, in which a significant decrease in the extracellular activity of laccase was observed; however, these mutants showed no loss of pathogenicity on tomato seedlings.⁸

To know the effect that some metals and chelants may have on the activity of laccases in *F. oxysporum*, we analyzed the effect of FeSO₄, the iron chelator bathophenanthrolinedisulfonic acid (BPDS), CuSO₄ and the copper chelator bathocuproinedisulfonic acid (BCDS) on the enzymatic activity of laccase in a wild strain and the mutant *rho1::hyg* (non-pathogenic) of *F. oxysporum* f. sp. *lycopersici*.

Materials and methods

Isolated fungi and growing conditions

Strain 4287 (race 2; wild-type) and the mutant strain *rho1::hyg* of *F. oxysporum* f. sp. *lycopersici* were obtained through Dr. Gonzalez Roncero (University of Cordoba, Spain). The mutant *rho1::hyg* has interrupted *rho1* gene (*rho1::hyg*) and presents loss of the pathogenicity in tomato seedlings.¹⁸

To preserve the strains, the wild-type strain was inoculated in the PDB medium and the mutant strain in the PDB medium with hygromycin (20 µg/mL; SIGMA, Saint Louis, Missouri, USA) for 4–5 days at 27 °C with shaking. The generated mycelia were removed through filtration under sterile conditions. The filtrate obtained was centrifuged at 6500 g for 10 min, and the sediment containing the spores was washed three times with sterile water. The strains were stored as microconidial suspensions in 30% glycerol at 70 °C until its usage.

For different growth conditions, the germinates were obtained from the PDB culture, which was inoculated with 5×10^6 microconidia mL⁻¹ for 19 h at 27 °C under shaking. The germs were transferred to a basal medium (BL) (0.4 g KH₂PO₄ L⁻¹, 0.2 g MgSO₄·7H₂O L⁻¹, 1 g NH₄NO₃ L⁻¹, 0.01 g FeSO₄ L⁻¹, 0.01 g ZnSO₄ L⁻¹ and 0.01 g MnSO₄ L⁻¹, 0.2 g KCl L⁻¹, 0.25% sucrose) for 6 days; BL without copper (incubation for 3 days)

and BL without iron (incubation for 3 days) at 27 °C, 220 rpm. On the third day of growth in the BL medium without copper, the copper chelator bathocuproinedisulfonic acid (BCDS; SIGMA, Saint Louis, Missouri, USA) at 1 mM and CuSO₄ at 25 µM was independently added in flasks. The same procedure is applied for the BL medium without iron, in which the iron chelator, bathophenanthrolinedisulfonic acid (BPDS; SIGMA, Saint Louis, Missouri, USA) was added at 1 mM; and for the duplicate of the BL medium without iron, FeSO₄ was added at 25 µM. After the addition of BCDS, CuSO₄, BPDS and FeSO₄, the incubation was continued for 3 days at 220 rpm at 27 °C. The mycelium and the extracellular fraction were obtained by centrifugation at 6500 g for 10 min at 4 °C.

Obtainment of the homogenate

The mycelia were resuspended in a Buffer Tris-HCl 50 mM solution with protease inhibitors (antipain, leupeptin, pepstatin and E64 at 5 µg/mL; SIGMA, Saint Louis, Missouri, USA). The cells were fragmented through sonication for 8–10 min for 1-min intervals. Subsequently an initial centrifugation was performed at 2500 × g for 5 min at a temperature of 4 °C. The pellet was discarded and the supernatant transferred to a new tube, where a second centrifugation was performed at 6500 × g for 45 min at 4 °C, the pellet discarded and the supernatant (intracellular fraction) obtained. The aforementioned protease inhibitors were added to the extracellular fraction. Both fractions were stored at –20 °C until its usage.

Quantification of enzyme activity

The laccase activity was determined using 2,2'-azino-bis(3-ethylbenzothiazoline-6-sulfonic acid) [ABTS; SIGMA, Saint Louis, Missouri, USA] as a substrate.⁷ The laccase activity was quantified through the oxidation of the ABTS substrate (1 mM) in 20 mM sodium acetate pH 3.5 buffer, by increasing the absorbance at 436 nm.¹⁴ Final reaction mixture contained 100 µL of the sample and 2 mL of the ABTS substrate,¹⁶ measuring the reaction at 0 and 10 min. A unit of laccase activity is defined as the amount of enzyme needed to oxidize a µmol of ABTS min⁻¹ at 25 °C.¹⁴ All assays were performed in triplicate and three independent batches were performed.

Statistic analysis

The data on enzyme activity were compared between the two strains using ANOVA and Tukey's tests, with Minitab ver. 15 software (Minitab Inc., State College, PA, USA), considering a level of significance at 0.05.

Results

Influence of iron on the enzymatic activity of laccases

From three independent batches, the enzymatic activity of laccases was measured in the intracellular and extracellular fraction of the wild and mutant strain *F. oxysporum* f. sp. *lycopersici*, in the presence of iron and the BPDS iron chelator. Through the ANOVA test, the mutant strain *rho1::hyg*

Table 1 – Relation between the activity in the medium of the *rho1::hyg* mutant and in that of the wild-type strain in the control and in the presence of BPDS and FeSO₄.

| Fraction | Condition | <i>rho1::hyg</i> activity/wild-type activity |
|---------------|-------------------|--|
| Intracellular | Control | 2.035218 |
| | BPDS | 2.813707 |
| | FeSO ₄ | 3.288776 |
| Extracellular | Control | 2.246674 |
| | BPDS | 2.162287 |
| | FeSO ₄ | 2.022699 |

demonstrated higher extracellular and intracellular enzyme activity (in the control condition, in the presence of BPDS and FeSO₄) compared to the wild-type strain. With a normal distribution of the data (p -value > 0.1) it is considered that the activity between both strains will always be greater for the mutant strain under the same conditions of the experiment (Table 1).

According to the statistical analysis through the Tukey method, which allows us to analyze the difference in enzymatic activity between the conditions studied for each strain (control group, in the presence of BPDS chelator and in the presence of FeSO₄), it was observed that in the extracellular fraction of the wild strain and the mutant strain there exists no difference between the enzymatic activity of the control group with respect to the group in the presence of BPDS or the group in the presence of FeSO₄. With respect to the enzymatic activity of the intracellular fraction, in both strains (wild and mutant) a bigger activity in the group with the BPDS chelator was observed with respect to control group (53.4 and 114.32%, respectively), whereas no difference of intracellular enzymatic activity was observed in both strains between the control group and the group with FeSO₄ (Fig. 1).

Influence of copper on the enzymatic activity of laccase

From three independent batches, the enzymatic activity of laccases was measured in the intracellular and extracellular




Fig. 1 – Enzymatic activity in the extracellular and intracellular fractions of wild-type and *rho1::hyg* of *F. oxysporum* f. sp. *lycopersici* strains with or without FeSO₄ and with BPDS. The data were grouped by the results of Tukey's test using a confidence level of 95%.

Table 2 – Relation between the activity in the medium of the *rho1::hyg* mutant and in that of the wild-type strain in the control and in the presence of BCDS and CuSO₄.

| Fraction | Condition | <i>rho1::hyg</i> activity/wild-type activity |
|---------------|-------------------|--|
| Intracellular | Control | 5.144615 |
| | BCDS | 2.424381 |
| | CuSO ₄ | 1.850917 |
| Extracellular | Control | 6.184008 |
| | BCDS | 6.805616 |
| | CuSO ₄ | 12.58875 |

fraction of the wild and mutant strain of *F. oxysporum* f. sp. *lycopersici*, in the presence of copper and the BCDS copper chelator. In both fractions, the ANOVA test performed on all the lots allowed us to know that the mutant strain *rho1::hyg* had a higher enzymatic activity for the different growth conditions compared to the wild strain. The data maintained a normal distribution (p -value > 0.1) (Table 2).

According to the statistical analysis performed through the Tukey method, it was observed that in the extracellular fraction, the enzymatic activity in the wild and mutant strain presented less activity in the group with CuSO₄ compared to their respective control groups (a reduction of the activity of 82 and 62.6% with respect the control group, respectively). There was no effect between the control group and the group with addition of the BCDS chelator of the extracellular fraction (in both strains). On the other hand, in the intracellular fraction of the wild-type strain there was no difference in laccase activity between the control group, the group with CuSO₄ and the group with addition of the BCDS chelator; however, in the mutant strain the CuSO₄ group demonstrated less activity with respect control group (a reduction of the activity of 54.8%). Also in the same fraction, the mutant strain showed less activity of laccase in the BCDS chelator group compared with control group (a reduction of 53.9%) (Fig. 2).




Fig. 2 – Enzymatic activity in the extracellular and intracellular fractions of wild-type and *rho1::hyg* of *F. oxysporum* f. sp. *lycopersici* strains with or without CuSO₄ and with BCDS. The data were grouped by the results of Tukey's test using a confidence level of 95%.




Fig. 3 – Oxidation of ABTS by laccase. ABTS cation radical (a) and ABTS dication (b).²²

Discussion

Laccases are a type of phenoloxidase enzymes widely distributed in nature, as they have been found in plants, bacteria and fungi.¹⁹ Among the functions attributed to laccases are lignification,²⁰ the protection of the fungus against phenolic compounds,²⁴ among others. ABTS is an organic redox mediator used in the laccase reaction. ABTS undergoes two step oxidation reaction during the enzymatic reaction by laccase (first for the obtention of ABTS cation radical and then the ABTS dication)²² (Fig. 3).

In studies by Cañero and Roncero in 2008⁸ demonstrated the expression of various laccase genes during the infection process of the tomato plant with *F. oxysporum* f. sp. *lycopersici* (*lcc1*, *lcc3* and *lcc9*). In addition, the location of each of

the laccase proteins through the amino acid sequence was identified, demonstrating that *Lcc1* and *Lcc2* have an intracellular localization, *Lcc3* is transmembrane and *Lcc4*, *Lcc5* and *Lcc9* have an extracellular localization. An analysis of the promoter regions of the *lcc1*, *lcc2*, *lcc3*, *lcc4*, *lcc5* and *lcc9* genes demonstrate metal-responsive elements (MRE), stress-responsive elements (STRE), xenobiotic-responsive elements (XRE), an ACE1 copper-responsive transcription factor and a PacC ambient pH response factor. Laccase activity has been reported to increase with low concentrations of sucrose, glycerol and polygalacturonic acid.⁸ Therefore, in this work a basal medium with a low concentration of sucrose as a carbon source was used.

Some trace elements are essential for fungal metabolism; however, at high concentrations they could be toxic.^{6,4} Some trace elements such as Fe, Zn and others act through metal-responsive promoter interaction, for post-transcriptional regulation.^{34,13,10} In fungi, it is known that regulation in the manifestation of laccases can be mediated by the presence of various metals.²⁵

In the results obtained in this work, a higher enzymatic activity of laccase was observed in the *rho1::hyg* mutant strain compared to the wild-type strain under all tested conditions (control group, presence of iron, copper, and BPDS and BCDS chelators). These results correspond with to the reported by Reyes-Medina and Macías-Sánchez,²⁶ since under the conditions utilized in that work they also detected greater activity in the mutant strain with respect to the wild-type, suggesting that this effect could be influenced by the absence of Rho protein. The results obtained at the extracellular and intracellular levels suggest that the presence of FeSO₄ has no apparent effect on the enzymatic activity of laccase in the wild strain of *F. oxysporum* f. sp. *lycopersici* in comparison to its basal condition, which could be due to FeSO₄ not acting as an inducer of the extracellular laccase (*Lcc4*, *Lcc5* and *Lcc9*) or intracellular laccases (*Lcc1* and *Lcc2*), or that the concentration used (25 μM) is not the concentration necessary to generate the induction of laccases. A similar behavior is observed for the mutant strain, where the addition of FeSO₄ does not appear to affect laccase activity. The effect of Fe on laccase activity differs depending on the microorganism and the concentration used; for example, the effect of this metal at a concentration of 2 mM on the activity of the purified laccase on *Aspergillus nidulans* demonstrated a relative activity of 139% with respect to that obtained in the control condition.³⁶ However, in *Streptomyces cyaneus* a reduction of the activity of laccase purified under the presence of FeSO₄ (concentrations since 0–50 mM) was observed.³ On the other hand,³⁸ observed in *Trametes velutina* an increase in the extracellular enzymatic activity when adding iron from a concentration of 0.04 mM.²⁷ tested the effect of FeSO₄ at a concentration of 5 mM on laccase activity in *Chalara* (syn. *Thielaviopsis*) *paradoxa*, observing an inhibition of the activity with respect to the control.

In *P. ostreatus* an increase in laccase activity was observed in the presence of FeSO₄ at 7 days of incubation; however, at 10 days of incubation there was less activity with respect to its control. On the other hand, the activity of laccase in *Pleurotus pulmonarius* was similar in presence of FeSO₄ and the control group at 7 days of incubation,

and there was a decrease of the activity in the presence of Fe compared to the control group at 10 days of incubation.³³

The BPDS is known as a potent iron chelating agent.³¹ In the present study, no differences were observed between the control group and the group with addition of BPDS in the extracellular fraction of the wild and mutant strain. In contrast, in the intracellular fraction, there was observed that in both strains the BPDS group presented greater activity compared to the control group, which would indicate that a lower iron concentration is required than the used in the control condition to induce the activity of the laccases.

Copper plays an important role in the regulation of laccases since it is known that the addition of low concentrations of copper to the culture medium can stimulate laccase production²¹; in addition, laccases have four copper atoms in their catalytic center, which have different properties.²³ With the data obtained in this work, in the group with the addition of CuSO₄, indicate a decrease in the activity of laccase in the extracellular fraction of the wild and mutant strain, as well as in the intracellular fraction of the mutant strain, indicating that CuSO₄ at a concentration of 25 μM inhibits the activity of laccases in *F. oxysporum* f. sp. *lycopersici*.¹¹ reported that an increase in the concentration of copper causes an increment in the laccases activity in *T. versicolor*. However, the use of different concentrations of the metal on the induction of laccases in other microorganisms has demonstrated that at a certain concentration a maximum is reached in the activity and after this the activity considerably decreases and in some cases it is canceled completely. In the *Grammothele subargentea* fungus, it was observed that in the concentration range of 0.6–1.2 mM of CuSO₄ major enzymatic activity of laccase occurred, whereas at higher concentrations of the metal (1.5 and 1.8 mM), the activity decreases³⁰; a similar situation is observed in *T. versicolor*, where the laccase activity is induced in the presence of CuSO₄ up to 17%; however, it decreases significantly with high levels of Cu²⁺ (80 mM).¹⁶ A particular case is that of *Trametes hirsuta*, in which there was observed the irregular behavior of enzymatic activity under induction conditions with CuSO₄ with respect to the culture time and the used concentration of the metal; in that experiment being possible to distinguish several peaks of activity, followed by periods of inhibition.²⁹ In this work, there is no difference in the activity of the laccase of the control group and the group with copper in the intracellular fraction of the wild strain, where possibly a higher concentration of copper is required to observe an effect on the enzymatic activity of the laccases. This also indicates that the laccases present in the intracellular fraction of the wild strain do not present as much sensitivity to copper, unlike the laccases present in the intracellular fraction of the mutant strain, and in the extracellular fraction (wild and mutant strain).

BCDS is a chelating agent for copper ions.³⁵ Endo et al.¹² used a chelator concentration similar to this work. They observed a reduction of the enzymatic activity between the condition and its control. With the results obtained in our work, in the presence of the BCDS chelator in the extracellular fraction, no differences were observed with respect to the control for the two strains utilized, as well as in the

intracellular fraction of the wild strain, not being the same case for the mutant strain, where there is a diminishing in the activity in the group with the addition of BCDS, being the only case where the inhibition of the enzymatic activity is observed with the presence of this chelator.⁹ analyzed the influence of copper in the expression of laccase genes in *F. oxysporum* utilizing CuSO₄ and BCDS; in both cases there were seen no differences in the expression of the *lcc1*, *lcc2*, *lcc3*, *lcc4*, *lcc5* and *lcc9* genes; however, if there was a higher extracellular enzymatic activity in the wild-type strain in the presence of CuSO₄ compared to the medium without CuSO₄ (growth for 5 days in a minimal medium containing sucrose with and without 250 μM of CuSO₄), these results differ from those obtained in this work, since we observed less activity in the presence of copper, even though a smaller amount of CuSO₄ (25 μM) was utilized; however, the incubation was performed for a shorter period of time (3 days), where it is possible to observe a different behavior over time, which may be due to regulation mechanisms of the metal.

In this work, it was of interest to learn about the *in vitro* behavior of the laccase enzymes' activity in the intra and extracellular fraction of the wild strain of *F. oxysporum* f. sp. *lycopersici* (pathogen strain), and of the mutant strain *rho1::hyg* (non-pathogen strain), and its response to inductors like Fe and Cu, and with the iron chelator (bathophenanthrolinedisulfonic acid), and copper chelator (bathocuproinedisulfonic acid). Nevertheless, these do not constitute a field treatment for the control of fusariosis. However, knowledge of the behavior of laccase activity in the wild and mutant strains under these conditions may be used as a guide in the analysis and modulation of these metallic ions in the soil, thus creating soil conditions that can be less appropriate for the establishment of fusariosis.

With the results obtained in this work it can be suggested that the intracellular laccases of the wild and mutant strain demonstrate a differential response to the presence of copper ions. The difference observed could be regulated through of *rho1* gene. Because the role of copper in the function of laccase proteins, this response could be attributed to its insertion in the inactive apoprotein. As perspectives of this work, the conditions that presented a bigger effect on the laccase activity can be used to observe the effect that present in the sporulation, pigmentation, and expression of transcription factors, among other processes.

Funding

This study was funded by the Instituto Politécnico Nacional, Secretaría de Investigación y Posgrado (20131056).

Conflicts of interest

The authors declare that they have no conflicts of interest.

Acknowledgements

This research was supported by the Secretaría de Investigación de Posgrado of the Instituto Politécnico Nacional (20131056).

REFERENCES

- Alcalde M. Laccase: biological functions, molecular structure and industrial applications. In: Polaina J, MacCabe AP, eds. *Industrial Enzymes: Structure Function and Applications*. Netherlands: Springer; 2007:461–476.
- Aramayo R, Timberlake WE. The *Aspergillus nidulans* yA gene is regulated by abaA. *EMBO J*. 1993;12:2039–2048.
- Arias ME, Arenas M, Rodríguez J, Soliveri J, Ball AS, Hernández M. Kraft pulp biobleaching and mediated oxidation of a nonphenolic substrate by laccase from *Streptomyces cyaneus* CECT 3335. *Appl Environ Microbiol*. 2003;69:1953–1958.
- Baldrian P. Interactions of heavy metals white-rot fungi. *Enzyme Microb Technol*. 2003;32:78–91.
- Baldrian P. Fungal laccases – occurrence and properties. *FEMS Microbiol Rev*. 2006;30:215–242.
- Baldrian P, Gabriel J. Copper and cadmium increase laccase activity in *Pleurotus ostreatus*. *FEMS Microbiol Lett*. 2002;206:69–74.
- Bourbonnais R, Paice MG. Oxidation of non-phenolic substrates. An expanded role for laccase in lignin biodegradation. *FEBS Lett*. 1990;267:99–102.
- Cañero DC, Roncero MI. Functional analyses of laccase genes from *Fusarium oxysporum*. *Phytopathology*. 2008;98:509–518.
- Cañero DC, Roncero MI. Influence of the chloride channel of *Fusarium oxysporum* on extracellular laccase activity and virulence on tomato plants. *Microbiology*. 2008;154:1474–1481.
- Catal T, Liu H, Bermek H. Selenium induces manganese-dependent peroxidase production by the white-rot fungus *Bjerkandera adusta* (Willdenow) P. Karsten. *Biol Trace Elem Res*. 2008;123:211–217.
- Collins PJ, Dobson ADW. Regulation of laccase gene transcription in *Trametes versicolor*. *Applied Environ Microbiol*. 1997;63:3444–3450.
- Endo K, Hayasho Y, Hibi T, Hosono K, Beppu T, Ueda K. Enzymological characterization of EpoA, a laccase-like phenol oxidase produced by *Streptomyces griseus*. *J Biochem*. 2003;133:671–677.
- Faraco V, Giardina P, Sannia G. Metal-responsive elements in *Pleurotus ostreatus* laccase gene promoters. *Microbiology*. 2003;149:2155–2162.
- Galhaup C, Goller S, Peterbauer CK, Strauss J, Haltrich D. Characterization of the major laccase isoenzyme from *Trametes pubescens* and regulation of its synthesis by metal ions. *Microbiology*. 2002;148:2159–2169.
- Geiser DM, Aoki T, Bacon CW, et al. One fungus, one name: defining the genus *Fusarium* in a scientifically robust way that preserves longstanding use. *Phytopathology*. 2013;103:400–408.
- Lorenzo M, Moldes D, Rodríguez CS, Sanromán MA. Inhibition of laccase activity from *Trametes versicolor* by heavy metals and organic compounds. *Chemosphere*. 2005;60:1124–1128.
- Mabrouk MA, Kheiralla HZ, Hamed RE, Youssry AA, Abd A. Characterization and kinetic properties of the purified *Trematosphaeria mangrovei* laccase enzyme. *Saudi J Biol Sci*. 2013;20:373–381.
- Martínez-Rocha AL, Roncero MI, López-Ramírez A, et al. Rho 1 has distinct functions in morphogenesis, cell wall biosynthesis and virulence of *Fusarium oxysporum*. *Cell Microbiol*. 2008;10:1339–1351.
- Mayer AM, Staples RC. Laccase: new functions for an old enzyme. *Phytochemistry*. 2002;60:551–565.
- O'Malley DM, Whetten R, Bao W, Chen CL, Seedorf RR. The role of laccase in lignification. *Plant J*. 1993;4:751–757.
- Palmieri G, Giardina P, Bianco C, Fontanella B, Sannia G. Copper induction of laccase isozymes in the ligninolytic fungus *Pleurotus ostreatus*. *Appl Environ Microbiol*. 2000;66:920–924.
- Pankaj KC, Rama SS, Sudha Y. A review on mechanism of laccase action. *Res Rev Biosci*. 2013;7:66–71.
- Pannu JS, Kapoor RK. Microbial laccases: a mini-review on their production, purification and applications. *Int J Pharm Biol Arch*. 2014;3:528–536.
- Pezet R, Pont V, Hoang-Van K. Enzymatic detoxication of stilbenes by *Botrytis cinerea* and inhibition by grape berries proanthocyanidins. In: Verhoeff K, Malathrakis NE, Williamson B, eds. *Recent Advances in Botrytis, Research*. Wageningen: Pudoc Scientific; 1992:87–92.
- Piscitelli A, Giardina P, Lettera V, Pezzella C, Sannia G, Faraco V. Induction and transcriptional regulation of laccases in fungi. *Curr Genom*. 2011;12:104–112.
- Reyes-Medina MA, Macías-Sánchez KL. GTPase Rho1 regulates the expression of xyl3 and laccase genes in *Fusarium oxysporum*. *Biotechnol Lett*. 2015;37:679–683.
- Robles A, Lucas R, Martínez-Cañamero M, Ben Omar N, Pérez R, Gálvez A. Characterisation of laccase activity produced by the hyphomycete *Chalara* (syn *Thielaviopsis*) *paradoxa* CH32. *Enzyme Microb Technol*. 2002;31:516–522.
- Roncero MIG, Hera C, Ruiz-Rubio M, et al. *Fusarium* as a model for studying virulence in soilborne plant pathogens. *Physiol Mol Plant Pathol*. 2003;62:87–98.
- Rosales E, Rodríguez S, Sanromán MA. Increased laccase production by *Trametes hirsuta* grown on ground orange peelings. *Enzyme Microb Technol*. 2007;40:1286–1290.
- Saparrat MCN. Optimizing production of extracellular laccase from *Grammothele subargentea* CLPS no. 436 strain. *World J Microbiol Biotechnol*. 2004;20:583–586.
- Severina IS, Bussygina OG, Pyatakova NV, Malenkova IV, Vanin AF. Activation of soluble guanylate cyclase by NO donors-S-nitrosothiols, and dinitrosyl-iron complexes with thiol-containing ligands. *Nitric Oxide*. 2003;8:155–163.
- Soden DM, Dobson ADW. Differential regulation of laccase gene expression in *Pleurotus sajor-caju*. *Microbiology*. 2001;147:1755–1763.
- Stajić M, Vukojević J, Knežević A, Milovanović I. Influence of trace elements on ligninolytic enzyme activity of *Pleurotus ostreatus* and *P. pulmonarius*. *Bioresources*. 2013;8:3027–3037.
- Thiele DJ. Metal-regulated transcription in eukaryotes. *Nucleic Acids Res*. 1992;20:1183–1191.
- Upadhyay S, Torres G, Lin X. Laccases involved in 1,8-dihydroxynaphthalene melanin biosynthesis in *Aspergillus fumigatus* are regulated by developmental factors and copper homeostasis. *Eukaryotic Cell*. 2013;12:1641–1652.

-
36. Vivekanandan KE, Sivaraj S, Kumaresan S. Characterization and purification of laccase enzyme from *Aspergillus nidulans* CASVK3 from vellar estuary south east coast of India. *Int J Curr Microbiol App Sci*. 2014;3:213–227.
 37. Williamson PR, Wakamatsu K, Ito S. Melanin biosynthesis in *Cryptococcus neoformans*. *J Bacteriol*. 1998;180:1570–1572.
 38. Yang Y, Wei F, Zhuo R, et al. Enhancing the laccase production and laccase gene expression in the white-rot fungus *Trametes velutina* 5930 with great potential for biotechnological applications by different metal ions and aromatic compounds. *PLoS One*. 2013;8:e79307.