



Short communication

Draft genome sequence of phenol degrading *Acinetobacter* sp. Strain V2, isolated from oil contaminated soil

Vikas Sharma, Johnson Lin*

University of KwaZulu-Natal (Westville), School of Life Sciences, Department of Microbiology, Durban, South Africa

ARTICLE INFO

Article history:

Available online 23 November 2016

Associate Editor: John McCulloch

Keywords:

Acinetobacter

Phenol degradation

Whole genome sequencing

Oil contaminated soil

ABSTRACT

We report here the draft genome sequence of *Acinetobacter* sp. Strain V2 isolated from the oil contaminated soil collected from ENGEN, Amanzimtoti, South Africa. Degradation of phenolic compounds such as phenol, toluene, aniline etc. at 400 ppm in 24 h and oil degrading capability makes this organism an efficient multifunctional bioremediator. Genome sequencing of *Acinetobacter* spp. V2 was carried out on Illumina HiSeq 2000 platform (performed by the Beijing Genomics Institute [BGI], Shenzhen, China). The data obtained revealed 643 contigs with genome size of 4.0 Mb and G + C content of 38.59%.

© 2016 Sociedade Brasileira de Microbiologia. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Genome announcement

Acinetobacter spp. have been involved in bioremediation of various pollutants such as phenols, benzoate, crude oil, acetonitrile etc.^{1,2} and biotechnological applications like production of extra-and-intracellular lipases, proteases, bio-emulsifiers and various types of biopolymers.^{3,4} Physiological and genetic characterisation of large number of phenol-degrading bacteria isolated from various sources have been done,^{5,6} however, little information on bacteria with a high phenol tolerance and high metabolising activity is available.⁷ Substrate inhibition and low degradation rate are the key factors which limits their applications.⁸ *Acinetobacter* sp. V2 strain is able to degrade 400 ppm of phenol within 24 h.⁹ Presence of commercially important proteins apart from its ability to

degrade diesel and engine oil makes this organism unique and thus genome sequencing.

Whole-genome sequencing was carried using Illumina HiSeq 2000 platform (Beijing Genomics Institute [BGI], Shenzhen, China) by generating paired-end libraries with an average insert size of 500bp following the manufacturer's instructions. The reads were then aligned with the reference sequence using SOAPaligner (version 2.21) software to calculate average depth and coverage ratio (<http://soap.genomics.org.cn/soapaligner.html>, version 2.21).¹⁰ The filtered short reads were first de novo assembled using SOAPdenovo v 2.04 according to the method described previously (<http://soap.genomics.org.cn/soapdenovo.html>),¹¹ and then contigs were manually connected according to their 500bp paired-end relationships. The draft genome sequence of *Acinetobacter* sp. V2 strain comprised of 643 contigs and

* Corresponding author at: Microbiology Discipline, School of Life Sciences, University of KwaZulu-Natal (Westville), Private Bag X54001, Durban 4000, South Africa.

E-mail: linj@ukzn.ac.za (J. Lin).

<http://dx.doi.org/10.1016/j.bjm.2016.06.015>

1517-8382/© 2016 Sociedade Brasileira de Microbiologia. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Table 1 – General features of the *Acinetobacter* sp. Strain V2 genome.

Sl. No.	Features	Values
1	Genome Size (bp)	4,007,850
2	DNA G + C content (%)	38.59
3	Total contigs	643
4	Total genes (CDS)	3717
5	rRNA	14
6	tRNA	71
7	Genes with predicted function	2971

16 scaffolds with the maximum contig size of 910,143 bp and scaffold size of 1,913,879 bp (Table 1). The genome size was 4,007,850 bp at 109.2 × coverage, with N₅₀ of 466,223 bp and N₉₀ of 97,271 bp and G+C content was 38.59%. A total of 3717 coding sequences (CDSs) or ORFs were predicted using Glimmer v3.02,¹² and homologous comparison to a non-redundant public database was performed by BLAST for function annotation. The genome annotation was performed using the (BASys) server (<https://www.basys.ca/>) and the output was downloaded in GenBank format resulting in 3742 (CDSs).¹³ The genome was further annotated with Rapid Annotation using Subsystems Technology (RAST) server (<http://rast.nmpdr.org/>).¹⁴ Among the predicted 3742 protein-coding genes by BASys, 79.4% (2971 genes) have been assigned putative functions according to the subsystem categorization. A total of 71 tRNA genes encompassing all 20 amino acids were identified using the tRNAscan-SE program,¹⁵ 14 rRNA genes were identified using RNAmmer¹⁶ and the insertion sequence (IS) elements were annotated by ISsaga.¹⁷ All the contigs were submitted to the Gene bank and NCBI has published sequence data in April 2015. The further analysis is going on.

Nucleotide sequence accession numbers: This WGS project has been deposited at DDBJ/EMBL/GenBank under the accession JZFB00000000. The version described in this paper is version JZFB01000000. Bioproject registered under accession: PRJNA275383 ID: 275383. The *Acinetobacter* sp. V2 isolate was deposited at the Leibniz Institute DSMZ-German Collection of Microorganisms and Cell Cultures and is available under the Accession No. DSM 101893.

REFERENCES

- Mandri T, Lin J. Isolation and characterization of engine oil degrading indigenous microorganisms in KwaZulu-Natal, South Africa. *Afr J Biotechnol.* 2007;6(1):23–27.

- Tabrez S, Ahmad M. Oxidative stress mediated genotoxicity of wastewaters collected from two different stations in northern India. *Mutat Res.* 2011;11:15–20.
- Saxena M, Gupta S, Mahmooduzzafar, Kumar R, Kumar A. Identification and genetic characterization of phenol-degrading bacterium isolated from oil contaminated soil. *Afr J Biotechnol.* 2013;12:791–797.
- Snellman EA, Colwell RR. *Acinetobacter* lipases: molecular biology, biochemical properties and biotechnological potential. *J Ind Microbiol Biotechnol.* 2004;31:391–400.
- Anli G, Ji LC. Proteome analysis of the adaptation of a phenol-degrading bacterium *Acinetobacter* sp. EDP3 to the variation of phenol loadings. *Chin J Chem Eng.* 2007;15:781–787.
- Zhan Y, Yan Y, Zhang W, et al. Comparative analysis of the complete genome of an *Acinetobacter calcoaceticus* strain adapted to a phenol-polluted environment. *Res Microbiol.* 2012;163:36–43.
- Wang Y, Tian Y, Han B, Zhaw HB, Bi JN, Cai BL. Biodegradation of phenol by free and immobilized *Acinetobacter* sp. strain PD12. *J Environ Sci.* 2007;19:222–225.
- Krastanov A, Alexieva Z, Yemendzhiev H. Microbial degradation of phenol and phenolic derivatives. *Eng Life Sci.* 2013;13:76–87.
- Lin J, Sharma V, Milase R, Mbhense N. Simultaneous enhancement of phenolic compound degradations by *Acinetobacter* strain V2 via a step-wise continuous acclimation process. *J Basic Microbiol.* 2015, <http://dx.doi.org/10.1002/jobm.201500263>.
- Li R, Yu C, Li Y, et al. SOAP2: an improved ultrafast tool for short read alignment. *Bioinformatics.* 2009;25:1966–1967.
- Li R, Li Y, Kristiansen K, Wang J. SOAP: short oligonucleotide alignment program. *Bioinformatics.* 2008;24:713–714.
- Delcher AL, Bratke KA, Powers EC, Salzberg SL. Identifying bacterial genes and endosymbiont DNA with Glimmer. *Bioinformatics.* 2007;23:673–679.
- van Domselaar GH, Stothard P, Shrivastava S, et al. BASys: a web server for automated bacterial genome annotation. *Nucleic Acids Res.* 2005;33:W455–W459.
- Aziz RK, Bartels D, Best AA, et al. The RAST Server: rapid annotations using subsystems technology. *BMC Genomics.* 2008;9:75.
- Schattner P, Brooks AN, Lowe TM. The tRNAscan-SE, snoscan and snoGPSweb servers for the detection of tRNAs and snoRNAs. *Nucleic Acids Res.* 2005;33:W686–W689.
- Lagesen K, Hallin PF, Rødland E, Stærfeldt HH, Rognes T, Ussery DW. RNAmmer: consistent annotation of rRNA genes in genomic sequences. *Nucleic Acids Res.* 2007;35:3100–3108.
- Varani AM, Siguier P, Goubeyre E, Charneau V, Chandler M. ISsaga is an ensemble of web-based methods for high throughput identification and semi-automatic annotation of insertion sequences in prokaryotic genomes. *Genome Biol.* 2011;12:R30.