

CLINICAL SCIENCE

Effect of continuous and interval exercise training on the PETCO₂ response during a graded exercise test in patients with coronary artery disease

Enéas A. Rocco,¹ Danilo M. L. Prado,^{1,3} Alexandre G. Silva,³ Jaqueline M. A. Lazzari,¹ Pedro C. Bortz,¹ Débora F. M. Rocco,³ Carla G. Rosa,¹ Valter Furlan¹

¹TotalCor Amil Hospital São Paulo, São Paulo/SP, Brazil. ²Hospital das Clínicas da Faculdade de Medicina da Universidade de São Paulo, São Paulo/SP, Brazil. ³Faculdade de Educação Física e Esporte da Universidade Santa Cecília, Santos/SP, Brazil.

OBJECTIVE: The purpose of this study was to evaluate the following: 1) the effects of continuous exercise training and interval exercise training on the end-tidal carbon dioxide pressure (PETCO₂) response during a graded exercise test in patients with coronary artery disease; and 2) the effects of exercise training modalities on the association between PETCO₂ at the ventilatory anaerobic threshold (VAT) and indicators of ventilatory efficiency and cardiorespiratory fitness in patients with coronary artery disease.

METHODS: Thirty-seven patients (59.7 ± 1.7 years) with coronary artery disease were randomly divided into two groups: continuous exercise training ($n=20$) and interval exercise training ($n=17$). All patients performed a graded exercise test with respiratory gas analysis before and after three months of the exercise training program to determine the VAT, respiratory compensation point (RCP) and peak oxygen consumption.

RESULTS: After the interventions, both groups exhibited increased cardiorespiratory fitness. Indeed, the continuous exercise and interval exercise training groups demonstrated increases in both ventilatory efficiency and PETCO₂ values at VAT, RCP, and peak of exercise. Significant associations were observed in both groups: 1) continuous exercise training (PETCO₂VAT and cardiorespiratory fitness $r=0.49$; PETCO₂VAT and ventilatory efficiency $r=-0.80$) and 2) interval exercise training (PETCO₂VAT and cardiorespiratory fitness $r=0.39$; PETCO₂VAT and ventilatory efficiency $r=-0.45$).

CONCLUSIONS: Both exercise training modalities showed similar increases in PETCO₂ levels during a graded exercise test in patients with coronary artery disease, which may be associated with an improvement in ventilatory efficiency and cardiorespiratory fitness.

KEYWORDS: Coronary Artery Disease; Exercise; Cardiorespiratory Fitness; Ventilatory Efficiency.

Rocco EA, Prado DM, Silva AG, Lazzari JM, Bortz PC, Rocco DF, et al. Effect of continuous and interval exercise training on the PETCO₂ response during a graded exercise test in patients with coronary artery disease. Clinics. 2012;67(6):623-627.

Received for publication on January 28, 2012; First review completed on February 15, 2012; Accepted for publication on February 15, 2012

E-mail: danielomprado@usp.br

Tel.: 55 11 3576-1049

INTRODUCTION

End-tidal carbon dioxide pressure (PETCO₂) is a non-invasive index that is considered to be a good indicator for evaluating the ventilation/perfusion relationship in patients over a wide range of conditions (1,2). Variations in PETCO₂ have been shown to reflect changes in both cardiac output and pulmonary blood flow in animals and humans under constant ventilation (3). For example, previous studies (4,5,6) have demonstrated alterations in the distribution of

ventilation and perfusion in the lungs (ventilation-perfusion mismatch) of cardiac patients. This observation has led to an increased ratio of physiologic dead space to tidal volume. In this regard, patients with cardiac disease have been shown to have an abnormally low PETCO₂ during exercise, especially those with an impaired response of cardiac output during exercise (5). Moreover, recent investigations have shown that reduced PETCO₂ is considered to be a strong predictor of adverse events in cardiac disease (7,8).

Aerobic exercise training has been recommended as a non-pharmacological treatment for patients with coronary artery disease (CAD) (9,10). In this context, continuous exercise training (CET) promotes beneficial cardiorespiratory adaptations in CAD patients (11). Interestingly, previous investigations (12) have also demonstrated the effectiveness of interval exercise training (IET) for improving cardiorespiratory fitness in CAD patients. IET involves alternating brief (2–5 min) higher-intensity ($\geq 75\% \text{ VO}_{2\text{peak}}$)

Copyright © 2012 CLINICS – This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

No potential conflict of interest was reported.

Table 1 - Patient characteristics.

	All patients	CET	IET
N	37	20	17
Age (years)	59.7 ± 1.7	62.5 ± 2.0	56.5 ± 3.0
Weight (kg)	79.8 ± 2.3	78.2 ± 3.1	81.6 ± 3.7
BMI (kg/m ²)	28.3 ± 0.6	28.1 ± 1.0	28.5 ± 0.9
Men/women	28/9	15/5	14/3
Hypertension	35 (84%)	19 (54%)	16 (46%)
Diabetes	11 (30%)	4 (36%)	7 (64%)
Myocardial infarction	17 (46%)	11 (65%)	6 (35%)
Drugs			
Beta-blockers	27 (72%)	16 (59%)	11 (41%)
ACE inhibitors	24 (65%)	10 (42%)	14 (58%)

Values are means ± SE. CET = continuous exercise training; IET = interval exercise training; BMI = body mass index. $p > 0.01$ for all variables between groups.

and moderate-intensity workloads throughout an exercise session. However, little is known regarding the impact of the exercise training modality on the PETCO₂ response during a graded exercise test in CAD patients. Thus, the purpose of this study was to evaluate the following: 1) the effects of continuous exercise training and interval exercise training on the PETCO₂ response during a graded exercise test in CAD patients; and 2) the effects of exercise modalities on the association between PETCO₂ at the ventilatory anaerobic threshold (VAT) and indicators of ventilatory efficiency and cardiorespiratory fitness in CAD patients.

MATERIALS AND METHODS

Population

The patients were admitted to the coronary care unit from the TotalCor Hospital for the diagnosis of coronary artery disease. Forty-five CAD patients (60.0 ± 1.7) were initially enrolled into this study. The inclusion criterion was stable coronary artery disease diagnosed by coronary angiography. The exclusion criteria were unstable angina pectoris, complex ventricular arrhythmias, pulmonary congestion, and orthopedic or neurological limitations to exercise. Thirty-seven patients meeting the inclusion/exclusion criteria were considered for this study. The CAD patients were randomly divided into two groups: CET (n = 20) and IET (n = 17). The patients remained on their standard medication throughout the study, and no changes were reported. The study participants read a detailed description of the protocol and completed a written informed consent (Table 1).

Graded Exercise Test

A maximal graded exercise test was performed on a programmable treadmill (DigiStress model pulsar, Gobernador Valadares, MG, Brazil). The gas exchange and ventilatory variables were measured continuously during the gas exchange tests, breath by breath, using an open-circuit spirometry procedure based on an exercise system (SensorMedics - model Vmax 229 Pulmonary Function/ Cardiopulmonary Exercise Testing Instrument, Yorba Linda, CA, USA). The following variables were obtained breath by breath and were expressed as 30-s averages: pulmonary oxygen uptake (VO₂ ml·kg⁻¹·min⁻¹ STPD), respiratory exchange ratio (RER), pulmonary ventilation (VE l·min⁻¹ BTPS), functional estimate of dead space (VD/VT), ventilatory equivalents for oxygen and carbon dioxide

(VE/VO₂ and VE/VCO₂) and end-tidal pressures for oxygen and carbon dioxide (PETO₂ and PETCO₂ mmHg). Before each test, the gas analyzers were calibrated using gases of known concentrations of carbon dioxide and oxygen balanced with nitrogen, and the flow meter was calibrated using a 3-L syringe. The heart rate was continuously recorded at rest, during the graded exercise testing and during the recovery period using a 12-lead ECG (HW Systems- HeartWare Ltd.). All tests in this study were performed in the same laboratory at the same room temperature (20–23 °C).

The subjects performed a ramp-like progressive exercise test until they were exhausted on the treadmill. The exercise workload (speed and/or slope) was increased every 1 minute; the incremental part of the exercise test was completed between approximately 8 and 12 minutes.

Ventilatory Anaerobic Threshold

The ventilatory anaerobic threshold (VAT) was determined to occur at the break point between the increase in the carbon dioxide output and VO₂ (V-Slope) or the point at which the VE/VO₂ reached the minimum value and began to rise without a concomitant rise in VE/VCO₂ (13).

Respiratory Compensation Point

The respiratory compensation point (RCP) was determined to be the point at which the VE/VCO₂ reached its minimum value and began to rise and the highest value of PETCO₂ before its progressive fall (14).

Peak Oxygen Consumption

The peak oxygen consumption (VO_{2peak}) was defined as the maximum attained VO₂ at the end of the exercise period, when the subject is exhausted (analog scale of the perceived exertion to the Borg scale).

Exercise Training Program

A supervised exercise training program was conducted at the cardiorespiratory rehabilitation center of the TotalCor Hospital. The exercise training program consisted of three 60-minute exercise sessions per week over a 3-month period. Each exercise session consisted of a 5 minute warm-up, 50 minutes of aerobic exercise and 5 minutes of cool-down exercises. The CET was performed on a treadmill with a 50-minute duration and intensity at VAT. The IET consisted of seven sets of 3 minutes at RCP and seven sets of 3 minutes of exercise at moderate intensity corresponding to the VAT totaling 42 minutes. Heart rate was monitored throughout the session to ensure that all patients exercised within the limits of intensity.

Statistical Analysis

The statistical procedures were performed using SPSS version 16.0 (SPSS Inc., Chicago, IL). The normality of the distribution was checked for all variables using the Kolmogorov-Smirnov test. A two-way analysis of variance (ANOVA) with repeated measures was performed to evaluate possible within-group and between-group differences for physical characteristics, cardiorespiratory fitness, ventilatory efficiency and PETCO₂ response during GET in the CAD patients subjected to CET and IET. When significance was identified, a Tukey's post hoc comparison was performed.

Table 2 - Physical and cardiorespiratory measurements in the CAD patients subjected to continuous exercise training and interval exercise training.

	CET		IET	
	Pre	Post	Pre	Post
<i>Physical measurements</i>				
Age (years)	62.5 ± 2.1	62.6 ± 2.1	56.5 ± 2.7	56.8 ± 2.6
Weight (kg)	78.2 ± 3.3	78.1 ± 3.3	81.6 ± 3.3	81.9 ± 3.0
BMI (kg/m ²)	28.1 ± 1.0	28.1 ± 1.0	28.5 ± 0.7	28.5 ± 0.7
<i>Cardiorespiratory measurements</i>				
VO _{2peak} mL/kg/min	18.0 ± 1.2	22.2 ± 1.3*	17.9 ± 1.0	22.3 ± 1.1*
VO ₂ (VAT) mL/kg/min	12.5 ± 0.7	15.0 ± 0.7*	12.3 ± 0.6	14.0 ± 0.7*
VEVCO ₂ (VAT)	33.0 ± 0.8	29.7 ± 0.6*	31.8 ± 0.4	29.1 ± 0.6*
VD/VT (VAT)	0.20 ± 0.0	0.16 ± 0.0*†	0.20 ± 0.0	0.18 ± 0.0*
PETCO ₂ (VAT) mmHg	37.5 ± 0.6	41.0 ± 0.4*†	38.0 ± 0.6	40.2 ± 0.3*

Values are means ± SE. Pre, pre-intervention; Post, post-intervention; CET, continuous exercise training; IET, interval exercise training; BMI, body mass index; VAT, ventilatory anaerobic threshold. *p < 0.05 vs. pre-intervention; †p < 0.01 vs. pre-intervention.

For both groups, the relative difference (Δ%) was calculated for PETCO₂ in the intervals from rest to VAT, rest to RCP, and rest to peak of exercise.

The Pearson-product moment correlation was performed to examine the association between PETCO₂ at VAT and the cardiorespiratory fitness and ventilatory efficiency after the interventions. The data are presented as means ± SE. A p-value of <0.05 was considered statistically significant.

RESULTS

Effects of interventions

Baseline measurements. Before the interventions, no differences in the physical characteristics between the groups studied were noted (Table 2). In addition, both groups exhibited similar levels of cardiorespiratory fitness and ventilatory efficiency (Table 2).

Physical characteristics. After the interventions, neither the CET nor IET showed a significant reduction in body weight (Table 2).

Cardiorespiratory response. After the interventions, both the CET and IET showed an increase in the VO_{2peak}. A comparison between the groups revealed a similar increase in the cardiorespiratory fitness (Table 2).

The ventilatory efficiency was analyzed at the VAT. After the exercise training program, the CAD patients subjected to either CET or IET showed significantly lower values for VEVCO₂ and VD/VT. In addition, both exercise training modalities demonstrated an increase in the PETCO₂ at VAT (Table 2). A comparison between the groups revealed a similar increase in the ventilatory efficiency (Table 2).

After the interventions, both the CET and IET showed values significantly greater for the PETCO₂ during the GET (Figure 1).

The changes in the PETCO₂ response from rest to VAT, rest to RCP, and rest to peak of exercise are shown in Figure 2. The statistical analysis determined that the relative difference of the PETCO₂ in all intervals analyzed were significantly higher after the interventions.

The results of the correlation analysis between PETCO₂ at VAT and the cardiorespiratory parameters after either CET or IET are shown in Table 3.

Positive associations were observed in both groups for cardiorespiratory fitness: 1) CET (PetCO₂VAT and VO_{2peak} r = 0.49; PetCO₂VAT and VO₂VAT r = 0.49; p < 0.05) and 2) IET (PetCO₂VAT and VO_{2peak} r = 0.39; PetCO₂VAT and VO₂VAT r = 0.33; p < 0.05). However, negative associations were observed in both groups for ventilatory efficiency: 1) CET (PetCO₂VAT and VEVCO₂VAT r = -0.80; PetCO₂VAT




Figure 1 - PETCO₂ response during a graded exercise test in patients with coronary artery disease subjected to either continuous exercise training or interval exercise training. VAT = ventilatory anaerobic threshold; RCP = respiratory compensation point. † p < 0.05 vs. rest (pre- and post-continuous); ‡ p < 0.05 vs. rest (pre- and post-interval); * p < 0.05 vs. post-continuous; § p < 0.05 vs. post-interval.




Figure 2 - Percentage difference from rest to ventilatory anaerobic threshold (rest/VAT), rest to respiratory compensation point (rest/RCP) and rest to peak of exercise (rest/Peak) in patients with coronary artery disease subjected to continuous exercise training (panel A) and interval exercise training (panel B). VAT=ventilatory anaerobic threshold; RCP=respiratory compensation point. * p<0.05 vs. post-intervention.

and VD/VT VAT r = -0.57; p<0.05) and 2) IET (PetCO₂VAT and VEVCO₂VAT r = -0.45; PetCO₂VAT and VD/VT VAT r = -0.42; p<0.05).

DISCUSSION

The main findings of this investigation were the following: 1) continuous exercise training and interval exercise training improved the PETCO₂ response during a graded exercise test in CAD patients; 2) both exercise modalities resulted in similar increases in the values of PETCO₂; and 3) after exercise training, the improvements in ventilatory efficiency and cardiorespiratory fitness in patients with CAD were associated with increased values of PETCO₂.

Table 3 - Correlations between PETCO₂ at ventilatory anaerobic threshold and the cardiorespiratory measurements in CAD patients after continuous exercise training and interval exercise training.

	CET		IET	
	r	p-value	r	p-value
VO _{2peak} mL/kg/min	0.49	0.01	0.39	0.01
VO ₂ (VAT) mL/kg/min	0.49	0.01	0.33	0.05
VEVCO ₂ (VAT)	-0.80	0.01	-0.45	0.01
VD/VT (VAT)	-0.57	0.01	-0.42	0.01

CET, continuous exercise training; IET, interval exercise training.

r = Pearson's product moment correlation coefficient.

The elucidation of the mechanisms involved in the improvement observed in the PETCO₂ response after exercise training in the CAD patients are beyond the scope of this study. However, we suggest that factors related to the improvement of the ventilation-perfusion (V/Q) mismatch may be responsible for the increase in the PETCO₂ during a graded exercise test (11).

In this regard, previous investigations have shown that the failure of pulmonary blood flow (cardiac output) to increase appropriately during exercise and increased physiological dead space are the main factors leading to a decrease in PETCO₂ in cardiac patients (1,15). For example, Matsumoto et al. (5) reported a positive association between cardiac output and PETCO₂ during a graded exercise test in cardiac patients. In addition, ETO et al. (11) observed a close relationship between increased PETCO₂ and cardiac output after aerobic exercise training in cardiac patients. In contrast, previous studies (1,5,15) reported an association between low levels of PETCO₂ during exercise with ventilatory inefficiency. Accordingly, our findings suggest that factors related to the increase in both ventilatory efficiency and cardiorespiratory fitness were mainly responsible for the increase in PETCO₂ during the graded exercise test. In fact, after 12 weeks of either CET or interval exercise training, the CAD patients had lower values for VEVCO₂ and VD/VT at VAT. Moreover, the CAD patients subjected to both exercise training modalities demonstrated increases in VO_{2peak}. After the interventions, we observed a positive association between PETCO₂ at VAT and VO_{2peak} and a negative association between PETCO₂ at VAT and VEVCO₂ at VAT. These findings suggest that the improvement in the V/Q mismatch after exercise training may be associated with an increase in the PETCO₂ at VAT. Another possible explanation for the increased PETCO₂ after exercise training would be raising the set point of PaCO₂ modulation by peripheral chemoreceptors and/or hyperventilation (11,16).

With regard to cardiorespiratory fitness, the CAD patients subjected to either CET or IET exhibited similar increases in VO_{2peak}. In this context, our findings are in agreement with previous studies that demonstrated the same effectiveness of both exercise training modalities to increase the cardiorespiratory fitness in CAD patients (12,17). However, other investigations (18,19,20) observed greater responsiveness to interval exercise training compared with continuous exercise training.

For example, Rogmo et al. (18) observed a superior increase in VO_{2peak} in CAD patients subjected to IET, in which 4 minutes at a higher intensity (80-90% VO_{2peak}) and 3 minutes at a moderate intensity (50-60% VO_{2peak}) were alternated throughout the exercise session. In the same study (18), the CAD patients subjected to CET performed 41 minutes at a moderate intensity (50-60% VO_{2peak}) over 10 weeks. In the same context, Ciolac et al. (20) observed that interval exercise training (80% of reserve heart rate) in hypertensive patients increased cardiorespiratory fitness to a higher degree than continuous exercise training (60% of reserve heart rate).

In this investigation, the CAD patients subjected to IET performed 3 minutes at RCP (80-90% VO_{2peak}) and 3 minutes at moderate intensity corresponding to the VAT. In contrast, the patients subjected to CET performed 50 minutes at VAT (70-80% VO_{2peak}) over 12 weeks.

These results conflict with previous studies (18,20) and may be associated with a greater relative intensity and higher training volume in the CET group of our study.

In fact, evidence has suggested that stroke volume is the main factor that limits cardiorespiratory fitness (20). Moreover, higher intensities of aerobic exercise promote an increase in central O₂ delivery (cardiac output), thereby contributing to the training-induced improvements in VO_{2peak} (20).

These findings demonstrate the need for a greater number of randomized trials to investigate the mode of aerobic exercise training that is more responsive in terms of increased cardiorespiratory fitness for patients with coronary artery disease.

Interestingly, after an exercise training program, the CAD patients did not display any reduction in body weight. Recent studies (21,22) have indicated that exercise-related caloric expenditure for patients enrolled in cardiac rehabilitation programs was only modest.

In both exercise training modalities, we observed energy expenditures of approximately 230 kcal/session. For example, Savage et al. (2000) reported that a caloric expenditure of 270 kcal/session during three months of a cardiac rehabilitation program had little impact on measures of obesity.

This study has limitations, such as the lack of evaluations of cardiac output and arterial blood gases during the graded exercise test.

In summary, these findings suggest that both continuous exercise training and interval exercise training promote an improved PETCO₂ response during a graded exercise test in patients with coronary artery disease. Taken together, these results demonstrate that the application of different modalities of aerobic exercise training may be important in improving cardiorespiratory efficiency in patients with coronary artery disease.

AUTHOR CONTRIBUTIONS

Rocco EA wrote the manuscript. Prado ML wrote the manuscript and was responsible for the statistical analyses. Silva AG wrote the manuscript. Lazzari JM and Bortz PC were responsible for the application of physical training in the patients. Rocco DF wrote the manuscript and reviewed the English version. Rosa CG was responsible for conducting the cardiopulmonary evaluation in the patients. Furlan V was responsible for the statistical analyses.

REFERENCES

1. Tanabe Y, Hosaka Y, Ito M, Ito E, Suzuki K. Significance of end-tidal P(CO₂) response to exercise and its relation to functional capacity in patients with chronic heart failure. *Chest*. 2001;119(3):811-7, <http://dx.doi.org/10.1378/chest.119.3.811>.
2. Hansen JE, Ulubay G, Chow BF, Sun XG, Wasserman K. Mixed-expired and end-tidal CO₂ distinguish between ventilation and perfusion defects during exercise testing in patients with lung and heart diseases. *Chest*. 2007;132(3):977-83, <http://dx.doi.org/10.1378/chest.07-0619>.
3. Idris AH, Staples ED, O'Brien DJ, Melker RJ, Rush WJ, Del Duca KD, et al. End-tidal carbon dioxide during extremely low cardiac output. *Annals of emergency medicine*. 1994;23(3):568-72, [http://dx.doi.org/10.1016/S0196-0644\(94\)70080-X](http://dx.doi.org/10.1016/S0196-0644(94)70080-X).
4. Wasserman K, Zhang YY, Gitt A, Belardinelli R, Koike A, Lubarsky L, et al. Lung function and exercise gas exchange in chronic heart failure. *Circulation*. 1997;96(7):2221-7.
5. Matsumoto A, Itoh H, Eto Y, Kobayashi T, Kato M, Omata M, et al. End-tidal CO₂ pressure decreases during exercise in cardiac patients: association with severity of heart failure and cardiac output reserve. *Journal of the American College of Cardiology*. 2000;36(1):242-9, [http://dx.doi.org/10.1016/S0735-1097\(00\)00702-6](http://dx.doi.org/10.1016/S0735-1097(00)00702-6).
6. Myers JN, Gujja P, Neelagaru S, Hsu L, Burkhoff D. Noninvasive measurement of cardiac performance in recovery from exercise in heart failure patients. *Clinics (Sao Paulo)*. 2011;66(4):649-56, <http://dx.doi.org/10.1590/S1807-59322011000400021>.
7. Arena R, Peberdy MA, Myers J, Guazzi M, Tevald M. Prognostic value of resting end-tidal carbon dioxide in patients with heart failure. *Int J Cardiol*. 2006;109(3):351-8, <http://dx.doi.org/10.1016/j.ijcard.2005.06.032>.
8. Arena R, Guazzi M, Myers J. Prognostic value of end-tidal carbon dioxide during exercise testing in heart failure. *Int J Cardiol*. 2007;117(1):103-8, <http://dx.doi.org/10.1016/j.ijcard.2006.04.058>.
9. Hagberg JM. Physiologic adaptations to prolonged high-intensity exercise training in patients with coronary artery disease. *Medicine and science in sports and exercise*. 1991;23(6):661-7.
10. Kendziorra K, Walther C, Foerster M, Mobius-Winkler S, Conradi K, Schuler G, et al. Changes in myocardial perfusion due to physical exercise in patients with stable coronary artery disease. *Eur J Nucl Med Mol*. 2005;32(7):813-9, <http://dx.doi.org/10.1007/s00259-005-1768-1>.
11. Eto Y, Koike A, Matsumoto A, Momomura S, Tajima A, Aizawa T, et al. Early aerobic training increases end-tidal CO₂ pressure during exercise in patients after acute myocardial infarction. *Circ J*. 2004;68(8):778-83, <http://dx.doi.org/10.1253/circj.68.778>.
12. Moholdt TT, Amundsen BH, Rustad LA, Wahba A, Lovo KT, Gullikstad LR, et al. Aerobic interval training versus continuous moderate exercise after coronary artery bypass surgery: A randomized study of cardiovascular effects and quality of life. *Am Heart J*. 2009;158(6):1031-7, <http://dx.doi.org/10.1016/j.ahj.2009.10.003>.
13. Wasserman K, Whipp BJ, Koyal SN, Beaver WL. Anaerobic Threshold and Respiratory Gas-Exchange during Exercise. *J Appl Physiol*. 1973;35(2): 236-43.
14. Braga A, Nunes N. Avaliação Cardiopulmonar. In: Negrão CE, Barreto ACP, editors. *Cardiologia do Exercício: do Atleta ao Cardiopata*. Manole; 2005. p. 128-47.
15. Myers J, Gujja P, Neelagaru S, Hsu L, Vittorio T, Jackson-Nelson T, et al. End-tidal CO₂ pressure and cardiac performance during exercise in heart failure. *Medicine and science in sports and exercise*. 2009;41(1):19-25.
16. Tomita T, Takaki H, Hara Y, Sakamaki F, Satoh T, Takagi S, et al. Attenuation of hypercapnic carbon dioxide chemosensitivity after postinfarction exercise training: possible contribution to the improvement in exercise hyperventilation. *Heart*. 2003;89(4):404-10, <http://dx.doi.org/10.1136/heart.89.4.404>.
17. Mirkova L, Siegelova J, Vymazalova L, Svacanova H, Vank P, Panovsky R, et al. [Interval and continuous training in cardiovascular rehabilitation]. *Vnitri lekarstvi*. 2006;52(1):44-50.
18. Rognmo Ø, Hetland E, Helgerud J, Hoff J, Slordahl SA. High intensity aerobic interval exercise is superior to moderate intensity exercise for increasing aerobic capacity in patients with coronary artery disease. *Eur J Cardiovasc Prev R*. 2004;11:216-222, <http://dx.doi.org/10.1097/01.hjr.0000131677.96762.0c>.
19. Wisloff U, Stoylen A, Loennechen JP, Bruvold M, Rognmo O, Haram PM, et al. Superior cardiovascular effect of aerobic interval training versus moderate continuous training in heart failure patients - A randomized study. *Circulation*. 2007;115(24):3086-94, <http://dx.doi.org/10.1161/CIRCULATIONAHA.106.675041>.
20. Ciolac EG, Guimaraes GV, D'Avila VM, Bortolotto LA, Doria EL, Bocchi EA. Acute effects of continuous and interval aerobic exercise on 24-h ambulatory blood pressure in long-term treated hypertensive patients. *Int J Cardiol*. 2009;133(3):381-7, <http://dx.doi.org/10.1016/j.ijcard.2008.02.005>.
21. Schairer JR, Kostelnik T, Proffitt SM, Faitel KI, Windeler S, Rickman LB, et al. Caloric expenditure during cardiac rehabilitation. *J Cardiopulm Rehabil*. 1998;18(4):290-4, <http://dx.doi.org/10.1097/00008483-199807000-00006>.
22. Savage PD, Brochu M, Scott P, Ades PA. Low caloric expenditure in cardiac rehabilitation. *Am Heart J*. 2000;140(3):527-33, <http://dx.doi.org/10.1067/mhj.2000.109219>.