



Karyotypic conservatism in samples of *Characidium cf. zebra* (Teleostei, Characiformes, Crenuchidae): Physical mapping of ribosomal genes and natural triploidy

José Carlos Pansonato-Alves, Claudio Oliveira and Fausto Foresti

Departamento de Morfologia, Instituto de Biociências, Universidade Estadual Paulista “Júlio de Mesquita Filho”, Botucatu, SP, Brazil.

Abstract

Basic and molecular cytogenetic analyses were performed in specimens of *Characidium cf. zebra* from five collection sites located throughout the Tietê, Paranapanema and Paraguay river basins. The diploid number in specimens from all samples was $2n = 50$ with a karyotype composed of 32 metacentric and 18 submetacentric chromosomes in both males and females. Constitutive heterochromatin was present at the centromeric regions of all chromosomes and pair 23, had additional interstitial heterochromatic blocks on its long arms. The nucleolar organizer regions (NORs) were located on the long arms of pair 23, while the 5S rDNA sites were detected in different chromosomes among the studied samples. One specimen from the Alambari river was a natural triploid and had two extra chromosomes, resulting in $2n = 77$. The remarkable karyotypic similarity among the specimens of *C. cf. zebra* suggests a close evolutionary relationship. On the other hand, the distinct patterns of 5S rDNA distribution may be the result of gene flow constraints during their evolutionary history.

Key words: *Characidium*, heterochromatin, 18S rDNA, 5S rDNA, natural triploidy.

Received: July 8, 2010; Accepted: October 29, 2010.

Introduction

Characidium is the richest genus in number of species within the subfamily Characidiinae (Crenuchidae). It comprises about 50 valid species widespread throughout the Neotropical region, from Panama to Argentina (Buckup, 2003). The phylogenetic status of this assemblage has been controversial since the first description of Characidiinae, which has already been regarded as a member of Hemiodontidae, a subfamily of Characidae or an independent family (Buckup, 1998). Studies based on morphological data using cladistic methods indicated that Characidiinae along with Crenuchinae formed the family Crenuchidae (Buckup, 1998).

Characidium zebra represents a dispersed and polymorphic species, often misidentified as *C. fasciatum* (Buckup and Reis, 1997). This species is characterized by the presence of a completely scaled isthmus. The presence of several plesiomorphic morphological traits led to the conclusion that this species was primitive in relation to its congeners, occupying a basal position in the phylogeny of *Characidium* (Buckup, 1993). The type locality of *C. zebra*

is in the Amazon drainage, but there are no studies defining its actual range (Buckup and Reis, 1997). Due to its phylogenetic position, the cytogenetic analysis of species related to *C. zebra* may represent an interesting starting point for understanding events related to the karyotypic evolution within the genus.

The objective of this work was to study the karyotypes of five samples of *C. cf. zebra* collected in three major Brazilian river basins (Tietê, Paranapanema and Paraguay) in order to contribute to the understanding of the processes involved in the karyotypic differentiation of *Characidium* species.

Material and Methods

Specimens of *Characidium cf. zebra* from five collection sites along distinct hydrographic basins in Brazil were analyzed. The specimens were fixed in 10% formaldehyde, stored in 70% ethanol for identification, and deposited in the fish collection of the Laboratório de Biologia e Genética de Peixes (LBP) at UNESP, Botucatu, São Paulo, Brazil (Table 1).

Chromosome preparations were obtained from kidney and gill tissue following the technique described by Foresti *et al.* (1981). Nucleolar organizer regions (NORs) were evidenced by silver nitrate staining according to

Send correspondence to José Carlos Pansonato Alves. Departamento de Morfologia, Instituto de Biociências, Universidade Estadual Paulista, “Júlio de Mesquita Filho”, Distrito de Rubião Junior s/n, 18618-970 Botucatu, SP, Brazil. E-mail: pansonato@ibb.unesp.br.

Table 1 - Specimens of *Characidium cf. zebra* analyzed.

LBP	Sample Localities	F	M	Coordinates
8704	Paraitinga river – Salesópolis – SP	12	7	S 23°30'40 W 45°51'32
8714	Alambari river – Botucatu – SP	25	10	S 22°56'06 W 48°19'18
6727	Novo river – Avaré – SP	10	4	S 23°01'26 W 48°49'32
8715	Araquá river – Botucatu – SP	8	5	S 22°47'13 W 48°28'89
8533	Juba river – Nova Fernandópolis – MT	6	5	S 14°58'23 W 57°44'40

LBP: deposit number at the fish collection of the Laboratório de Biologia e Genética de Peixes, Instituto de Biociências, UNESP. F: females and M: males; SP – São Paulo; MT – Mato Grosso.

Howell and Black (1980) and C-banding was performed based on Sumner (1972).

Molecular cytogenetic analyses comprised the identification of GC- and AT-rich regions by fluorochrome staining with chromomycin A₃ (Schweizer, 1976) and DAPI (Schweizer *et al.*, 1978), respectively. Mapping of both 18S and 5S rDNA sites on metaphase chromosomes was performed by fluorescent *in situ* hybridization (FISH) according to Pinkel *et al.* (1986), with slight modifications. The probes used were a 18S rDNA sequence obtained after PCR from the nuclear DNA of *Prochilodus argenteus* (Hatanaka and Galetti Jr, 2004) using the primers NS1 5'-GTAGTCATATGCTTGTCTC-3' and NS8 5'-TCCGCAGGTTACCTACGGA-3' (White *et al.*, 1990) and a 5S rDNA sequence amplified using the primers A (5'-TACGCCGATCTCGTCCGATC-3') and B (5'-GCTGGTATGGCCGTAGC-3') of *Leporinus elongatus* (Martins and Galetti Jr, 1999). The 18S rDNA probe was labeled with 14-dATP biotin by nick translation, according to the manufacturer's instructions (Bionick Labeling System – Invitrogen). Signal detection and amplification were performed with fluorescein-conjugated avidin (FITC) and biotinylated anti-avidin antibodies. The 5S rDNA probe was labeled with digoxigenin-11-dUTP (Roche Applied Science) by PCR and the hybridization signal was detected with anti-digoxigenin conjugated with rhodamine.

Chromosomes were counterstained with DAPI or propidium iodide (in the case of the triploid specimen) and analyzed under an epifluorescence photomicroscope (Olympus BX61). Images were captured using Image Pro Plus, 6.0 software (MediaCybernetics). The chromosomes were morphologically classified according to their arms ratio, as established by Levan *et al.* (1964), and were arranged in decreasing size.

Results

The five samples of *Characidium cf. zebra* presented a constant diploid number of $2n = 50$ and a karyotype composed of 32 metacentric and 18 submetacentric chromosomes. Constitutive heterochromatin was identified only at the centromeric regions of most chromosomes in individuals of both sexes. Pair 23, presented additional interstitial

heterochromatic blocks on its long arms in some metaphases, which colocalized with the NORs (Figure 1). No sex chromosome heteromorphism was identified. The Ag-NORs, evidenced after silver nitrate staining (Figure 1, inset), CMA₃ staining (Figure 2a) and FISH with a 18S rDNA probe (Figure 3d), were located at an interstitial position in the long arm of pair 23. Chromomycin A₃ staining allowed the demonstration that the NORs were located in GC-rich segments in all the populations studied.

FISH with the 5S rDNA probe showed that the specimens from the Paraitinguinha river presented 5S rRNA genes at interstitial positions of five chromosomes (pairs 1 and 6 and one homologue of pair 17) (Figure 3b). In the individuals from the Novo, Alambari and Araquá rivers (Figure 3a), this ribosomal sequences were located at an




Figure 1 - Karyotype present in the five populations of *Characidium cf. zebra* analyzed after Giemsa staining (a) and C-banding (b). In the inset, the NOR-bearing chromosomes. Bar = 10 μ m.




Figure 2 - Metaphases of *Characidium cf. zebra* after chromomycin A₃ (a) and DAPI (b) staining. The arrows indicate GC-rich (a) and AT-poor (b) sites. Bar = 10 μ m.




Figure 3 - Karyotype of *Characidium cf. zebra* showing the 5S rDNA on pair 17, typical of the populations from the Alambari, Novo and Araquá rivers (a); in pairs 1, 6 and one homologue of pair 17, from the Paratinguinha river (b) and in pairs 1, 6, 9, 17 and 18 from the Juba river (c). In (d), the karyotype present in the five populations of *Characidium cf. zebra* with the 18S rDNA on pair 23. Bar = 10 µm.

interstitial position on pair 17. The 5S rDNA sites were localized at interstitial regions of five chromosome pairs (1, 6, 9, 17 and 18) in the samples from the Juba river (Figure 3c).

All the cells of one specimen collected in the Alambari river presented 77 chromosomes, characterizing a case of natural triploidy (Figure 4). This individual also carried two extra chromosomes (Figure 4a, inset). C-banding revealed heterochromatic blocks only in the centromeric regions of all chromosomes of the normal set (Figure 4b) in both diploid and triploid specimens. The extra chromo-




Figure 4 - Karyotype of a triploid specimen of *Characidium cf. zebra* from the Alambari river after Giemsa staining (a) and C-banding (b). In the inset, the supernumerary chromosomes. Bar = 10 µm.




Figure 5 - Metaphases of a triploid specimen of *Characidium cf. zebra* showing NORs after: silver nitrate staining (a), FISH with a 18S rDNA probe (b) and CMA₃ staining (c). Note that there are only two NOR-bearing chromosomes. Bar = 10 µm.

somes were almost completely euchromatic (Figure 4b, inset), except for their centromeric regions. Although the specimen was a triploid, its NORs, identified by silver nitrate staining, CMA₃, and FISH with the 18S rDNA probe, were restricted to two chromosomes (Figure 5).

Discussion

Cytogenetic studies in five samples of *Characidium cf. zebra* showed a constant $2n = 50$ with similar karyotypes comprising metacentric and submetacentric chromosomes. The largest chromosome pair (pair 1) in this karyotype is remarkably larger than the second pair (pair 2) in all the individuals, thus representing an important marker for the species. The largest chromosomes in species such as *Characidium cf. gomesi* (Centofante *et al.*, 2001; Maistro *et al.*, 1998, Pansonato-Alves *et al.*, 2010b), *C. lanei* (Noletto *et al.*, 2009, Pansonato-Alves *et al.*, 2010a) and *C. schubarti* (Pansonato-Alves *et al.*, 2010a) are those from the 17th submetacentric pair. The presence of the marker chromosome pair 1, the similar distribution pattern of heterochromatin and of the NORs suggest that the studied samples of *C. cf. zebra* are closely related.

Based on the phylogenetic position of *C. zebra* proposed by Buckup (1993), the absence of differentiated sex chromosomes in this species would represent a plesiomorphic trait within the genus, whereas the presence of heteromorphic sex chromosomes in some representatives of the genus should be regarded as an apomorphy (Centofante *et al.*, 2003). According to Centofante *et al.* (2003), the occurrence of a ZW sex chromosome system with a heterochromatic W chromosome is restricted to a few species closely related to *C. gomesi*. However, among the ten *Characidium* species cytogenetically studied (Miyazawa and Galetti Jr, 1994; Maistro *et al.*, 1998; Centofante *et al.*, 2001, 2003 and Pansonato-Alves *et al.*, 2010a,b), only *C. cf. zebra* and *C. lagsantensis* lack heteromorphic sex chromosomes. This suggests that the presence of differentiated sex chromosomes is the most common condition of the character among the species of the genus.

The NOR-bearing chromosome pair 23 was identical in the five studied samples of *Characidium cf. zebra*. The region corresponding to the NOR was faintly stained by DAPI, reinforcing its AT-poor nature (Figure 2b). NORs

have been shown to be usually stained by GC-specific fluorochromes, such as chromomycin A₃ and mithramycin in fish (Pendás *et al.*, 1993; Vicari *et al.*, 2005) and amphibians (Schmid, 1982). However, some heterochromatic segments unrelated to NORs may also brightly fluoresce after GC-specific fluorochrome staining (Artoni *et al.*, 1999).

FISH with 5S rDNA probes in metaphase chromosomes of *Characidium* revealed a peculiar distribution of these sequences. Vicari *et al.* (2008) mapped the 5S rDNA genes to an interstitial position in a single metacentric pair of a *C. cf. gomesi* population from the Tibagi river. Pansonato-Alves *et al.* (2010b) located the same genes at an interstitial position in the 25th submetacentric pair in specimens from two populations of *C. cf. gomesi* from the Parapanema river basin and at an interstitial position in pairs 20 and 25 in individuals from the Tietê river basin. In the present study, the populations of *C. cf. zebra* from the Alambari, Araquá and Novo rivers had 5S rDNA sites at an interstitial position on pair 17. The 5S rDNA clusters were located in several chromosomes in populations from the Paraitinguinha (in pairs 1, 6 and one homologue of pair 17) and Juba rivers (in pairs 1, 23, 6, 9, 17 and 18). These findings indicate that, in spite of a close relationship among the studied samples, specific modifications have occurred on the basic genome and restrictions to gene flow during their evolutionary history may have determined the currently observed patterns. Furthermore, the 5S rDNA probes produced signals at centromeric heterochromatic regions in the populations from the Juba and Paraitinguinha rivers. This suggests that, some of them may correspond to pseudogenes representing inactive 5S rDNA-like sequences, as proposed for *Centropyge aurantonotus* (Affonso and Galetti Jr, 2005). The 5S rDNA and 18S rDNA sequences were located in distinct chromosome positions. This is in agreement with the most frequent pattern found in vertebrates, in which the two ribosomal gene families follow independent evolutionary pathways (Lucchini *et al.*, 1993; Suzuki *et al.*, 1996).

The presence of multiple 5S rDNA sites at interstitial regions of several chromosomes, a derived situation as compared to the plesiomorphic condition of a single interstitial site, may be due to structural rearrangements in the interphase nucleus (Schweizer and Loidl, 1987). This hypothesis contradicts the suggestion that an interstitial position would protect 5S rDNA sites (Martins and Galetti Jr, 2001), as chromosomal domains and heterochromatin dispersion could play a major role in the distribution of these genes.

The occurrence of natural triploidy in a single specimen of *C. cf. zebra* from the Alambari river may be related to the geography and climate of the Botucatu region, where other natural triploids have already been reported (Maistro *et al.*, 1994). In fish, external fecundation and completion of the second mitotic division only take place after spawning. Environmental temperature changes may facilitate the

retention of the second polar body and lead to the formation of natural triploids (Cuellar and Uyeno, 1972). Triploid fish are artificially produced by several techniques, such as thermal or pressure shocks and chemical treatment after fertilization, resulting in a large number of phenotypically normal polyploids (Purdom, 1983). In these procedures, the second meiotic division is disrupted, inhibiting anaphase II (Thorgaard *et al.*, 1981). The outcome is a triploid fish bearing three chromosome sets: two maternal and one paternal, as an unreduced diploid oocyte is fertilized by a normal haploid spermatozoon. The same features were observed in natural triploids of the genus *Cyprinus* (Vasilev *et al.*, 1975) and *Salmo* (Thorgaard and Allen Jr, 1987).

The occurrence of natural triploidy in *Characidium cf. zebra* seems to result from a process similar to that reported for *C. gomesi* (Centofante *et al.*, 2001), although the former species lacks heteromorphic sex chromosomes. However, in addition to the 75 chromosomes of the triploid specimen, two supranumerary chromosomes were observed, resulting in a $2n = 77$ (Figure 4). Cases of triploidy associated with B chromosomes have already been reported in *Astyanax scabripinnis* (Fauaz *et al.*, 1994; Maistro *et al.*, 1994) and *Curimata modesta* (Venere and Galetti Jr, 1985). In these studies, the extra chromosomes were easily identified by C-banding due to their heterochromatic nature. All the chromosomes of the triploid *C. cf. zebra* specimen presented centromeric heterochromatin, hindering the precise identification of the extra elements by C-banding (Figure 4). Nevertheless, two acro/subtelocentric chromosomes could be detected, contrasting to the meta- and submetacentric chromosomes commonly reported in *Characidium* karyotypes.

An extra element was also found in a single diploid *C. cf. zebra* from the Passa Cinco river, Ipeúna, SP by Miyazawa and Galetti Jr (1994). That chromosome was identified as an euchromatic acrocentric B chromosome, similar to those observed in our triploid specimen. There were no B chromosomes in the other 35 diploid specimens of *C. cf. zebra* from the Alambari river. This may indicate that the occurrence of extra chromosomes constitute a unique condition present in the triploid specimen and that they are a consequence of the same event that led to the formation of the triploid fish.

Conspicuous nucleolar organizer regions were located in only two chromosomes of the triploid specimen after silver nitrate and CMA₃ staining and FISH with a 18S rDNA probe. This may indicate a complete functional inactivation of the NOR in the third NOR-bearing homologue, as already reported in a Cyprinidae (Mayr *et al.*, 1986). However, the lack of CMA₃ and 18S rDNA signals in the third chromosome of our specimen suggests that other events, such as unequal crossovers, may be responsible for the loss or significant deletion of rDNA sequences in this chromosome.

The remarkable karyotypic similarities in the five populations of *C. cf. zebra* studied reinforces the possibility of a close relationship among them. Nevertheless, the distinct pattern of 5S rDNA distribution in the samples from the Paraitinguinha and Juba rivers most likely derive from constraints in gene flow during their evolutionary history. This, in turn, may indicate a tendency of fixation of allopatric differentiation mechanisms among populations isolated in headwaters of different hydrographic systems, which is related to the lifestyle of the species.

References

- Affonso PRAM and Galetti Jr PM (2005) Chromosomal diversification of reef fishes from genus *Centropyge* (Perciformes, Pomacanthidae). *Genetica* 123:227-233.
- Artoni RF, Molina WF, Bertollo LAC and Galetti Jr PM (1999) Heterochromatin analysis in the fish species *Liposarcus anisitsi* (Siluriformes) and *Leporinus elongatus* (Characiformes). *Genet Mol Biol* 22:39-44.
- Buckup PA (1993) The monophyly of the Characidiinae, a Neotropical group of characiform fishes (Teleostei, Ostariophysii). *Zool J Linn Soc* 108:224-245.
- Buckup PA (1998) Relationships of the Characidiinae and phylogeny of characiform fishes (Teleostei, Ostariophysii). In: Malabarba LR, Reis RE, Vari RP, Lucena CAS and Lucena ZMS (eds) *Phylogeny and Classification of Neotropical Fishes*. Edipucrs, Porto Alegre, pp 123-144.
- Buckup PA (2003) Family Crenuchidae (South American Darters). In: Reis RE, Kullander ESO and Ferraris Jr CJ (eds) *Check List of the Freshwater Fishes of South and Central America*. 2nd edition. Edipucrs, Porto Alegre, pp 87-95.
- Buckup PA and Reis RE (1997) Characidiin genus *Characidium* (Teleostei, Characiformes) in southern Brazil, with description of three new species. *Copeia* 3:531-548.
- Centofante L, Bertollo LAC and Moreira-Filho O (2001) Comparative cytogenetics among sympatric species of *Characidium* (Pisces, Characiformes). Diversity analysis with the description of a ZW sex chromosome system and natural triploidy. *Caryologia* 54:253-260.
- Centofante L, Bertollo LAC, Buckup PA and Moreira-Filho O (2003) Chromosomal divergence and maintenance of sympatric *Characidium* fish species (Crenuchidae, Characidiinae). *Hereditas* 138:213-218.
- Cuellar O and Uyeno T (1972) Triploidy in rainbow trout. *Cytogenetics* 11:508-515.
- Fauaz G, Vicente VE and Moreira-Filho O (1994) Natural triploidy and B chromosomes in the Neotropical fish genus *Astyanax* (Characidae). *Braz J Genet* 17:157-163.
- Foresti F, Almeida-Toledo LF and Toledo-Filho AS (1981) Polymorphic nature of nucleolus organizer regions in fishes. *Cytogenet Cell Genet* 31:137-144.
- Howell WM and Black DA (1980) Controlled silver staining of nucleolus organizer regions with a protective colloidal developer: A 1-step method. *Experientia* 36:1014-1015.
- Hatanaka T and Galetti Jr PM (2004) Mapping of the 18S and 5S ribosomal RNA genes in the fish *Prochilodus argenteus* Agassiz, 1829 (Characiformes, Prochilodontidae). *Genetica* 122:239-244.
- Levan A, Fredga K and Sandberg AA (1964) Nomenclature for centromeric position on chromosomes. *Hereditas* 52:201-220.
- Lucchini S, Nardi I, Barsacchi G, Batistoni R and Andronico F (1993) Molecular cytogenetics of the ribosomal (18S + 28S and 5S) DNA loci in primitive and advanced urodele amphibians. *Genome* 36:762-773.
- Martins C and Galetti Jr PM (1999) Chromosomal localization of 5s rDNA genes in *Leporinus* fish (Anostomidae, Characiformes). *Chromosome Res* 7:363-367.
- Martins C and Galetti Jr PM (2001) Two rDNA arrays in Neotropical fish species: Is it a general rule for fishes? *Genetica* 111:439-446.
- Maistro EL, Dias AL, Foresti F, Oliveira C and Moreira O (1994). Natural triploidy in *Astyanax scabripinnis* (Pisces, Characidae) and simultaneous occurrence of macro B-chromosomes. *Caryologia* 47:233-239.
- Maistro EL, Mata EP, Oliveira C and Foresti F (1998) Unusual occurrence of a ZZ/ZW sex-chromosome system and supernumerary chromosomes in *Characidium cf. fasciatum* (Pisces, Characiformes, Characidiinae). *Genetica* 104:1-7.
- Mayr B, Rab P and Kalat M (1986) NORs and counterstain-enhanced fluorescence studies in Cyprinidae of different ploidy level. *Genetica* 69:111-118.
- Miyazawa CS and Galetti Jr PM (1994) First cytogenetical studies in *Characidium* species (Pisces, Characiformes, Characidiinae). *Cytologia* 59:73-79.
- Noieto RB, Amorin AP, Vicari MR, Artoni RF and Cestari MM (2009) An unusual ZZ/ZW sex chromosome system in *Characidium* fishes (Crenuchidae, Characiformes) with the presence of rDNA sites. *J Fish Biol* 75:448-453.
- Pansonato-Alves JC, Paiva LRS, Oliveira C and Foresti F (2010a) Interspecific chromosomal divergences in the genus *Characidium* (Teleostei, Characiformes, Crenuchidae). *Neotrop Ichthyol* 8:77-86.
- Pansonato-Alves JC, Vicari MR, Oliveira C and Foresti F (2010b) Chromosomal diversification in samples of *Characidium cf. gomesi* (Teleostei, Crenuchidae). *J Fish Biol*, epub.
- Pendás AM, Morán P and Garcia-Vásquez G (1993) Ribosomal RNA genes are interspersed throughout a heterochromatic chromosome arm in Atlantic salmon. *Cytogenet Cell Genet* 63:128-130.
- Pinkel D, Straume T and Gray JW (1986) Cytogenetic analysis using quantitative, high-sensitivity, fluorescence hybridization. *Proc Natl Acad Sci USA* 83:2934-2938.
- Purdum CE (1983) Genetic engineering by the manipulation of chromosomes. *Aquaculture* 33:287-300.
- Schmid M (1982) Chromosome banding in Amphibia. VII. Analysis of the structure and variability of NORs in Anura. *Chromosoma* 87:327-344.
- Schweizer D (1976) Reverse fluorescent chromosome banding with chromomycin and DAPI. *Chromosoma* 58:307-324.
- Schweizer D and Loidl J (1987) A model for heterochromatin dispersion and the evolution of C band patterns. *Chromosome Today* 9:61-74.
- Schweizer D, Ambros P and Andrlé M (1978) Modification of DAPI banding on human chromosomes by prestaining with a DNA-binding oligopeptide antibiotic, Distamycin A. *Exp Cell Res* 111:327-332.
- Sumner AT (1972) A simple technique for demonstrating centromeric heterochromatin. *Exp Cell Res* 75:304-306.

- Suzuki H, Sakurai S and Matsuda Y (1996) Rat rDNA spacer sequences and chromosomal assignment of the genes to the extreme terminal region of chromosome 19. *Cytogenet Cell Genet* 72:1-4.
- Thorgaard GH and Allen Jr SK (1987) Chromosome manipulation and markers in fishery management. In: Ryman N and Utter F (eds) *Population Genetic and Fishery Management*. University of Washington Press, Seattle, pp 319-331.
- Thorgaard GH, Jazmin ME and Stier AR (1981) Polyploidy induced by heat shock in rainbow trout. *Trans Am Fish Soc* 110:546-550.
- Vasilev VP, Makeeva AP and Ryabov IN (1975) On the triploidy of remote hybrids of carp (*Cyprinus carpio*) with other representative Cyprinidae. *Genetika* 11:49-56.
- Venere PC and Galetti Jr PM (1985) Natural triploidy and chromosome B in the fish *Curimata modesta* (Curimatidae, Characiformes). *Rev Bras Genet* 7:681-687.
- Vicari MR, Artoni RF and Bertollo LAC (2005) Comparative cytogenetics of *Hoplias malabaricus* (Pisces, Erythrinidae): A population analysis in adjacent hydrographic basins. *Genet Mol Biol* 28:103-110.
- Vicari MR, Artoni RF, Moreira-Filho O and Bertollo LAC (2008) Diversification of a ZZ/ZW sex chromosome system in *Characidium* fish (Crenuchidae, Characiformes). *Genetica* 134:311-317.
- White TJ, Bruns T, Lee S and Taylor L (1990) Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: *PCR Protocols: A Guide to Methods and Applications*. Academic Press Inc, London/New York, pp 312-315.

Associate Editor: Yatiyo Yonenaga-Yassuda

License information: This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.