



Jornal de Pediatria

www.jped.com.br



EDITORIAL

Strengthening RSV prevention in early life through new generation strategies*



Respiratory syncytial virus (RSV) remains a leading cause of viral lower respiratory tract infection (LRTI, including bronchiolitis and pneumonia) in infants and toddlers.^{1–3} Prior to the implementation of universal RSV prophylaxis for young infants, RSV was estimated to cause globally 6.6 new million episodes of acute LRTI each year in infants < 6 months, leading to 1.4 million hospitalizations and ~ 45,000 deaths annually, accounting for 3.6% of all deaths in this age group.⁴ Over 97% of RSV-attributable deaths occur in low middle income countries (LMIC). Importantly, community based-studies have shown that a significant proportion of RSV-related deaths occur outside hospital settings, especially in regions with limited access to care.⁵ These data emphasizes the need to provide universal protection against RSV from birth and through early infancy.

Up until recently, the only strategy approved for RSV prevention was passive prophylaxis with palivizumab. Palivizumab is a neutralizing humanized monoclonal antibody (IgG₁) that targets antigenic site II located in the RSV F (fusion) protein in both the pre-F and post-F conformations, and has a half-life of 28 days, requiring monthly intramuscular administration during the RSV season. Palivizumab was approved by the FDA in 1998 for prevention of severe RSV disease in preterm infants and children with chronic lung disease (CLD), and subsequently in 2003, in those with congenital heart diseases (CHD).^{6,7} Since then, palivizumab has been implemented in more than 60 countries. Both placebo-controlled and observational comparative studies have consistently demonstrated its efficacy at preventing RSV-hospitalizations in these high-risk children, as well as an excellent safety record.⁸ Nonetheless, due to palivizumab high costs and the need for monthly administration studies have reported different challenges in the implementation and adherence to palivizumab prophylaxis in high risk-children.⁹

The study by Feitosa and Vieira published in this issue of the journal¹⁰ provides clear evidence that access and adherence to palivizumab in Brazil, remain major challenges.

Authors conducted a retrospective study including ~ 900 infants that had received palivizumab between 2008 and 2019 at the Vaccine and Immunobiological Unit in the Child and Adolescent Institute as part of the Medical School of Sao Paulo University. Their main goal was to analyze adherence to palivizumab prophylaxis and the barriers to its implementation. They found that fewer than half (44.5%) of eligible infants were completely adherent to the recommended prophylaxis regimen with palivizumab, defined as receiving the correct number of doses at the appropriate intervals. Specifically, geographic barriers and maternal sociodemographic factors — particularly younger maternal age — were associated with poor adherence, while infants with extreme prematurity and very low birth weight demonstrated somewhat better compliance. These findings underscore that structural and social determinants of health weigh as heavily as clinical eligibility in determining outcomes. In addition, the study did not find a clear association between full adherence to palivizumab and bronchiolitis-related hospitalizations, raising the question of whether partial prophylaxis may still confer protection, as reported in studies using abbreviated palivizumab dosing schedules.¹¹ Nonetheless, although some of the bronchiolitis-related hospitalizations may have been caused by RSV, the lack of RSV testing precludes confirmation, especially considering that palivizumab confers protection only against RSV, and has no activity against other respiratory viruses.

Altogether, the study by Feitosa and Vieira underscores that reliance on an effective, costly, multi-dose monthly regimen that required administration in centralized centers contributes to inequities that ultimately limits the impact of palivizumab administration high-risk children.

The timing of this study is particularly relevant. As mentioned above, while there are well-defined risk factors for severe RSV disease, the majority of children that are hospitalized for RSV LRTI have no identifiable risk factors, underscoring the critical need for universal RSV prevention.^{12–14} In the last decade the development of RSV preventive strategies has witnessed a marked resurgence, overcoming the

* See paper by Feitosa and Ventura in 101405.

setbacks of the earlier vaccine trials conducted in the mid-1960s'. On the one hand, the recognition of the impact of RSV disease globally, and on the other hand, the identification of key antigenic sites in the RSV fusion (F) protein in its pre-F conformation, have been major drivers of this notable resurgence.¹⁵ There are currently three products approved for the prevention of RSV LRTI in young infants in the United States: a maternal bivalent preF vaccine, and two extended half-life human monoclonal antibodies, nirsevimab, and clesrovimab.^{16–18} Nirsevimab is a highly potent pre-F specific mAb that targets antigenic site Ø, while clesrovimab targets antigenic site IV located in the preF and postF conformations, and has enhanced penetration into the nasal epithelial lining fluid, which is crucial for neutralizing RSV at the primary site of infection.¹⁹ Recent studies evaluating the impact of the maternal RSV vaccine and of nirsevimab in the real world have demonstrated effectiveness that exceeds that reported in the original phase-III clinical trials, reducing RSV-related hospitalizations in infants by 70%–90% during their first RSV season.^{20–22}

Nirsevimab, and maternal RSV vaccination are now approved in Brazil and offer scalable and are superior alternatives to the monthly palivizumab regimen. These strategies address adherence challenges and broaden protection to all infants including those with high-risk factors. Yet, the potential of these strategies needs to be weighed against their cost and the capacity of national health systems to implement and integrate them equitably, challenges that are specially pronounced in LMIC.^{23,24} In the U.S, both nirsevimab and the maternal RSV vaccine were approved in the summer of 2023. However, data from regional and national surveillance systems during the 2023-2024 season showed overall low uptake: only 29% of infants were protected, with 19% receiving nirsevimab and 10% covered through maternal vaccination.²⁵ When both interventions were offered coverage increased substantially to 70%–80%. Nonetheless, lower uptake was identified among infants born to non-Hispanic Black and Middle Eastern or North African mothers, underscoring that inequities persist even within high-income countries.²⁶

The study by Feitosa and Vieira highlights that preventing RSV hospitalizations requires not only effective drugs but also ensuring that children have access to them. As these new preventive strategies are introduced in Brazil and beyond, countries should strive to integrate RSV prevention within frameworks guided by equity and sustainability. Cost-effectiveness analyses, as shown in the US or Canada,²⁷ proactive maternal immunization campaigns, and integration of RSV long-acting monoclonal antibodies into national immunization schedules are key steps. Only then will we bridge the gap between scientific advances and their real-world impact. Every year 140 million infants are born worldwide. The introduction of single-dose long-acting monoclonal antibodies and maternal vaccines against RSV have been transformative; the priority now is to ensure that no infant is left behind.

Declaration of competing interest

AM reports funding from NIH, Merck and Gates foundation for research support; fees for participation in advisory boards from Merck, Pfizer, Enanta, Moderna, Astra-Zeneca and Sanofi-Pasteur; and fees for lectures from Sanofi-Pasteur

and Pfizer. O.R. has received research grants from the NIH, Gates Foundation and Merck; and fees for participation in advisory boards from Merck, Sanofi-Pasteur, Pfizer and Moderna; and fees for lectures from Pfizer, Merck, and Sanofi-Pasteur.

Editor

R. Soibelman Procianoy

References

1. Pneumonia Etiology Research for Child Health Study G. Causes of severe pneumonia requiring hospital admission in children without HIV infection from Africa and Asia: the PERCH multi-country case-control study. *Lancet*. 2019;394(10200):757–79.
2. Hall CB, Weinberg GA, Iwane MK, Blumkin AK, Edwards KM, Staat MA, et al. The burden of respiratory syncytial virus infection in young children. *N Engl J Med*. 2009;360(6):588–98.
3. Brenes-Chacon H, Eisner M, Acero-Bedoya S, Ramilo O, Mejias A. Age-specific predictors of disease severity in children with respiratory syncytial virus infection beyond infancy and through the first 5 years of age. *Pediatr Allergy Immunol*. 2024;35(2):e14083.
4. Li Y, Wang X, Blau DM, Caballero MT, Feikin DR, Gill CJ, et al. Global, regional, and national disease burden estimates of acute lower respiratory infections due to respiratory syncytial virus in children younger than 5 years in 2019: a systematic analysis. *Lancet*. 2022;399(10340):2047–64.
5. Blau DM, Baillie VL, Els T, Mahtab S, Mutevedzi P, Keita AM, et al. Deaths attributed to respiratory syncytial virus in young children in high-mortality rate settings: Report from Child Health and Mortality Prevention Surveillance (CHAMPS). *Clin Infect Dis*. 2021;73(Suppl_3):S218–28.
6. Palivizumab, a Humanized respiratory syncytial virus monoclonal antibody, reduces hospitalization from respiratory syncytial virus infection in high-risk infants. *Pediatrics*. 1998;102(3):531–7.
7. Feltes TF, Cabalka AK, Meissner HC, Piazza FM, Carlin DA, Top FH Jr, et al. Palivizumab prophylaxis reduces hospitalization due to respiratory syncytial virus in young children with hemodynamically significant congenital heart disease. *J Pediatr*. 2003;143(4):532–40.
8. Carbonell-Estrany X, Simões EAF, Bont L, Manzoni P, Zar HJ, Greenough A, et al. Twenty-five years of palivizumab: a global historic review of its impact on the burden of respiratory syncytial virus disease in children. *Expert Rev Anti Infect Ther*. 2025;23(6):359–78.
9. Jah F, Mejias A, Nzirawa T, Silveira RC, Berger A, Travan L, et al. Improving the quality of care of respiratory syncytial virus in the neonatal and pediatric populations globally. *Front Pediatr*. 2025;13:1590842.
10. Feitosa DC, Vieira SE. Challenges in the prophylaxis of severe respiratory syncytial virus infections. *J Pediatr (Rio J)*. 2025;101(5):101415.
11. Lavoie PM, Solimano A, Taylor R, Kwan E, Claydon J, Turvey SE, et al. Outcomes of respiratory syncytial virus immunoprophylaxis in infants using an abbreviated dosing regimen of Palivizumab. *JAMA Pediatr*. 2016;170(2):174–6.
12. Wang EE, Law BJ, Stephens D. Pediatric Investigators Collaborative Network on Infections in Canada (PICNIC) prospective study of risk factors and outcomes in patients hospitalized with respiratory syncytial viral lower respiratory tract infection. *J Pediatr*. 1995;126(2):212–9.

13. García CG, Bhole R, Soriano-Fallas A, Trost M, Chason R, Ramilo O, Mejias A. Risk factors in children hospitalized with RSV bronchiolitis versus non-RSV bronchiolitis. *Pediatrics*. 2010;126(6): e1453–60.
14. Curns AT, Rha B, Lively JY, Sahni LC, Englund JA, Weinberg GA, et al. Respiratory Syncytial virus-associated hospitalizations among children < 5 years old: 2016 to 2020. *Pediatrics*. 2024;153(3):e2023062574.
15. Mejias A, Ramilo O. RSV Prevention within reach for older infants and toddlers: The role of active immunization. *J Pediatric Infect Dis Soc*. 2024;13(Supplement_2):S125–30.
16. Kampmann B, Madhi SA, Munjal I, Simões EAF, Pahud BA, Llapur C, et al. Bivalent prefusion F Vaccine in pregnancy to prevent RSV illness in infants. *N Engl J Med*. 2023;388(16):1451–64.
17. Muller WJ, Madhi SA, Seoane Nuñez B, Baca Cots M, Bosheva M, Dagan R, et al. Nirsevimab for Prevention of RSV in Term and Late-Preterm Infants. *N Engl J Med*. 2023;388(16):1533–4.
18. Moulia DL, Link-Gelles R, Chu HY, Jamieson D, Brooks O, Meyer S, et al. Use of Clesrovimab for prevention of severe respiratory syncytial virus-associated lower respiratory tract infections in infants: Recommendations of the Advisory Committee on Immunization Practices - United States, 2025. *MMWR Morb Mortal Wkly Rep*. 2025;74(32):508–14.
19. Phuah JY, Maas BM, Tang A, Zhang Y, Caro L, Railkar RA, et al. Quantification of clesrovimab, an investigational, half-life extended, anti-respiratory syncytial virus protein F human monoclonal antibody in the nasal epithelial lining fluid of healthy adults. *Biomed Pharmacother*. 2023;169:115851.
20. Ares-Gómez S, Mallah N, Santiago-Pérez MI, Pardo-Seco J, Pérez-Martínez O, Otero-Barrós MT, et al. Effectiveness and impact of universal prophylaxis with nirsevimab in infants against hospitalisation for respiratory syncytial virus in Galicia, Spain: initial results of a population-based longitudinal study. *Lancet Infect Dis*. 2024;24(8):817–28. Erratum in: *Lancet Infect Dis*. 2024;24(7):e419.
21. Torres JP, Sauré D, Goic M, Thraves C, Pacheco J, Burgos J, et al. Effectiveness and impact of nirsevimab in Chile during the first season of a national immunisation strategy against RSV (NIRSE-CL): a retrospective observational study. 2025. doi: 10.1016/S1473-3099(25)00233-6. Online ahead of print.
22. Pérez Marc G, Vizzotti C, Fell DB, Di Nunzio L, Olszevicki S, Maniewicz SW, et al. Real-world effectiveness of RSVpreF vaccination during pregnancy against RSV-associated lower respiratory tract disease leading to hospitalisation in infants during the 2024 RSV season in Argentina (BERNI study): a multicentre, retrospective, test-negative, case-control study. *Lancet Infect Dis*. 2025;25(9):1044–54.
23. Barsosio HC, Bont LJ, Groome MJ, Karron RA, Kragten-Tabatabaie L, Madhi SA, et al. How Gavi support for RSV immunisation will advance health equity. *Lancet*. 2025;406(10499):127–8.
24. Terstappen J, Hak SF, Bhan A, Bogaert D, Bont LJ, Buchholz UJ, et al. The respiratory syncytial virus vaccine and monoclonal antibody landscape: the road to global access. *Lancet Infect Dis*. 2024;24(12):e747–61.
25. Boundy EO, Fast H, Jatlaoui TC, Razzaghi H, Harris L, Nguyen K, et al. Respiratory syncytial virus immunization coverage among infants through receipt of Nirsevimab monoclonal antibody or maternal vaccination - United States, October 2023-March 2024. *MMWR Morb Mortal Wkly Rep*. 2025;74(31):484–9.
26. Irving SA, Crane B, Weintraub ES, Patel SA, Razzaghi H, Daley MF, et al. Infant respiratory syncytial virus immunization coverage in the vaccine safety datalink: 2023-2024. *Pediatrics*. 2025;155(6):e2024070240.
27. Bugden S, Mital S, Nguyen HV. Cost-effectiveness of nirsevimab and maternal RSVpreF for preventing respiratory syncytial virus disease in infants across Canada. *BMC Med*. 2025;23(1):102.

Asuncion Mejias ^{*}, Octavio Ramilo 
St. Jude Children's Research Hospital, Department of Infectious Diseases, Memphis, TN, USA

^{*} Corresponding author.
E-mail: asuncion.mejias@stjude.org (A. Mejias).