



ORIGINAL ARTICLE

Alterations in the pulmonary histoarchitecture of neonatal mice exposed to hyperoxia[☆]

Renata B. Reis^a, Akinori C. Nagato^b, Clarissa R. Nardeli^c,
Isadora C.P. Matias^c, Wanderson G. Lima^d, Frank S. Bezerra^{e,*}

^a Acadêmica, Curso de Fisioterapia, Laboratório de Biomorfologia e Patologia Experimental, Centro de Ciências da Saúde, Universidade Severino Sombra (USS), Vassouras, RJ, Brasil

^b Mestre em Ciências, Professor Assistente, Laboratório de Biomorfologia e Patologia Experimental, Centro de Ciências da Saúde, Universidade Severino Sombra (USS), Vassouras, RJ, Brasil

^c Acadêmica, Curso de Biomedicina, Laboratório de Biomorfologia e Patologia Experimental, Centro de Ciências da Saúde, Universidade Severino Sombra (USS), Vassouras, RJ, Brasil

^d Doutor em Patologia, Professor Adjunto, Laboratório de Imunopatologia, Departamento de Ciências Biológicas, Universidade Federal de Ouro Preto (UFOP), Ouro Preto, MG, Brasil

^e Doutor em Ciências Morfológicas, Professor Adjunto, Laboratório de Bioquímica Metabólica, Departamento de Ciências Biológicas, Universidade Federal de Ouro Preto (UFOP), Ouro Preto, MG, Brasil

Received 8 August 2012; accepted 29 October 2012

Available online 26 April 2013

KEYWORDS

Hyperoxia;
Lung;
Newborn animals

Abstract

Objectives: To analyze the effects of exposure to hyperoxia (100% oxygen) on the lung histoarchitecture of neonatal mice.

Methods: Neonatal Balb/c mice were exposed to hyperoxia (HG) (100% oxygen) (n = 10) in a chamber (15 x 20 x 30 cm) for 24 hours with a flow of 2 L/min. The control group (CG) (n = 10) was exposed to normoxia in the same type of chamber and for the same time. After exposure, the animals were euthanized by decapitation; the lungs were removed and processed for histological examination according to the laboratory routine. Three-mm thick sections were stained with hematoxylin and eosin (H&E). The morphometric analysis was performed with in order to analyze the macrophages present in the alveolar lumen, surface density (S_v) of gas exchange, volume density (V_v) of lung parenchyma, and areas of atelectasis.

Results: A decrease in the number of alveolar macrophages ($M\emptyset$) was observed in the HG ($HG = 0.08 \pm 0.01 M\emptyset/mm^2$, $CG = 0.18 \pm 0.03 M\emptyset/mm^2$, $p = 0.0475$), S_v of gas exchange in HG ($HG = 8.08 \pm 0.12 mm^2/mm^3$, $CG = 8.65 \pm 0.20 mm^2/mm^3$, $p = 0.0233$), V_v of lung parenchyma in HG ($HG = 54.7/33.5/83.5\%/mm^2$; $CG = 75/56.7/107.9\%/mm^2$, $p < 0.0001$) when compared with the CG. However, there was an increase in areas of atelectasis in HG ($HG = 17.5/11.3/38.4 atelectasis/mm^2$, $CG = 14/6.1/24.4 atelectasis/mm^2$, $p = 0.0166$) when compared with the CG.

[☆] Please cite this article as: Reis RB, Nagato AC, Nardeli CR, Matias IC, de Lima WG, Bezerra FS. Alterations in the pulmonary histoarchitecture of neonatal mice exposed to hyperoxia. J Pediatr (Rio J). 2013;89:300–6.

* Corresponding author.

E-mail: frank@iceb.ufop.br (F.S. Bezerra).

PALAVRAS-CHAVE

Hiperóxia;
Pulmão;
Animais
recém-nascidos

Conclusion: The present results indicate that hyperoxia caused alterations in lung histoarchitecture, increasing areas of atelectasis and diffuse alveolar hemorrhage.

© 2013 Sociedade Brasileira de Pediatria. Published by Elsevier Editora Ltda. All rights reserved.

Alterações da histoarquitetura pulmonar de camundongos neonatos expostos à hiperóxia

Resumo

Objetivos: Analisar os efeitos da exposição à hiperóxia (100% de oxigênio) sobre a histoarquitetura pulmonar de camundongos neonatos.

Métodos: Camundongos neonatos da linhagem Balb/c foram expostos à hiperóxia (GH) (100% de oxigênio) (n = 10) em uma câmara (15 x 20 x 30 cm) por 24 h, com fluxo de 2 L/min. O grupo controle (GC) (n = 10) foi exposto a normóxia em um mesmo tipo de câmara e pelo mesmo tempo. Após a exposição, os animais foram sacrificados por decapitação, os pulmões foram removidos para análise histológica e processados de acordo com a rotina do laboratório. Cortes de 3 µm de espessura foram corados com hematoxilina e eosina (H&E). A análise morfométrica foi realizada com o objetivo de analisar macrófagos presentes na luz alveolar, densidade de superfície (S_v) de trocas gasosas, densidade de volume (V_v) de parênquima pulmonar e áreas de atelectasias.

Resultados: Foi verificada diminuição do número de macrófagos alveolares ($M\emptyset$) no GH ($GH = 0,08 \pm 0,01 \text{ } M\emptyset/\text{mm}^2$; $GC = 0,18 \pm 0,03 \text{ } M\emptyset/\text{mm}^2$; $p = 0,0475$), S_v de troca gasosa no GH ($GH = 8,08 \pm 0,12 \text{ mm}^2/\text{mm}^3$; $GC = 8,65 \pm 0,20 \text{ mm}^2/\text{mm}^3$; $p = 0,0233$), V_v de parênquima pulmonar no GH ($GH = 54,7/33,5/83,5\%/\text{mm}^2$; $GC = 75/56,7/107,9\%/\text{mm}^2$; $p < 0,0001$) quando comparado com o GC. Entretanto, houve aumento de áreas de atelectasias no GH ($GH = 17,5/11,3/38,4 \text{ atelectasia}/\text{mm}^2$; $GC = 14/6,1/24,4 \text{ atelectasia}/\text{mm}^2$; $p = 0,0166$) quando comparado com o GC.

Conclusão: Nossos resultados indicam que a hiperóxia promoveu alterações na histoarquitetura pulmonar, aumentando áreas de atelectasia e hemorragia alveolar difusa.

© 2013 Sociedade Brasileira de Pediatria. Publicado por Elsevier Editora Ltda. Todos os direitos reservados.

Introduction

It is estimated that 3.9 of the 10.8 million child's deaths worldwide occur in the first 28 days of life. Over 96% of these deaths occur in developing countries. Pneumonia may be associated with a low Apgar score (severe respiratory alterations at birth), which is commonly associated with chorioamnionitis, inflammation of the fetal membranes (chorion and amnion) due to a bacterial infection, usually related to prolonged vaginal deliveries and also to inhalation of infected amniotic fluid. In most cases, this leads to fetal asphyxia. One of the most obvious manifestations is hypoxemia, followed by chest indrawing and cyanosis.

As treatment, studies have shown that drug therapies using antibiotics are effective, as well as the use of oxygen in order to reverse the initial hypoxemia and reduce the risk of mortality.¹ The use of oxygen is among the first lines of therapy for hypoxemia caused by pulmonary and heart disease,²⁻⁴ which involves treating hypoxia by oxygen inhalation at a pressure greater than that of ambient air, which facilitates hematosis and reduces ventilatory work, in order to maintain adequate oxygenation with $\text{PaO}_2 > 50 \text{ mmHg}$ and $< 70 \text{ mmHg}$.⁵

Although O_2 therapy is considered essential for life, studies show an association between oxygen toxicity and retinopathy of prematurity, chronic lung disease, bronchopulmonary dysplasia, atelectasis by resorption, tracheobronchitis, depression of mucociliary activity, nausea, anorexia, headache, lung epithelial damage, damage to the blood-air barrier, and pulmonary edema. Moreover, the injury can be intensified when combined with mechanical ventilation.⁶ Recent studies suggest that the pulmonary epithelial damage induced by exposure to high concentrations of oxygen, specifically, has been associated with oxidative stress,⁷ based on the hypothesis that hyperoxia induces an increase in the number of oxygen free radicals, reactive species capable of reacting with biomolecules and causing direct damage to membrane proteins and DNA.⁸

After pulmonary epithelium lesion, in particular, there is activation of macrophages and an inflammatory cascade, followed by pulmonary edema and presence of fibrin, collagen, and neutrophilic aggregate.⁹ The literature describes animal models exposed to hyperoxia only in adult mice, when their lungs are already fully formed. The effects of high concentrations of oxygen at the time of lung formation, *i.e.*, the lungs of newborns, are yet to be clearly described in Balb/c mice. Therefore, this study aimed to evaluate the histological patterns in lungs of neonatal mice 12 hours after birth, exposed to hyperoxia for 24 hours.

Methods

Ethical aspects

The experiment was performed in accordance with the provisions of the Brazilian Society of Science in Laboratory Animals, and was approved by the Ethics Committee for Animal Research of the University.

Animals

Twenty Balb/c neonatal mice, approximately 12 hours after birth, with a mean weight of 1.5 g (despite the low weight of newborn mice, their anatomical structures are well-defined, allowing for experimental manipulation) were obtained from the Laboratory of Experimental Pathology and Biomorphology of the Centro de Ciências da Saúde (CCS) of the Universidade Severino Sombra, Brazil. The animals' nutrition in the postnatal period until euthanasia was provided by *ad libitum* breastfeeding (breastfeeding in mice lasts on average 19 to 21 days after birth).

Experimental design

Exposure to oxygen

The animals were divided into two groups: control group (CG) - mice exposed to ambient air and to the same conditions of the experimental group and the hyperoxia group (HG) - mice exposed to hyperoxia for 24 h.

For the animals exposed to hyperoxia, an acrylic inhalation chamber was used (30 cm long, 20 cm wide, and 15 cm high), as described by Nagato.¹⁰ Oxygen 100% was purchased from White Martins® (White Martins Praxair Inc. - São Paulo, Brazil). The oxygen cylinder was coupled to the oxygen inhalation chamber through a silicone conduit. The gas was released into the chamber with a constant flow of 2 L/min, thus ensuring an oxygen flow that would supply and saturate the environment.

After a period of time, when oxygen had filled the chamber space, all mice (except the control group, which inhaled ambient air) were placed in the inhalation chamber and removed after 24 h. The oxygen concentration was measured continuously through an oxygen cell (C3 - Middlesbrough, England). The mice received water and food *ad libitum*, and were kept in individual cages with controlled temperature and humidity ($21 \pm 2^\circ\text{C}$, $50 \pm 10\%$, respectively), and submitted to inverted 12-h cycles of light/dark (artificial lights, 7 p.m. to 7 a.m.).

This experimental model was designed to mimic the conditions of supplemental oxygen that neonates receive in intensive care units in the first days of life when associated with a pathological picture, during lung formation.

Euthanasia and organ removal

At the end of the O_2 exposure time, euthanasia was performed by decapitation. A ventral midline incision was performed to remove the skin, starting from the chest to the abdominal region. Access to the thoracic cavity was made using a subxiphoid incision with removal of the diaphragm

and costal osteotomy, to expose the mediastinum. After exposure of the mediastinum, a lung perfusion was performed by sectioning the left atrium, followed by a puncture in the right ventricle, injecting 1 mL of saline solution with 0.9% NaCl with constant pressure controlled by a pump (Sykam - Gewerbering, Germany). After perfusion, the lungs were removed by plucking through the mediastinum.

Histological analysis

Both lungs were fixed through a cannula inserted in the trachea by instillation of formalin (Vetec Química Fina - Duque de Caxias, Brazil) buffered (10%) with constant pressure controlled by a pump. After 48 h, they were processed through the following steps: dehydrated in increasing concentrations of ethanol (50%, 70%, 92.8%, and 99.3%), diaphonized in xylene, and paraffin-embedded. Three- μm -thick sections were stained with hematoxylin and eosin (H & E).

Morphometric analysis

Representative and proportional lung samples stained with H & E were randomly studied; 15 random fields were assessed by histological slide, under $40\times$ magnification. The analyzed section came from the lungs equally embedded in paraffin and sectioned in the same direction, aiming to analyze all of the representative parts of the entire lung and proportionally in all analyses. Analyses were performed to determine the number of macrophages present in the alveolar lumen, the volume density (V_v) of parenchyma, surface density (S_v) of gas exchange, and areas of atelectasis and erythrocytes in airspaces. Both analyses were performed by observing the microscope slide on a TV monitor (Sharp - 14") where a test system was superimposed on the screen, and the analyses were made in relation to tissues. S_v of gas exchange was verified through the test system of cycloid arcs with proportional orientation to the sine of the vertical axis angle. The measurement was performed by counting coincident points on the portion of the gas exchange surface, with the test system superimposed on the lung tissue image.¹¹

The V_v of lung parenchyma was measured through the M42 test system in an irregular arrangement of points. Analysis was conducted by overlapping the test system to an image of lung tissue, and coincident points of the lung parenchyma were counted.¹¹

Statistical analysis

To determine the sample size, a statistical power of 0.9 was adopted. The data used for this calculation were obtained from studies performed by this group,¹⁰ taking into account the expected data from larger standard error of the mean. Therefore, a resulting sample size of 10 for each group was obtained. The normality of all data was tested through the Kolmogorov-Smirnov test. Parametric data were expressed as mean \pm standard error of mean, followed by unpaired Student's *t*-test. Nonparametric data were expressed as median/minimum/maximum value, and the Mann-Whitney test was used. The difference was considered significant when $p\text{-value} < 0.05$.




Figure 1 Mean number of alveolar macrophages per microscopic field in the lung parenchyma in neonatal mice exposed to ambient air and hyperoxia for 24 h.

CG = control group, animals exposed to ambient air; HG = hyperoxia group, animals exposed to 100% oxygen for 24 hours.

* Means difference between the CG and HG, with a p -value = 0.0475 in the unpaired Student's t -test.

Results

After exposure to hyperoxia for 24 h, it was observed that the HG ($HG = 0.08 \pm 0.01 \text{ MØ/mm}^2$) showed a decrease in alveolar macrophages in the alveolar lumen ($p = 0.0475$) compared to CG ($CG = 0.18 \pm 0.03 \text{ MØ/mm}^2$) (Fig. 1).

As for morphometric analyses, a decrease in the V_v of lung parenchyma was observed in the HG ($HG = 54.7/33.5/83.5\%/mm^2$, $CG = 75/56.7/107.9\%/mm^2$, $p < 0.0001$) (Fig. 2A), and the S_v of gas exchange of HG ($HG = 8.08 \pm 0.12 \text{ mm}^2/\text{mm}^3$; $CG = 8.65 \pm 0.20 \text{ mm}^2/\text{mm}^3$, $p = 0.0193$) (Fig. 2B).

Histologically, the CG was characterized by presence of lung parenchyma of normal aspect (Fig. 3A). The HG showed diffuse parenchymal abnormalities, with varying degrees of intensity (from mild to severe). The presence of areas of atelectasis and the presence of red blood cells in the alveolar lumen, as shown in Fig. 3B, were the most frequent alterations and were significantly increased in HG ($HG = 17.5/11.3/38.4 \text{ atelectasis/mm}^2$, $CG = 14/6.1/24.4 \text{ atelectasis/mm}^2$, $p = 0.0166$) when compared to the CG (Fig. 4).

Discussion

The present study analyzed the effects of exposure to high concentrations of oxygen on lung histological patterns of neonatal Balb/c mice.

It was observed that hyperoxia induced a decrease in the number of alveolar macrophages, modified the lung histoarchitecture, and increased the number of red cells in the air spaces.

The HG showed a decrease of macrophages in the alveolar space after 24 h of exposure. In their studies, Petrache et al.¹² demonstrated *in vitro* (after 24 hour-exposure to $O_2 > 95\%$) and *in vivo* (after the mice were exposed for 72 h to 100% O_2) that alveolar macrophages undergo apoptosis when compared to macrophages in normoxia. It was also demonstrated that, in the first 30 minutes of hyperoxia, the increased activity of ERK (extracellular signal-regulated kinase) protected alveolar macrophages, decreasing the rate of apoptosis. However, the same did not occur in the period from 8 to 24 h, because ERK activity returned to normal values.

Nyunoya et al.¹³ observed that the decreased activity of phosphatases during hyperoxia, including PP2A and MKP-3, is related to ERK inhibition, which decreases macrophage survival. Similar results were observed in the present study, as there was a significant decrease in alveolar macrophages in the lung of newborn animals exposed to hyperoxia for 24 h. In another study, macrophages cultures exposed to 95%




Figure 2 Volume density (V_v) and surface density (S_v) in the pulmonary parenchyma in neonatal mice exposed to ambient air and hyperoxia for 24 h.

CG = control group (animals exposed to ambient air); HG = hyperoxia group, (animals exposed to 100% oxygen for 24 hours).

*** Means difference between CG and HG, with p -value < 0.0001 at Mann-Whitney test.

* Means difference enters the CG and HG, with p -value = 0.0193 at unpaired Student's t -test.




Figure 3 Photomicrographs representative of histological sections of lung parenchyma of neonatal mice exposed to ambient air and hyperoxia for 24 h. A V_v decrease can be observed in the lung parenchyma, gas exchange S_v , and atelectasis increase in HG (b) compared with the CG (a). Hematoxylin and eosin (H & E).

oxygen (hyperoxia) showed reduced proliferation when compared to cultures exposed to 21% oxygen (normoxia). This reduction was probably mediated by actin polymerization induced by oxidative stress, which altered the phagocytic capacity to pathogens.¹⁴ Thus, it is suggested that hyperoxia may influence both the increase in apoptosis and the decrease in proliferation of alveolar macrophages.

Another study by Thébaud et al.¹⁵ demonstrated that exposure to oxygen therapy at high concentrations interferes with the development of lung parenchyma, as newborn rats had a lower expression of vascular endothelial growth factor (VEGF) and, consequently, a decrease in the number of blood capillaries, which resulted in increased air spaces. Mascaretti et al.¹⁶ also reported this decrease in the number of alveoli in an experimental model of exposure to hyperoxia in preterm rabbits of the New Zealand

lineage. Animal models have demonstrated structural lung abnormalities resulting from exposure to hyperoxia.^{17,18}

Neonates are subject to alterations caused by oxygen exposure, since their antioxidant system develops later. These alterations make the neonate vulnerable to such lesions, including parenchymatous lesions, which may be irreversible.¹⁹ Dauger et al.²⁰ studied mice exposed to hyperoxia at 65% over a period of 28 days after birth, demonstrating a smaller number of alveoli, albeit with increased alveolar lumen. The alterations lasted for seven months after exposure, evidencing that hyperoxia causes permanent alterations in lung structure. Neonatal mice are at the saccular stage of lung development, and decreased alveolarization is a prevalent characteristic.²¹ This pattern was demonstrated in the present study. However, exposure to hyperoxia exacerbated the decrease in volume density of the lung parenchyma and gas exchange surface area, compared to animals exposed to ambient air.

In clinical practice, atelectasis is often found during general anesthesia, especially in the postoperative period and/or during mechanical ventilation.⁶ The present results indicate that exposure to hyperoxia for 24 h resulted in an increase in areas of pulmonary atelectasis. This can be explained by the induction of atelectasis by resorption, a mechanism responsible for impairment of gas exchange and of structural lung parenchyma.²² Loewen et al.²³ studied rabbits of the New Zealand lineage and demonstrated the beneficial effect of supplementation of exogenous surfactant in lungs exposed to hyperoxia. In their study, animals exposed to hyperoxia at 100% associated with surfactant supplementation presented a decrease in areas of atelectasis, when compared to animals exposed to hyperoxia alone. This suggests that reduction in surfactant production induced by high doses of oxygen promotes increased areas of atelectasis, which was also confirmed by Buonocore et al.²⁴

Studies in experimental animals have demonstrated that hyperoxia promotes increased capillary permeability, extravasation of plasma proteins into the interstitium and alveolar space, and later, after a prolonged exposure, fibrosis in the alveolar wall.²⁵ Tokieda et al.²⁶ studied mice with deficiency of surfactant protein B (SP-B) exposed to hyperoxia at 95%, and observed a susceptibility to pulmonary congestion and bleeding. However, Lian et al.²⁷




Figure 4 Mean number of alveoli with atelectasis per microscopic field in the lung parenchyma of neonatal mice exposed to ambient air and hyperoxia for 24 h.

CG, control group (animals exposed to ambient air); HG, hyperoxia group, (animals exposed to 100% oxygen for 24 h).

* Means the difference between CG and HG with p-value = 0.0166 in Mann-Whitney test.

demonstrated the protective effects against damage caused by exposure to oxygen therapy. In that study, transgenic animals overexpressing signal transducer and activator of transcription 3 (Stat3C) and, consequently, with an increased production of SP-B protein, presented resistance to alveolar hemorrhage. Alveolar hemorrhage,²⁰ non-cardiogenic pulmonary edema,²⁸ and damage to type I pneumocytes²⁷ and type II pneumocyte hyperplasia have been mentioned as alterations resulting from high oxygen concentrations in clinical practice.²⁹

The stages of lung structural development are similar in humans and mice. In the mouse, after the ninth day of gestational development, lung formation begins, characterized by embryonic events that depend on the interaction between epithelial and mesenchymal cells. The 12-hour postnatal period, which was chosen for the start of the present intervention, is described as crucial for the development of histological and biochemical alterations that can be evaluated in this experimental model. Moreover, at this time, the animals are at the intermediate saccular period of lung development and lung structures are significantly formed.³⁰ The objective of the present study was to investigate how the developing lung would be able to respond to exposure to oxygen at high concentrations, considering that in clinical practice newborns receive such treatment (supplemental oxygen therapy) in intensive care units.

The limitation to mimic the time and intensity of oxygen administration in experimental models is due to the lack of clinical studies that indicate the mean time and mean fraction of inspired oxygen used by newborns during the hospitalization period. However, this study stimulates the development of other clinical and experimental findings, for example, feasibility studies and specific markers of apoptosis for alveolar macrophages, all with the aim of achieving therapeutic alternatives for the treatment of medical exposure to oxygen at high concentrations.

Conflicts of interest

The authors declare no conflicts of interest.

Acknowledgements

To FAPERJ for the scientific initiation grant to undergraduate student Renata Reis and to FAPEMIG for the postdoctoral grant to Professor Frank Silva Bezerra DECBI/UFOP through the project approved by Edict 15/2010 "Programa Primeiros Projetos".

References

1. Duke T. Neonatal pneumonia in developing countries. *Arch Dis Child Fetal Neonatal Ed.* 2005;90:F211–9.
2. Frey B, Shann F. Oxygen administration in infants. *Arch Dis Child Fetal Neonatal Ed.* 2003;88:F84–8.
3. Duke T, Graham SM, Cherian MN, Ginsburg AS, English M, Howie S, et al. Oxygen is an essential medicine: a call for international action. *Int J Tuberc Lung Dis.* 2010;14:1362–8.
4. O'Reilly MA. DNA damage and cell cycle checkpoints in hyperoxic lung injury: braking to facilitate repair. *Am J Physiol Lung Cell Mol Physiol.* 2001;281:L291–305.
5. Usen S, Weibert M. Clinical signs of hypoxaemia in children with acute lower respiratory infection: indicators of oxygen therapy. *Int J Tuberc Lung Dis.* 2001;5:505–10.
6. Reissmann H, Böhm SH, Suárez-Sipmann F, Tusman G, Buschmann C, Maisch S, et al. Suctioning through a double-lumen endotracheal tube helps to prevent alveolar collapse and to preserve ventilation. *Intensive Care Med.* 2005;31:431–40.
7. Hay Jr WW, Bell EF. Oxygen therapy, oxygen toxicity, and the STOP-ROP trial. *Pediatrics.* 2000;105:424–5.
8. Kim Y, Kim H, Yoo HY, Kang JS, Kim SJ, Kim JK, et al. Suppression of CFTR-mediated Cl secretion of airway epithelium in vitamin C-deficient mice. *J Korean Med Sci.* 2011;26:317–24.
9. Minamino T, Komuro I. Regeneration of the endothelium as a novel therapeutic strategy for acute lung injury. *J Clin Invest.* 2006;116:2316–9.
10. Nagato AC, Bezerra FS, Lanzetti M, Lopes AA, Silva MA, Porto LC, et al. Time course of inflammation, oxidative stress and tissue damage induced by hyperoxia in mouse lungs. *Int J Exp Pathol.* 2012;93:269–78.
11. Mandarim-de-Lacerda CA. Stereological tools in biomedical research. *An Acad Bras Cienc.* 2003;75:469–86.
12. Petrache I, Choi ME, Otterbein LE, Chin BY, Mantell LL, Horowitz S, et al. Mitogen-activated protein kinase pathway mediates hyperoxia-induced apoptosis in cultured macrophage cells. *Am J Physiol.* 1999;277:L589–95.
13. Nyunoya T, Monick MM, Powers LS, Yarovinsky TO, Hunninghake GW. Macrophages survive hyperoxia via prolonged ERK activation due to phosphatase down-regulation. *J Biol Chem.* 2005;280:26295–302.
14. O'Reilly PJ, Hickman-Davis JM, Davis IC, Matalon S. Hyperoxia impairs antibacterial function of macrophages through effects on actin. *Am J Respir Cell Mol Biol.* 2003;28:443–50.
15. Thébaud B, Ladha F, Michelakis ED, Sawicka M, Thurston G, Eaton F, et al. Vascular endothelial growth factor gene therapy increases survival, promotes lung angiogenesis, and prevents alveolar damage in hyperoxia-induced lung injury: evidence that angiogenesis participates in alveolarization. *Circulation.* 2005;112:2477–86.
16. Mascaretti RS, Mataloun MM, Dolnikoff M, Rebello CM. Lung morphometry, collagen and elastin content: changes after hyperoxic exposure in preterm rabbits. *Clinics (Sao Paulo).* 2009;64:1099–104.
17. Wispe JR, Roberts RJ. Molecular basis of pulmonary oxygen toxicity. *Clin Perinatol.* 1987;14:651–66.
18. Crapo JD. Morphologic changes in pulmonary oxygen toxicity. *Annu Rev Physiol.* 1986;48:721–31.
19. Monte LF, Silva Filho LV, Miyoshi MH, Rozov T. Displasia broncopulmonar. *J Pediatr (Rio J).* 2005;81:99–110.
20. Dauger S, Ferkdadjji L, Saumon G, Vardon G, Peuchmaur M, Gaultier C, et al. Neonatal exposure to 65% oxygen durably impairs lung architecture and breathing pattern in adult mice. *Chest.* 2003;123:530–8.
21. Rogers LK, Tipple TE, Nelin LD, Welty SE. Differential responses in the lungs of newborn mouse pups exposed to 85% or > 95% oxygen. *Pediatr Res.* 2009;65:33–8.
22. Carvalho CR, de Paula Pinto Schettino G, Maranhão B, Bethlem EP. Hyperoxia and lung disease. *Curr Opin Pulm Med.* 1998;4:300–4.
23. Loewen GM, Holm BA, Milanowski L, Wild LM, Notter RH, Matalon S. Alveolar hyperoxic injury in rabbits receiving exogenous surfactant. *J Appl Physiol.* 1989;66:1087–92.
24. Buonocore G, Perrone S, Tataranno ML. Oxygen toxicity: chemistry and biology of reactive oxygen species. *Semin Fetal Neonatal Med.* 2010;15:186–90.
25. Yamada M, Kubo H, Kobayashi S, Ishizawa K, Sasaki H. Interferon-gamma: a key contributor to hyperoxia-induced

- lung injury in mice. *Am J Physiol Lung Cell Mol Physiol*. 2004;287:L1042-7.
26. Tokieda K, Iwamoto HS, Bachurski C, Wert SE, Hull WM, Ikeda K, et al. Surfactant protein-B-deficient mice are susceptible to hyperoxic lung injury. *Am J Respir Cell Mol Biol*. 1999;21:463-72.
27. Lian X, Qin Y, Hossain SA, Yang L, White A, Xu H, et al. Overexpression of Stat3C in pulmonary epithelium protects against hyperoxic lung injury. *J Immunol*. 2005;174:7250-6.
28. Song Y, Fukuda N, Bai C, Ma T, Matthay MA, Verkman AS. Role of aquaporins in alveolar fluid clearance in neonatal and adult lung, and in oedema formation following acute lung injury: studies in transgenic aquaporin null mice. *J Physiol*. 2000;525:771-9.
29. Babu PB, Chidekel A, Shaffer TH. Hyperoxia-induced changes in human airway epithelial cells: the protective effect of perflubron. *Pediatr Crit Care Med*. 2005;6:188-94.
30. Minoo P, King RJ. Epithelial-mesenchymal interactions in lung development. *Annu Rev Physiol*. 1994;56:13-45.



ERRATUM

Erratum of “Alterations in the pulmonary histoarchitecture of neonatal mice exposed to hyperoxia”

Renata B. Reis^a, Akinori C. Nagato^b, Clarissa R. Nardeli^c, Isadora C.P. Matias^c, Wanderson G. Lima^d, Frank S. Bezerra^{e,*}

^a Acadêmica, Curso de Fisioterapia, Laboratório de Biomorfologia e Patologia Experimental, Centro de Ciências da Saúde, Universidade Severino Sombra (USS), Vassouras, RJ, Brasil

^b Mestre em Ciências, Professor Assistente, Laboratório de Biomorfologia e Patologia Experimental, Centro de Ciências da Saúde, Universidade Severino Sombra (USS), Vassouras, RJ, Brasil

^c Acadêmica, Curso de Biomedicina, Laboratório de Biomorfologia e Patologia Experimental, Centro de Ciências da Saúde, Universidade Severino Sombra (USS), Vassouras, RJ, Brasil

^d Doutor em Patologia, Professor Adjunto, Laboratório de Imunopatologia, Departamento de Ciências Biológicas, Universidade Federal de Ouro Preto (UFOP), Ouro Preto, MG, Brasil

^e Doutor em Ciências Morfológicas, Professor Adjunto, Laboratório de Bioquímica Metabólica, Departamento de Ciências Biológicas, Universidade Federal de Ouro Preto (UFOP), Ouro Preto, MG, Brasil

In the original article “Alterations in the pulmonary histoarchitecture of neonatal mice exposed to hyperoxia” (J Pediatr (Rio J). 2013;89(3):300–306), in affiliation ‘e’, where it reads *Laboratório de Medicina Metabólica*, it should read *Laboratório de Bioquímica Metabólica*.

Doi: 10.1016/j.jpmedp.2012.10.010

DOI of refers to article: <http://dx.doi.org/10.1016/j.jpmedp.2012.10.010>

* Corresponding author.

E-mail: frank@iceb.ufop.br (F.S. Bezerra).