

# ***Candidatus Rickettsia andeanae*, a spotted fever group agent infecting *Amblyomma parvum* ticks in two Brazilian biomes**

**Fernanda Aparecida Nieri-Bastos<sup>1</sup>, Marcos Gomes Lopes<sup>1</sup>, Paulo Henrique Duarte Cançado<sup>2</sup>, Giselle Ayres Razera Rossa<sup>1</sup>, João Luiz Horácio Faccini<sup>3</sup>, Solange Maria Gennari<sup>1</sup>, Marcelo Bahia Labruna<sup>1/+</sup>**

<sup>1</sup>Faculdade de Medicina Veterinária e Zootecnia, Universidade de São Paulo, São Paulo, SP, Brasil

<sup>2</sup>Embrapa Gado de Corte, Campo Grande, MS, Brasil <sup>3</sup>Universidade Federal Rural do Rio de Janeiro, Seropédica, RJ, Brasil

*Adult ticks of the species Amblyomma parvum were collected from the vegetation in the Pantanal biome (state of Mato Grosso do Sul) and from horses in the Cerrado biome (state of Piauí) in Brazil. The ticks were individually tested for rickettsial infection via polymerase chain reaction (PCR) targeting three rickettsial genes, gltA, ompA and ompB. Overall, 63.5% (40/63) and 66.7% (2/3) of A. parvum ticks from Pantanal and Cerrado, respectively, contained rickettsial DNA, which were all confirmed by DNA sequencing to be 100% identical to the corresponding fragments of the gltA, ompA and ompB genes of Candidatus Rickettsia andeanae. This report is the first to describe Ca. R. andeanae in Brazil.*

Key words: ticks - *Amblyomma parvum* - *Candidatus Rickettsia andeanae*

The bacterial genus *Rickettsia* (Rickettsiales: Rickettsiaceae) includes Gram-negative coccobacilli organisms in obligatory associations with eukaryote cells. A number of *Rickettsia* species genetically classified into the spotted fever group (SFG) are agents of tick-borne diseases in the world (Parola et al. 2005). In Brazil, the most important SFG agents are *Rickettsia rickettsii*, the causative agent of Brazilian spotted fever, which is the most deadly rickettsiosis globally (Guedes et al. 2005, Labruna 2009), and the strain Atlantic rainforest, an *Rickettsia parkeri*-like agent that was only recently shown to cause spotted fever in humans (Spolidorio et al. 2010, Silva et al. 2011). Other tick-associated rickettsiae, namely *Rickettsia rhipicephali*, *Rickettsia amblyommii*, *Rickettsia monteiroi* and *Rickettsia bellii*, have also been reported in Brazil, but have not been associated with human illness (reviewed by Labruna et al. 2011). It is noteworthy that during the last three decades, approximately 10 SFG *Rickettsia* species or subspecies previously known to infect only ticks were further identified as emerging agents of tick-borne rickettsioses throughout the world (Parola et al. 2005, Shapiro et al. 2010). An outstanding example is *R. parkeri*, which has been known to infect *Amblyomma* ticks for more than 60 years, but was recognised as a human pathogen only during the last 10 years (Paddock et al. 2004). Based on these findings, it is prudent to consider any new tick-associated *Rickettsia* species as a potential human pathogen until further studies are conducted.

The tick species *Amblyomma parvum* is distributed from southern Mexico to Argentina (Guglielmone et al. 2003). While the *A. parvum* adult stage has a preference for parasitising medium to large-sized mammals (ruminants, horses and carnivores), larvae and nymphs seem to prefer small mammals (Aragão 1936, Nava et al. 2008). Recent studies in Argentina and Brazil have indicated that rodents of the genera *Galea* and *Trichomys* are important hosts for *A. parvum* sub-adults (Nava et al. 2008, Horta et al. 2011). Of note, the *A. parvum* adult stage is an important human-biting tick in Argentina and Brazil (Guglielmone et al. 2006); therefore, it is a potential pathogen vector for humans.

To evaluate rickettsial infection in *A. parvum* ticks from Brazil, tick specimens were collected in two Brazilian biomes. In the Pantanal biome, free-living ticks were collected by CO<sub>2</sub> traps set on the vegetation in the Nhumirim Farm (18°59'S 56°39'W) at the Nhecolândia Pantanal, state of Mato Grosso do Sul, central-western Brazil, from March 2006-January 2007, as detailed previously (Cançado et al. 2008). In the *Cerrado* biome, ticks were collected from horses on a farm (04°76'91''S 42°58'44''W) in the municipality of Teresina, state of Piauí (PI), northeastern Brazil, during August 2010. The ticks were preserved in absolute ethanol until they were processed in the laboratory.

Each adult *A. parvum* tick was subjected to DNA extraction by the guanidine isothiocyanate-phenol technique, as previously described (Sangioni et al. 2005). Purified DNA was stored at -20°C until it was used as template for polymerase chain reaction (PCR) amplifications. Five microlitres of each tick DNA template (approximately 500 ng of DNA) was used for PCR using the primers CS-78 (forward) GCAAGTATCGGTGAGGAT-GTAAT and CS-323 (reverse) GCTTCCTTAAAT-TCAATAAATCAGGAT, which amplify a 398-bp fragment of the citrate synthase gene (*gltA*) of, most likely, all

doi: 10.1590/0074-0276140283

Financial support: FAPESP, CNPq, CAPES

+ Corresponding author: labruna@usp.br

Received 28 May 2013

Accepted 27 November 2013

known *Rickettsia* species (Labruna et al. 2004). If an expected product was observed following gel electrophoresis, the tick was tested using two other PCR protocols: one targeting a fragment of the rickettsial 190-kDa outer membrane protein gene (*ompA*) using primers Rr190.70F ATGGCGAATATTTCTCCAAAA and Rr190.701R GT-TCCGTTAATGGCAGCATCT (Eremeeva et al. 2006) and the other targeting a fragment of the rickettsial 135-kDa outer membrane protein gene (*ompB*) using primers 120-M59 CCGCAGGGTTGGTAAGTGC and 120-807 CCTTTTAGATTACCGCCTAA (Roux & Raoult 2000). Ten microlitres of each PCR product was electrophoretically separated in a 1.5% agarose gel stained with ethidium bromide and examined using ultraviolet transillumination. PCR products compatible with the expected band size were purified using ExoSap (USB Corporation, Cleveland, OH, USA) and sequenced in an automated sequencer (model ABI Prism 310 Genetic, Applied Biosystems, Perkin Elmer, CA, USA) according to the manufacturer's protocol. The partial sequences that were obtained were subjected to BLAST analyses (ncbi.nlm.nih.gov/blast) to determine the closest similarities to other *Rickettsia* species available in GenBank.

Sixty-three (37 males, 26 females) out of 81 *A. parvum* adult specimens collected from the vegetation in the Pantanal were tested in this study. From the *Cerrado* area of PI, 3 *A. parvum* specimens (2 males, 1 female) were collected from three out of 27 examined horses; other tick species (not tested in this study) collected from these horses were *Amblyomma cajennense* (8 adults) and *Dermacentor nitens* (570 adults). The *gltA*-PCR indicated that 40 (63.5%) *A. parvum* specimens (20 males, 20 females) from Pantanal and two *A. parvum* males (66.7%) from PI were infected by *Rickettsia*. From these 42 *gltA*-PCR positive ticks, 23 were positive in the *ompA*-PCR and 17 were also positive in the *ompB*-PCR. High-quality DNA sequences were obtained from 19 *gltA*-positive ticks, 12 *ompA*-positive ticks and 11 *ompB*-positive ticks, including the two males from PI. For each rickettsial gene, the sequences from different individual ticks were identical to each other and were also 100% identical to the corresponding sequences of *Candidatus Rickettsia andeanae* from the United States of America (USA), Peru and Argentina, (e.g., GU169051, EF451001 for *gltA*; EU826513, EF372578 for *ompA*; GU395297, EF451003 for *ompB*) when submitted to BLAST analyses. The sequences generated in this study have been submitted to GenBank under the accessions KF030931-KF030933.

This study reports the identification of the SFG rickettsial agent *Ca. R. andeanae* for the first time in Brazil. Taxonomic identification of the agent was confirmed by sequencing fragments of one conserved housekeeping gene (*gltA*) and two highly polymorphic outer membrane protein genes (*ompA* and *ompB*). Among the 42 ticks that were PCR-positive for the *gltA* gene, only 23 were positive in the *ompA*-PCR and 17 in the *ompB*-PCR. Because fragments of approximately 398, 630 and 820 bp were targeted by the *gltA*, *ompA* and *ompB*-PCR protocols, respectively, the different results obtained with these protocols are possibly related to different PCR sensitivities, as it is well known that the shorter the PCR fragment, the more efficient the PCR reaction (Bustin 2000).

*Ca. R. andeanae* has been reported to infect ticks in Peru, USA, Argentina and Chile (Blair et al. 2004, Pacheco et al. 2007, Paddock et al. 2010, Abarca et al. 2012). More recently, Luce-Fedrow et al. (2012) and Ferrari et al. (2013) reported evidence for the continuous in vitro cultivation of *Ca. R. andeanae*. A number of different tick species have been reported to harbour *Ca. R. andeanae in nature*. These species include *Amblyomma maculatum*, *Ixodes boliviensis* and *Rhipicephalus sanguineus* in Peru (Blair et al. 2004, Flores-Mendoza et al. 2013), *A. maculatum* in USA (Paddock et al. 2010, Luce-Fedrow et al. 2012), *A. parvum* and *Amblyomma pseudoconcolor* in Argentina (Pacheco et al. 2007, Tomassone et al. 2010) and *Amblyomma triste* in Chile (Abarca et al. 2012). According to these references, different designations have been given to *Ca. R. andeanae*, such as *Rickettsia* sp. strain Argentina or the rickettsial endosymbiont of *A. maculatum*. Because the DNA sequences representing these different designations were 100% identical to each other, they are all considered to represent *Ca. R. andeanae* (Paddock et al. 2010, Luce-Fedrow et al. 2012, Ferrari et al. 2013), which was first reported in Peru by Blair et al. (2004). Because *Ca. R. andeanae*-infected ticks were found in distant areas of two Brazilian biomes, it is possible that the distribution of this organism within Brazil is much broader than is reported here.

The majority of the ticks in this study were infected with *Ca. R. andeanae*. This finding is in accordance with previous studies that reported that most of the *A. parvum* ticks from Argentina (69.2%) were infected with *Ca. R. andeanae*. However, infection rates by this agent among *A. maculatum* ticks in USA have been much lower, usually between 1-10% (Paddock et al. 2010, Jiang et al. 2012). Regardless, the role of *Ca. R. andeanae* as a human pathogen is unknown. Because *A. parvum* is an important human-biting tick in South America (Guglielmo et al. 2006), further studies are needed to evaluate the capacity of *Ca. R. andeanae* to be tick transmitted and to infect humans or other vertebrate hosts.

## REFERENCES

- Abarca K, López J, Acosta-Jamett G, Lepe P, Soares JF, Labruna MB 2012. A third *Amblyomma* species and the first tick-borne *Rickettsia* in Chile. *J Med Entomol* 49: 219-222.
- Aragão HB 1936. Ixodidas brasileiros e de alguns paizes limitrophes. *Mem Inst Oswaldo Cruz* 31: 759-843.
- Blair PJ, Jiang J, Schoeler GB, Moron C, Anaya E, Cespedes M, Cruz C, Felices V, Guevara C, Mendoza L, Villaseca P, Sumner JW, Richards AL, Olson JG 2004. Characterization of spotted fever group rickettsiae in flea and tick specimens from northern Peru. *J Clin Microbiol* 42: 4961-4967.
- Bustin SA 2000. Absolute quantification of mRNA using real-time reverse transcription polymerase chain reaction assays. *J Mol Endocrinol* 25: 169-193.
- Cançado PH, Piranda EM, Mourão GM, Faccini JL 2008. Spatial distribution and impact of cattle-raising on ticks in the Pantanal region of Brazil by using the CO<sub>2</sub> tick trap. *Parasitol Res* 103: 371-377.
- Eremeeva ME, Bosserman EA, Demma LJ, Zambrano ML, Blau DM, Dasch GA 2006. Isolation and identification of *Rickettsia massiliae* from *Rhipicephalus sanguineus* ticks collected in Arizona. *Appl Environ Microbiol* 72: 5569-5577.

- Ferrari FAG, Goddard J, Moraru GM, Smith WEC, Varela-Stokes AS 2013. Isolation of “*Candidatus Rickettsia andeanae*” (Rickettsiales: Rickettsiaceae) in embryonic cells of naturally infected *Amblyomma maculatum* (Ixodida: Ixodidae). *J Med Entomol* 50: 1118-1125.
- Flores-Mendoza C, Florin D, Felices V, Pozo EJ, Graf PC, Burrus RG, Richards AL 2013. Detection of *Rickettsia parkeri* from within Piura, Peru, and the first reported presence of *Candidatus Rickettsia andeanae* in the tick *Rhipicephalus sanguineus*. *Vector Borne Zoonotic Dis* 13: 505-508.
- Guedes E, Leite RC, Prata MCA, Pacheco RC, Walker DH, Labruna MB 2005. Detection of *Rickettsia rickettsii* in the tick *Amblyomma cajennense* in a new Brazilian spotted fever-endemic area in the state of Minas Gerais. *Mem Inst Oswaldo Cruz* 100: 841-845.
- Guglielmone AA, Beati L, Barros-Battesti DM, Labruna MB, Nava S, Venzal JM, Mangold AJ, Szabo MP, Martins JR, Gonzalez-Acuna D, Estrada-Pena A 2006. Ticks (Ixodidae) on humans in South America. *Exp Appl Acarol* 40: 83-100.
- Guglielmone AA, Estrada-Peña A, Keirans JE, Robbins RG 2003. *Ticks (Acari: Ixodida) of the Neotropical zoogeographic region*, Universiteit Utrecht/J Bovy-Verbeek, Utrecht, 173 pp.
- Horta MC, Nascimento GF, Martins TF, Labruna MB, Machado LCP, Nicola PA 2011. Ticks (Acari: Ixodida) parasitizing free-living wild animals in the *Caatinga* biome in the state of Pernambuco, northeastern Brazil. *Syst Appl Acarol* 16: 207-211.
- Jiang J, Stromdahl EY, Richards AL 2012. Detection of *Rickettsia parkeri* and *Candidatus Rickettsia andeanae* in *Amblyomma maculatum* Gulf Coast ticks collected from humans in the United States. *Vector Borne Zoonotic Dis* 12: 175-182.
- Labruna MB 2009. Ecology of *Rickettsia* in South America. *Ann N Y Acad Sci* 1166: 156-166.
- Labruna MB, Mattar S, Nava S, Bermudez SM, Dolz G, Abarca K, Romero L, Oteo J, Zavala-Castro J 2011. Rickettsiosis in Latin America, Caribbean, Spain and Portugal. *Rev MVZ Cordoba* 16: 2435-2457.
- Labruna MB, Whitworth T, Horta MC, Bouyer DH, McBride JW, Pinter A, Popov V, Gennari SM, Walker DH 2004. *Rickettsia* species infecting *Amblyomma cooperi* ticks from an area in the state of São Paulo, Brazil, where Brazilian spotted fever is endemic. *J Clin Microbiol* 42: 90-98.
- Luce-Fedrow A, Wright C, Gaff HD, Sonenshine DE, Hynes WL, Richards AL 2012. In vitro propagation of *Candidatus Rickettsia andeanae* isolated from *Amblyomma maculatum*. *FEMS Immunol Med Microbiol* 64: 74-81.
- Nava S, Mangoldi AJ, Guglielmone AA 2008. Aspects of the life cycle of *Amblyomma parvum* (Acari: Ixodidae) under natural conditions. *Vet Parasitol* 156: 270-276.
- Pacheco RC, Moraes-Filho J, Nava S, Brandao PE, Richtzenhain LJ, Labruna MB 2007. Detection of a novel spotted fever group rickettsia in *Amblyomma parvum* ticks (Acari: Ixodidae) from Argentina. *Exp Appl Acarol* 43: 63-71.
- Paddock CD, Fournier PE, Sumner JW, Goddard J, Elshenawy Y, Metcalfe MG, Loftis AD, Varela-Stokes A 2010. Isolation of *Rickettsia parkeri* and identification of a novel spotted fever group *Rickettsia* sp. from Gulf Coast ticks (*Amblyomma maculatum*) in the United States. *Appl Environ Microbiol* 76: 2689-2696.
- Paddock CD, Sumner JW, Comer JA, Zaki SR, Goldsmith CS, Goddard J, McLellan SLF, Tammenga CL, Ohl CA 2004. *Rickettsia parkeri*: a newly recognized cause of spotted fever rickettsiosis in the United States. *Clin Infect Dis* 38: 805-811.
- Parola P, Davoust B, Raoult D 2005. Tick and flea-borne rickettsial emerging zoonoses. *Vet Res* 36: 469-492.
- Roux V, Raoult D 2000. Phylogenetic analysis of members of the genus *Rickettsia* using the gene encoding the outer membrane protein rOmpB (*ompB*). *Int J Syst Evol Microbiol* 50: 1449-1455.
- Sangioni LA, Horta MC, Vianna MCB, Gennari SM, Soares RM, Galvão MAM, Schumaker TTS, Ferreira F, Vidotto O, Labruna MB 2005. Rickettsial infection in animals and Brazilian spotted fever endemicity. *Emerg Infect Dis* 11: 265-270.
- Shapiro MR, Fritz CL, Tait K, Paddock CD, Nicholson WL, Abramowicz KF, Karpathy SE, Dasch GA, Sumner JW, Adem PV, Scott JJ, Padgett KA, Zaki SR, Eremeeva ME 2010. *Rickettsia* 364D: a newly recognized cause of eschar-associated illness in California. *Clin Infect Dis* 50: 541-548.
- Silva N, Eremeeva ME, Rozental T, Ribeiro GS, Paddock CD, Ramos EA, Favacho AR, Reis MG, Dasch GA, de Lemos ER, Ko AI 2011. Eschar-associated spotted fever rickettsiosis, Bahia, Brazil. *Emerg Infect Dis* 17: 275-278.
- Spolidorio MG, Labruna MB, Mantovani E, Brandao PE, Richtzenhain LJ, Yoshinari NH 2010. Novel spotted fever group rickettsiosis, Brazil. *Emerg Infect Dis* 16: 521-523.
- Tomassone L, Nuñez P, Ceballos LA, Gürtler RE, Kitron U, Farber M 2010. Detection of “*Candidatus Rickettsia* sp. strain Argentina” and *Rickettsia bellii* in *Amblyomma* ticks (Acari: Ixodidae) from Northern Argentina. *Exp Appl Acarol* 52: 93-100.