

Asymptomatic *Leishmania chagasi* Infection in Relatives and Neighbors of Patients with Visceral Leishmaniasis

Argemiro D'Oliveira Júnior, Sérgio Ricardo M Costa, Aurinha Bispo Barbosa*, Maria de La Glória Orge Orge, Edgar M Carvalho/+

Laboratório de Imunologia, Hospital Universitário Prof. Edgar Santos, Rua João das Botas s/nº, 40110-160 Salvador, BA, Brasil *Posto de Saúde de Monte Gordo, Camaçari, BA, Brasil

The frequency of asymptomatic infection among relatives and neighbors of cases of visceral leishmaniasis (VL) was compared and characterization of the immunological response in these subjects was performed. Cases were from a new endemic area, close to the beach and near Salvador, capital of the State of Bahia, Brazil. The characterization of asymptomatic infection was made using a skin reaction test and detection of antibody to Leishmania chagasi by the ELISA test. To characterize the immunological response of these subjects with asymptomatic L. chagasi infection the cytokines profile and the lymphoproliferative response were determined after stimulation of lymphocytes by L. chagasi antigen. There was no difference in the frequency of L. chagasi infection in relatives (45%) and in neighbors (27%) of cases of VL (P>0.05). The immunological response from these subjects was characterized by high production of IFN- γ and a low production of IL-10 and a good lymphoproliferative response to L. chagasi antigen.

Key words: visceral leishmaniasis - asymptomatic infection - relatives' infections

There is evidence that in endemic areas of visceral leishmaniasis (VL) only about 20% of the subjects infected by *Leishmania chagasi* will develop classical VL. The majority of the infected individuals have a sub-clinical infection that may remain completely asymptomatic or have an oligosymptomatic form of the disease. Subjects with oligosymptomatic infection may develop clinical VL months after the seroconversion or may self heal their infections one or two years after the seroconversion (Badaró et al. 1986a). In Brazil the majority of patients who develop VL are malnourished children with mean age of 3 years (Badaró et al. 1986b). Depression of cellular immune response is also involved in the development of the disease since infected subjects whose lymphocytes do not proliferate and do not produce IFN- γ upon leishmania stimulus are in the group who will progress from the infection to disease (Carvalho et al. 1992). The evidence of VL cluster in families could be an indication that genetic factors may predispose the development of the disease (Evans et al. 1992). Alternatively a great number of cases of VL in one family may be related to increased exposure since *L. chagasi* transmission is predominantly

peridomiciliar. It was demonstrated an intrafamilial pattern of infection in VL in an endemic area (Jacobina, BA, Brazil) with a strong suggestion of at least partial genetic involvement (Cabello et al. 1995).

The majority of the cases of VL in Brazil occur in communities localized in the interior of the Northeastern region of the country. In 1989, an outbreak of VL was reported in villages of the municipality of Camaçari, approximately 2 km from the beach and 50 km north of Salvador, the capital of the State of Bahia. An epidemiological and clinical study in one of the villages has documented autochthonous cases of disease and infected dogs (Cunha et al. 1995). In the present study we have determined by serological test and intradermal reaction the frequency of *L. chagasi* infection in relatives and neighbors of patients with VL who lived in these areas. In addition we characterized the cytokines profile secreted by lymphocytes from subjects with asymptomatic *L. chagasi* infection upon stimulation with leishmania antigen.

MATERIALS AND METHODS

The study was conducted in the villages of Monte Gordo and Barra do Jacuípe that are located at sea level along the Coco road, which links Salvador with coastal towns of the State of Bahia. These villages have a population nearly 6000 inhabitants and 10 to 12 cases of VL have been reported annually since 1989. There are no cases of

+Corresponding author. Fax: +55-71-245.7110

Received 29 April 1996

Accepted 21 August 1996

cutaneous or mucosal leishmaniasis in this area. Participants of the present study included relatives and neighbors of 12 patients who had VL diagnosed between June of 1992 and May of 1993. The patients with VL were diagnosed by finding amastigotes of *L. chagasi* in material obtained from bone marrow or spleen aspiration stained by Giemsa. In order to determine the frequency of infection in relatives and neighbors we evaluated the father, mother, siblings, brother and sister living in the same house of the cases and the neighbors who lived no more than 100 m of the cases of VL. This distance of no more than 100 m was chosen based on evidence of the mean flight range of phlebotomines (Southgate & Oriedo 1967). The patients, parents and neighbors share the same socio-cultural, economic and environmental conditions. The subjects evaluated in this study, 100% of the relatives and 95% of the neighbors of the cases of VL had a clinical history and examination, a serological test to detect antibodies against *L. chagasi* and an intradermal skin test with leishmania antigen. Additionally we characterized the cytokine profile in 11 subjects with asymptomatic infection and compared these findings with those observed in 5 subjects cured of VL and 5 with active VL.

Leishmania antigen, antibody detection and skin test - Leishmania lysate was obtained from promastigotes of a strain isolated from a patient with VL and characterized by monoclonal antibodies and isoenzymes as *L. chagasi* (MHOM Ba 62). Briefly, promastigotes in the stationary phase of growth from cultures were washed three times in PBS, re-suspended and rapidly frozen (-70°C) and thawed (37°C) six times (Reed et al. 1986). After sonication the lysate was centrifuged (12000xg) for 20 min, the supernatant was collected, and protein content was determined by using the method of Lowry et al. (1951).

An ELISA to detect antibodies against leishmania using 1:100 dilution of assayed sera was done as previously described (Badaró et al. 1986c). A serological test was considered positive when the OD was >0.040 which represents the mean plus 3 SD of absorbancies obtained of sera from individuals without exposure to leishmania.

Lymphocyte blastogenesis test and cytokine determination - Peripheral blood mononuclear cells were obtained from heparinized venous blood by centrifugation using lymphocyte separation medium (Bionetics Laboratory Products Kensington, MD) as previously described (Carvalho & Horwitz 1980). After washing in RPMI 1640 (GIBCO, Grand Island, NY) and adjusted to a concentration of 10^6 cells/ml in medium supplemented with 10% pooled human AB serum, triplicate samples

(aliquots 0.2ml) of this cell suspension were cultured in microtiter plates (Limbro Chemical, New Heaven, CT). Cell cultures were incubated (37°C, 5% CO₂) for 5 days with *L. chagasi* antigen (1mg/ml of protein). The cells were then pulsed with 1mCi ³[H]-thymidine for the final 6 hr of culture. The results are expressed as the mean response of ³[H]-thymidine incorporation in triplicate cultures.

For IFN- γ and IL-10 detection, the PBMC were adjusted to 3×10^6 cells/ml and stimulated with 20mg of *L. chagasi* antigen. After 48 hr the cells were centrifuged and the supernatant filtered and stored at -20°C until use. Gamma-interferon and IL-10 levels were determined by ELISA using a sandwich technique (Russo et al. 1993). A standard curve was used to express the results in pg/ml.

Statistical analysis - Data on frequency of positive ELISA and/or positive intradermal skin test in relative and neighbors was performed by χ^2 test with correction of Yates. Data on lymphocyte proliferation and cytokine production were compared by rank sum test. To calculate the risk of infection for *L. chagasi* we used the odds ratio (OR) with 95% confidence intervals. Levels of statistical significance were designated at P<0.05 (two-tailed-test).

RESULTS

The clinical and demographic findings of 12 cases of VL are shown in Table I. The age of the patients ranged from 2 to 46 years with median of 26 years. In five cases the age was equal or lower than 10 years. Seven patients lived in the villages since they were born. The others were living in the villages for at least the last six years. The duration of the illness in such cases ranged from 15 to 90 days. There was a cluster of cases of VL in families since 8 of the 12 patients belonged to three families. Two families with two cases and one family with four cases. ELISA test to detect *L. chagasi* antibodies and intradermal skin with leishmania antigen test were performed in relatives and neighbors (n=135) of the VL cases. All subjects tested had no present complain related to VL and they denied previous diagnosis of VL. Evidence of *L. chagasi* infection determined by a positive ELISA or intradermal skin test (Table II) was documented in 18 of 40 relatives (45%) and in 26 of 95 (27%) of the neighbors of the VL cases (P>0.07; OR=2.17: 1.0 - 4.7). The age distribution of subjects infected by *L. chagasi* followed the age distribution of the sample (data not shown). In all age groups there was a higher number of cases with positive skin test than that subjects with positive serology (data not shown). *L. chagasi* infection, determined by positive serological test and or positive skin test, was observed in 31 (39.7%) of the

TABLE I
Clinical features of visceral leishmaniasis in a recent area of *Leishmania chagasi* transmission

Patients initials	Age/Sex	No. of years living in the area	Hepatomegaly	Splenomegaly	ELISA (OD)	Parasite isolation
JPC	2/F	2	Yes	Yes	0,194	Yes
DSB	4/F	4	Yes	Yes	0,50	Yes
RCS	5/F	5	Yes	Yes	0,208	Yes
LSC	6/F	6	Yes	Yes	0,67	Yes
LSC	10/F	6	Yes	Yes	0,46	Yes
ANS	19/M	19	Yes	Yes	0,154	Yes
IBS	32/F	30	Yes	Yes	0,164	Yes
JLF	39/M	8	No	No	0,180	No
MDS	39/M	39	Yes	Yes	0,137	Yes
RDS	41/M	41	No	Yes	0,49	Yes
ACS	55/M	6	No	No	0,256	Yes
LES	58/F	10	Yes	Yes	0,96	Yes

TABLE II

Frequency of relatives and neighbors of visceral leishmaniasis cases with evidence of antibodies or positive skin test to *Leishmania chagasi*

Population	Evidence of infectivity ^a		Total
	Positive (%)	Negative (%)	
Relatives ^b	18 (45)	22 (55)	40
Neighbors	26 (27)	69 (73)	95
Total	44	91	135

a: serology and/or intradermal skin test positive; b: parents, children and brothers.

78 subjects older than 13 years and in 13 (22.8%) of 57 children aged 13 years or less (P=0.04; OR=2.23:1.03- 4.8). None of the population had splenomegaly and none developed VL after one year and half of follow-up.

To characterize the T cell response in these subjects with asymptomatic *L. chagasi* infection the lymphocyte proliferative response and the cytokine profile produced after *in vitro* stimulation with *L. chagasi* were evaluated (Table III). Eleven subjects, mean age of 28±20, participated of the immunological study being five with negative ELISA test and positive intradermal skin test, five with positive ELISA and negative intradermal reaction and one case with both tests positive. As control T cell response was performed in five subjects cured of VL (mean age 29±11) and five VL patients (mean age 16±6).

The ³[H]-thymidine uptake of lymphocyte cultures ranged from 2205 to 54373 cpm with mean and SD of 30966±15111. The IFN-γ production ranged from <20 to 1234 pg/ml with mean of 601±391 pg/ml. The magnitude of lymphocyte

TABLE III
Lymphoproliferative response and cytokine profile of subjects with asymptomatic *Leishmania chagasi* infection

Age/Sex	ELISA	Skin test	3H-Thymidine		Cytokine levels (pg/ml)	
			Media	<i>L. chagasi</i>	IFN-γ	IL-10
7/F	-	+	864±315	41380±1265	670	25
11/F	-	+	1119±332	45735±939	410	48
13/F	-	+	2596±126	54373±4800	851	26
16/M	-	+	297±85	21631±1842	1234	47
19/M	-	+	567±115	44382±2765	785	18
19/M	+	-	861±110	22141±1353	300	19
24/F	+	-	367±111	15857±1589	1220	25
35/F	+	-	940±103	32921±9633	237	19
43/M	+	-	478±130	31370±3280	480	24
55/M	+	-	684±75	2205±638	<20	26
72/F	+	+	731±182	28654±1285	410	21
Active visceral leishmaniasis (n=5)			743±587	926±334	15±8	387±118
Cured visceral leishmaniasis (n=5)			862±316	37286±12345	547±263	27±11

proliferation and cytokine production was not different when we compared subjects with positive skin test with those with negative skin test ($p > 0.05$). Of the eleven subjects evaluated with asymptomatic infection six were neighbors and five were relatives from patients with VL. The lymphoproliferative response and cytokine production in the neighbors was not different when compared with the relatives of the VL cases ($P > 0.05$). The IFN- γ production by subjects with asymptomatic *L. chagasi* infection was higher ($P < 0.05$) than that observed in patients with active VL (15 ± 8 pg/ml) and similar ($P > 0.05$) to that observed in subjects cured of VL (547 ± 263 pg/ml). In contrast with the high production of IFN- γ observed in the subjects with asymptomatic *L. chagasi* infection there was a very low production of IL-10. The IL-10 levels in subjects with asymptomatic *L. chagasi* infection (27 ± 11 pg/ml) were lower ($p < 0.01$) than those observed in patients with active VL (387 ± 118 pg/ml).

DISCUSSION

In the present study we showed that although cases of VL tend to occur in the same family, the frequency of the *L. chagasi* infection in relatives in the same house of patients with VL was similar to that observed in neighbors of this same patients. Additionally, the immunological studies in subjects with asymptomatic *L. chagasi* infection showed that upon leishmania antigen stimulation, these individuals produce high amount of IFN- γ and very little IL-10.

Several factors have been associated with predisposition of development of VL. Among these factors are: low age, malnourishment and inability of lymphocytes to proliferate and to produce IFN- γ upon exposition to *L. chagasi* antigen (Badaró et al. 1986a, Carvalho et al. 1992). In contrast, very little information is known regarding factors involved in acquisition of *L. chagasi* infection and why the majority of the population infected by *L. chagasi* is able to control the progression of the infection to the disease.

Epidemiological studies performed in areas of *L. chagasi* transmission have shown that VL occurs predominantly in children (Guedes et al. 1974, Wijers & Killu 1984, Navin et al. 1985, Jahn et al. 1986, Badaró et al. 1986b, Corredor et al. 1989, Beer et al. 1991, Evans et al. 1992). In the present study performed in a recent area of *L. chagasi* transmission we observed that the disease was more frequent in adults than in children and that the adult population was also more infected than the children ($P = 0.04$). Similar findings have been observed by others in outbreaks in areas previously free of the disease (Zijlstra et al. 1991, Perea et al. 1991).

Since the majority of our patients with VL were born in the villages or have lived there for at least the last six years, a period much higher than the illness duration, and since transmission of *L. chagasi* is active (dogs have been found infected and *Lutzomyia longipalpis* are frequent in the area) the autochthony of all cases is highly likely. VL in adult is common in Africa where wild or domestic animals have not been identified as reservoir of *L. donovani* (WHO 1990). In the Americas transmission of *L. chagasi* infection is predominantly peridomiciliar (Corredor et al. 1989, WHO 1990) and the dog is the more frequent reservoir of *L. chagasi* infection (Alencar et al. 1975). However several studies have failed to determine that killing of dogs decrease seroconversion and the incidence of VL indicating that others reservoirs may be involved in *L. chagasi* transmission (Deane & Deane 1962, Lainson 1983, Ward 1985, Evans et al. 1985). The findings from this study showing that VL and asymptomatic infection occur more frequently in the adult population than in children, suggests that in recent areas of *L. chagasi* transmission, a wild reservoir may have an important role. After years of transmission a high percentage of the adult population have been already infected and have control or had acquired the disease. Children then become the major target of the leishmania and malnourishment, low age and concomitant infections may now contribute to the increase incidence of the disease in childhood (Badaró et al. 1986b, Cerf et al. 1987, Evans et al. 1992).

Genetic susceptibility to leishmania infection have been reported in experimental animals (Bradley 1977, Blackwell 1982) and cases of VL have been documented in the same family. However, families not related living closely have different incidence of disease and infection (Thakur 1984, Evans et al. 1985, Costa et al. 1986, Badaró et al. 1986b, Dhiman & Sen 1991) and evidence of a genetic predisposition for development of the human disease is lacking. In this study 8 of the 12 cases of VL that were diagnosed in a period of 12 months occurred in three families, indicating that genetic studies need to be performed. In spite of the cluster of cases of VL, there was no evidence that infection by *L. chagasi* determined by serodiagnosis or skin test was significantly more frequent in relatives than in neighbors of the patients with VL.

Depression of Th1 response characterized by absence of lymphoproliferative response, IL-2 and IFN- γ production and secretion of cytokines of Th2 type such IL-4, IL-5 and IL-10 are documented in VL (Carvalho et al. 1989, Karp et al. 1993, Ghalib et al. 1993, Carvalho et al. 1994). Previously we have observed that children recently exposed to *L.*

chagasi and able to produce IFN- γ are able to control the infection or even a sub-clinical disease preventing the development of VL (Carvalho et al. 1992). This present study confirms that IFN- γ production is a marker of individuals with asymptomatic *L. chagasi* infection. Additionally we extend the characterization of the immune responses showing that IL-10 production is low in such subjects. Marked expression of mRNA for IL-10 is observed in VL (Karp et al. 1993, Carvalho et al. 1994) and the role of this cytokine in down regulate T cell responses in these patients have been documented (Ghalib et al. 1993, Carvalho et al. 1994). The determination of high IFN- γ and low IL-10 production in subjects with asymptomatic *L. chagasi* infection indicate that balance between the production of these cytokines may be an important key in determine whether subjects will or not develop disease.

IFN- γ production after leishmania antigen stimulation was documented in all subjects but one and low production of IL-10 was observed in the asymptomatic subjects. These observations were similar to those observed in cured subjects and different from patients with VL that had high IL-10 and absence of IFN- γ production. The only asymptomatic subject who did not produce IFN- γ did not produce IL-10. This subject remained asymptomatic up one year and half of seroconversion and was only once immunologically evaluated. It is possible that if subsequent evaluations were performed, this patient became a responder as has been previously observed (Carvalho et al. 1992).

ACKNOWLEDGMENTS

To the Director of the Health Post of Monte Gordo and the other staff members specially Mr João Silva Araújo for their cooperation. To Mrs Elbe Silva and Mr Jackson Lemos for preparation of the manuscript.

REFERENCES

- Alencar JE, Almeida YM, Silva ZF, Paiva AS, Fonseca MF 1975. Aspectos atuais do calazar no Ceará. *Rev Bras Malariol Doença Trop* 25/26: 27-48.
- Badaró R, Jones TC, Carvalho EM, Sampaio D, Reed SG, Barral A, Teixeira R, Johnson WD 1986a. New perspective on a subclinical form of visceral leishmaniasis. *J Infect Dis* 154: 1003-1011.
- Badaró R, Jones TC, Lorenço R, Cerf BJ, Sampaio D, Carvalho EM, Rocha H, Teixeira R, Johnson WD 1986b. A prospective study of visceral leishmaniasis in an endemic area of Brazil. *J Infect Dis* 154: 639-649.
- Badaró R, Reed SG, Barral A, Orge G, Jones TC 1986c. Evaluation of the micro enzyme linked immunosorbent assay (ELISA) for antibodies in American visceral leishmaniasis antigen selection for detection of infection specific responses. *Am J Trop Med Hyg* 35: 79-85.
- Beer P, de Harit AEC, Deng LL, Semião-Santos SJ, Chantal B, Grootheest M 1991. A killing disease epidemic among displaced sudanese population identified as visceral leishmaniasis. *Am J Trop Med Hyg* 44: 283-289.
- Blackwell JM 1982. Genetic control of recovery from visceral leishmaniasis. *Trans R Soc Trop Med Hyg* 76: 147-154.
- Bradley DJ 1977. Regulation of the leishmania populations within the host. II. Genetic control of acute susceptibility of mice to *L. donovani* infection. *Clin Exp Immunol* 30: 130-140.
- Cabello PH, Lima AMVM, Azevedo ES, Krieger H 1995. Familial aggregation of *Leishmania chagasi* infection in northeastern Brazil. *Am J Trop Med Hyg* 52: 364-365.
- Carvalho EM, Bacellar O, Barral A, Badaró R, Johnson WD 1989. Antigen-specific immunosuppression in visceral leishmaniasis is cell mediated. *J Clin Invest* 83: 860-864.
- Carvalho EM, Bacellar O, Brownell CE, Coffman RL, Reed SG 1994. Restoration of IFN-gamma production and lymphocyte proliferation in visceral leishmaniasis. *J Immunol* 152: 5949-5956.
- Carvalho EM, Barral A, Pedral-Sampaio D, Barral-Netto M, Badaró R, Rocha H, Johnson WD 1992. Immunologic markers of clinical evolution in children recently infected with *Leishmania donovani chagasi*. *J Infect Dis* 165: 535-540.
- Carvalho EM, Horwitz DA 1980. Characterization of a non-T, non-B human blood lymphocyte that mediates the enhancing effects of immune complexes on lymphocyte blastogenesis. *J Immunol* 124: 1656-1661.
- Cerf BJ, Jones TC, Badaró R, Sampaio D, Carvalho EM, Rocha H, Teixeira R, Johnson WD 1987. Malnutrition as a risk factor for severe visceral leishmaniasis. *J Infect Dis* 156: 1030-1032.
- Corredor A, Gallego JF, Tesh RB, Morales A, Carrasquilla CF, Young DG, Kreutzen RD, Boshell J, Palan MT, Carceres E, Palaez D 1989. Epidemiology of visceral leishmaniasis in Colombia. *Am J Trop Med Hyg* 40: 480-486.
- Costa JML, Osaho NK, Vale KC, Lago EL, França F, Vaxenat JA, Marsden PD 1986. Ocorrência familiar da leishmaniose tegumentar americana em uma região endêmica, Corte de Pedra, Bahia. *Rev Soc Bras Med Trop* 19: 197-198.
- Cunha S, Freire M, Eulálio C, Cristovão J, Netto E, Johnson WD, Reed SF, Badaró R 1995. Visceral leishmaniasis in a new ecological niche near a major metropolitan area of Brazil. *Trans R Soc Trop Med Hyg* 89: 155-158.
- Deane LM, Deane MP 1962. Visceral leishmaniasis in Brazil: Geographic distribution and transmission. *Rev Inst Med Trop São Paulo* 4: 198-212.
- Dhiman RC, Sen AB 1991. Epidemiology of kala-azar in rural Bihar (India) using village as a component unit of study. *Indian J Med Res* 93: 155-160.
- Evans T, Reis M de F, de Alencar JE, Naidu TG, de Jesus JA, Mc Auliffe JF, Pearson RD. 1985. American visceral leishmaniasis (kala-azar). *West J Med*

- 142: 777-781.
- Evans TG, Teixeira MJ, McAuliffe IT, Vasconcelos IAB, Vasconcelos AW, Sousa AQ, Lima JWO, Pearson RD 1992. Epidemiology of visceral leishmaniasis in northeast Brazil. *J Infect Dis* 166: 1124-1132.
- Ghalib HW, Piuvezan MR, Skeiky YAW, Siddig M, Hashmi FA, El-Hassam AM, Russo DM, Reed SG 1993. Interleukin-10 production correlates with pathology in human *L. donovani* infections. *J Clin Invest* 92: 324-329.
- Guedes GE, Maroja A, Chaves E, Estélio J, Cunha MJ, Arcoverde S 1974. Calazar no litoral do estado da Paraíba, Brasil. Encontro de 70 casos humanos e 16 caninos. *Rev Inst Med Trop São Paulo* 16: 265-269.
- Jahn A, Lelmett JM, Diesfeld HJ 1986. Seroepidemiological study of kala-azar in Beringo District, Kenya. *J Trop Med Hyg* 89: 91-94.
- Karp CL, El-Safi SH, Wynn TA, Satt MMH, Kardogani AM, Hashmi FA, Hag-Ali M, Neva FA, Nutman TB, Sacks DL 1993. *In vivo* cytokine profiles in patients with kala-azar. *J Clin Invest* 91: 1644-1648.
- Lainson R 1983. The American leishmaniasis: some observations on their ecology and epidemiology. *Trans R Soc Trop Med Hyg* 77: 569-596.
- Lowry DH, Rosembrongh NG, Farr AL, Randall RG 1951. Protein measurement with the folin phenol reagent. *J Biol Chem* 193: 265-275.
- Navin TR, Sierra M, Custodio R, Steurer F, Porter CH, Ruebush TK 1985. Epidemiologic study of visceral leishmaniasis in Honduras, 1975-1983. *Am J Trop Med Hyg* 34: 1069-1075.
- Perea WA, Ancelle T, Moren A, Nagelkerke, Sondorp E 1991. Visceral leishmaniasis in southern Sudan. *Trans R Soc Trop Med Hyg* 85: 48-53.
- Reed SG, Badaró R, Masur H, Carvalho EM, Lorenço R, Lisboa A, Teixeira R, Johnson WD, Jones TC 1986. Selection of a skin test antigen for american visceral leishmaniasis. *Am J Trop Med Hyg* 35: 79-85.
- Russo DM, Jardim A, Carvalho EM, Sleath PR, Armitage RJ, Olafson RW, Reed SG 1993. Mapping human T cell epitopes in leishmania gp63. Identification of cross-reactive and species-specific epitopes. *J Immunol* 150: 932-939.
- Southgate BA, Oriedo BVE 1962. Studies in the epidemiology of East African leishmaniasis. I. The circumstantial epidemiology of Kala-azar in the Kitui District of Kenya. *Trans R Soc Trop Med Hyg* 56: 30-47.
- Thakur CP 1984. Epidemiological, clinical and therapeutic features of Bihar kala-azar (including post kala-azar dermal leishmaniasis). *Trans R Soc Trop Med Hyg* 78: 391-398.
- Ward RD 1985. Vector biology and control leishmaniasis, p.199-212. In KP Chang, RS Bray (eds) *Leishmaniasis*. Elsevier, New York.
- Wijers DJB, Killu G 1984. Studies on the vector of kala-azar in Kenya. VIII. The outbreak in Machakos District; epidemiological features and possible way of control. *An Trop Med Parasitol* 78: 597-604.
- WHO - World Health Organization 1990. *Control of leishmaniasis*. WHO Tech Rep Ser 793: 54
- Zijlstra EE, Ali MS, El-Hassan AM, El-Toum IA, Satti M, Ghalib HW, Sondorp E, Winkler A 1991. Kala-azar in displaced people from southern Sudan: epidemiological, clinical and therapeutic findings. *Trans R Soc Trop Med Hyg* 85: 365-369.