Inherited neurometabolic diseases and the importance of imaging-based classification systems

Nina Ventura¹

Inherited neurometabolic disorders (INMDs) constitute a heterogeneous group of diseases that includes the so-called inborn errors of metabolism, leukodystrophies, mitochondrial diseases, and deposition diseases⁽¹⁾. Although INMDs are rare individually, they collectively account for a significant proportion of neurological diseases in children^(1,2). The clinical and imaging findings of INMDs can be nonspecific and can overlap, making the diagnosis challenging. However, because there are treatments available for some INMDs, early diagnosis is fundamental in order to minimize or prevent neurological damage⁽¹⁾. Therefore, it is essential that neuroradiologists are able to recognize the imaging findings that can narrow the differential diagnosis.

Magnetic resonance imaging is the modality of choice for the evaluation of various diseases of the central nervous system⁽³⁻⁷⁾, including INMDs, being crucial for the localization, identification, and characterization of alterations. It is still the main method employed for monitoring patients with INMDs, either to evaluate the natural progression of the disease or to assess the response to treatment, as well as being used in order to screen family members for genetic counseling⁽⁸⁻¹⁰⁾.

Over the years, a number of classifications based on magnetic resonance imaging findings have been proposed, in attempts to facilitate the approach to INMDs and demystify the topic. Recent articles have proposed divisions by disease groups, such as leukodystrophies or mitochondrial diseases^(10,11), or an epidemiological division, with separate categories for disorders that occur only in adults or children^(8,12). Some authors have also shown that specific imaging findings, such as the pattern of involvement of the white matter and basal ganglia, as well as the presence or absence of cysts and a T2-weighted signal in the corpus callosum, are quite useful for guiding the diagnosis^(9,13).

Radiol Bras. 2022 Mai/Jun;55(3):VII-VIII

The article authored by Pedri et al.⁽¹⁴⁾, entitled "Classification of inherited neurometabolic disorders based on radiological aspects: pictorial essay" and published in the previous issue of **Radiologia Brasileira**, proposes a new classification system that encompasses all INMDs, of any epidemiology or subtype. That system is easily applied in daily practice because it is based on the patterns seen on diagnostic imaging. The authors identified 10 distinct patterns: macrocrania; cysts; distribution of white matter involvement; contrast enhancement; calcifications; basal ganglia involvement; restricted diffusion; vascular abnormalities; cranial nerve involvement; and the metabolic profile on spectroscopy. The article presents practical divisions and could be an excellent resource when we encounter any suspicion of an INMD.

Despite the fact that INMD is a complex subject that is avoided by many radiologists, it is inevitable that we will come across some cases over the course of our careers. Knowledge of the key findings and imaging patterns can facilitate the investigation and make the diagnosis of INMDs less challenging, as can reliable, concise sources for consultation with practical guides, such as that provided by Pedri et al.⁽¹⁴⁾.

REFERENCES

- Biswas A, Malhotra M, Mankad K, et al. Clinico-radiological phenotyping and diagnostic pathways in childhood neurometabolic disorders-a practical introductory guide. Transl Pediatr. 2021;10:1201–30.
- Aksoy DÖ, Alkan A. Neurometabolic diseases in children: magnetic resonance imaging and magnetic resonance spectroscopy features. Curr Med Imaging Rev. 2019;15:255–68.
- Pereira RG, Niemeyer B, Pereira TRGC, et al. Magnetic resonance imaging evaluation of spinal cord lesions: what can we find? – Part 1. Neoplastic, vascular, metabolic, and traumatic injuries. Radiol Bras. 2021;54:406–11.
- 4. Pereira RG, Niemeyer B, Pereira TRGC, et al. Magnetic resonance imaging evaluation of spinal cord lesions: what can we find? Part 2. Inflammatory and infectious injuries. Radiol Bras. 2021;54:412–7.
- Dalaqua M, Nascimento FBP, Miura LK, et al. Magnetic resonance imaging of the cranial nerves in congenital, traumatic, and vascular diseases: a pictorial essay. Radiol Bras. 2021;54:185–92.
- Pereira RG, Niemeyer B, Hollanda RTL, et al. Non-neoplastic intracranial cystic lesions: not everything is an arachnoid cyst. Radiol Bras. 2021;54: 49–55.
- Santana LM, Valadares EJA, Rosa-Júnior M. Differential diagnosis of temporal lobe lesions with hyperintense signal on T2-weighted and FLAIR sequences: pictorial essay. Radiol Bras. 2020;53:129–36.
- Resende LL, Paiva ARB, Kok F, et al. Adult leukodystrophies: a step-by-step diagnostic approach. Radiographics. 2019;39:153–68.

^{1.} Instituto Estadual do Cérebro Paulo Niemeyer, Rio de Janeiro, RJ, Brazil; Universidade Federal do Rio de Janeiro (UFRJ), Rio de Janeiro, RJ, Brazil; Hospital Samaritano, Grupo Fleury, Rio de Janeiro, RJ, Brazil. E-mail: niventuraa@ gmail.com.

https://orcid.org/0000-0003-2364-1612.

- Roosendaal SD, van de Brug T, Alves CAPF, et al. Imaging patterns characterizing mitochondrial leukodystrophies. AJNR Am J Neuroradiol. 2021; 42:1334-40.
- van der Knaap MS, Bugiani M. Leukodystrophies: a proposed classification system based on pathological changes and pathogenetic mechanisms. Acta Neuropathol. 2017;134:351–82.
- 11. Gonçalves FG, Alves CAPF, Heuer B, et al. Primary mitochondrial disorders of the pediatric central nervous system: neuroimaging findings. Radiographics. 2020;40:2042-67.
- (cc)) BY

- Cheon JE, Kim IO, Hwang YS, et al. Leukodystrophy in children: a pictorial review of MR imaging features. Radiographics. 2002;22:461–76.
- Paprocka J, Machnikowska-Sokołowska M, Gruszczyńska K, et al. Neuroimaging of basal ganglia in neurometabolic diseases in children. Brain Sci. 2020;10:849.
- 14. Pedri AF, Guedes MS, Castro CC. Classificação das doenças neurometabólicas hereditárias baseada em aspectos radiológicos: ensaio iconográfico. Radiol Bras. 2022;55:113–9.